

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 June 2007 (28.06.2007)

PCT

(10) International Publication Number
WO 2007/072997 A1

(51) International Patent Classification:
C07K 14/47 (2006.01) A61K 38/16 (2006.01)

(21) International Application Number:
PCT/JP2006/326176

(22) International Filing Date:
21 December 2006 (21.12.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2005-370388 22 December 2005 (22.12.2005) JP
2006-275843 6 October 2006 (06.10.2006) JP

(71) Applicant (for all designated States except US): TAKEDA
PHARMACEUTICAL COMPANY LIMITED [JP/JP];
1-1, Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka,
5410045 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ASAMI, Taiji
[JP/JP]; c/o TAKEDA PHARMACEUTICAL COMPANY
LIMITED, 10, Wadai, Tsukuba-shi, Ibaraki, 3004293
(JP). NISHIZAWA, Naoki [JP/JP]; c/o TAKEDA PHAR-
MACEUTICAL COMPANY LIMITED, 10, Wadai,
Tsukuba-shi, Ibaraki, 3004293 (JP).

(74) Agents: KOBAYASHI, Hiroshi et al.; c/o ABE, IKUBO
& KATAYAMA, Fukuoka Bldg. 9th Fl. 8-7, Yaesu
2-chome, Chuo-ku, Tokyo, 1040028 (JP).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS,
JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS,
LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY,
MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS,
RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METASTIN DERIVATIVES AND USE THEREOF

(57) Abstract: The present invention provides stable metastin derivatives having excellent biological activities (a cancer metasta-
sis suppressing activity, a cancer growth suppressing activity, a gonadotrophic hormone secretion stimulating activity, sex hormone
secretion stimulating activity, etc.). By substituting the constituent amino acids of metastin with specific amino acids in the metastin
derivative of the present invention, blood stability, solubility, etc. are more improved, gelation tendency is reduced, pharmacokinet-
ics are also improved, and an excellent cancer metastasis suppressing activity or a cancer growth suppressing activity is exhibited.
Furthermore, the metastin derivative of the present invention has the effects of suppressing gonadotropic hormone secretion, sup-
pressing sex hormone secretion, etc.

WO 2007/072997 A1

DESCRIPTION

METASTIN DERIVATIVES AND USE THEREOF

5

TECHNICAL FIELD

The present invention relates to metastin derivatives and use thereof.

BACKGROUND ART

Human-derived metastin (also called KiSS-1 peptide) (WO 00/24890) and
10 mouse or rat-derived metastin (WO 01/75104) are known. Sustained release
preparations containing metastin are also known (WO 02/85399).

Reportedly, metastin has an effect of suppressing cancer metastasis and is
thus effective for preventing or treating cancers (for example, lung cancer, gastric
cancer, liver cancer, pancreatic cancer, colorectal cancer, rectal cancer, colonic cancer,
15 prostate cancer, ovarian cancer, cervical cancer, breast cancer, renal cancer, bladder
cancer, brain tumor, etc.); metastin also has an effect of controlling the pancreatic
function and is effective for preventing or treating pancreatic diseases (e.g., acute or
chronic pancreatitis, pancreatic cancer, etc.); and metastin further has an effect of
controlling placental function and is effective for preventing or treating
20 choriocarcinoma, hydatid mole, invasive mole, miscarriage, fetal hypoplasia,
abnormal glucose metabolism, abnormal lipid metabolism or abnormal delivery (WO
00/24890, WO 01/75104 and WO 02/85399).

DISCLOSURE OF THE INVENTION


25 The present invention aims at providing stable metastin derivatives having
excellent biological activities (a cancer metastasis suppressing activity, a cancer
growth suppressing activity, a gonadotrophic hormone secretion stimulating activity,
sex hormone secretion stimulating activity, etc.).

The present inventors have made extensive studies to solve the foregoing
30 problems and as a result, have found that by substituting the constituent amino acids
of metastin with specific amino acids, unexpectedly blood stability, solubility, etc.
are more improved, gelation tendency is reduced, pharmacokinetics are also
improved, and an excellent cancer metastasis suppressing activity or a cancer growth
suppressing activity is exhibited. The present inventors have further found that
35 unexpectedly these metastin derivatives have an effect of suppressing the

gonadotropic hormone secretion, an effect of suppressing the sex hormone secretion, etc., which are totally different from the effects known so far. Based on these findings, the present inventors have continued further investigations and come to accomplish the present invention.

5 That is, the present invention provides the following features, and so on.

(1) A metastatin derivative represented by formula:



10 (wherein:

XX0 represents formyl, a C₁₋₂₀ alkanoyl, cyclopropanecarbonyl, 6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl, 6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl, 3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl, 15 acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl, 6-aminocaproyl, 6-acetylaminocaproyl, 4-[bis-(2-pyridylmethyl)aminomethyl]benzoyl or 4-ureidobenzoyl;

XX2 represents Tyr, D-Tyr, D-Ala, D-Leu, D-Phe, D-Lys, D-Trp or a chemical bond;

20 XX3 represents:

i) an amino acid selected from Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Ser, Thr, Trp, Tyr and Val wherein the α -amino group may optionally be methylated;

25 ii) a cyclic amino acid selected from Pro, Aze(2), Aze(3), Pic(2), Pic(3), Hyp, Thz, Abz(2), Abz(3), Pzc(2), Pro(4NH₂), Hyp(Bzl), cisHyp, Pro(4F) and lzc;

iii) an amino acid selected from D-Dap, D-Pya(4), DL-Ala(Pip), Orn, Aib and Tyr(PO₃H₂); or,

iv) a chemical bond;

30 XX4 represents Asn, 2-amino-3-ureidopropionic acid, N ^{β} -formyl- β -diaminopropionic acid, N ^{β} -acetyl- β -diaminopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine, 2,4-diaminobutanoic acid, 2,3-diaminopropionic acid, His, Gln, Gly, Arg, Cit, Nva, D-Asn or a chemical bond;

35 XX5 represents Ser, Thr, Val, NMeSer, Gly, Ala, Hyp, D-Ala, D-Thr, D-Pro or a chemical bond;

XX6 represents Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4),
threo-Ser(3Phenyl), erythro-Ser(3Phenyl), or optionally substituted phenylalanine ;

AzaGly represents azaglycine;

XX8 represents Leu, Nva, Val or Ala(cPr);

5 XX9 represents optionally substituted arginine, optionally substituted lysine
or optionally substituted ornithine; and,

XX10 represents 2-naphthylalanine, 2-thienylalanine, tyrosine, optionally
substituted phenylalanine or optionally substituted tryptophan); or a salt thereof.

(2) A metastatin derivative-represented by formula:

10

XX0-XX2-XX3-XX4-XX5-XX6-AzaGly-XX8-XX9-XX10-NH₂

(wherein:

XX0 represents formyl, a C₁₋₂₀ alkanoyl, cyclopropanecarbonyl,
15 6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl,
6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl,
3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl,
acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl,
6-aminocaproyl, 6-acetylaminocaproyl,
20 4-[bis-(2-pyridylmethyl)aminomethyl]benzoyl or 4-ureidobenzoyl;

XX2 represents Tyr, D-Tyr, D-Ala, D-Leu, D-Phe, D-Lys, D-Trp or a
chemical bond;

XX3 represents D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-Trp, D-Tyr,
D-Pya(4), D-NMeAla, D-NMePhe, Aze(2), Aze(3), Pic(2), Pic(3), Hyp, Thz, Gly,
25 Aib, Abz(2), Abz(3), Sar, Izc, Léu, Lys, Glu, Thr, Trp, Ser, Ala, NMeAla, β-alanine,
DL-Ala(Pip), Pzc(2), Orn, His(3Me), Tyr(PO₃H₂), Pro(4NH₂), Hyp(Bzl), cisHyp,
Pro(4F) or a chemical bond;

XX4 represents Asn, 2-amino-3-ureidopropionic acid,
N^β-formyl-β-diaminopropionic acid, N^β-acetyl-β-diaminopropionic acid,
30 N^ω-pentylasparagine, N^ω-cyclopropylasparagine, N^ω-benzylasparagine,
2,4-diaminobutanoic acid, 2,3-diaminopropionic acid, His, Gln, Gly, Arg, Cit, Nva,
D-Asn or a chemical bond;

XX5 represents Ser, Thr, Val, NMeSer, Gly, Ala, Hyp, D-Ala, D-Thr, D-Pro
or a chemical bond;

35 XX6 represents Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), Phe(2F),

Phe(3F), Phe(4F), Phe(4Cl), α MePhe, Phe(2Me), Phe(3Me), Phe(4Me),
threo-Ser(3Phenyl), erythro-Ser(3Phenyl) or D-Phe;

AzaGly represents azaglycine;

XX8 represents Leu, Nva, Val or Ala(cPr);

5 XX9 represents Arg, Orn, Arg(Me), D-Arg or Arg(asymMe₂); and,

XX10 represents Phe, Trp, 2-naphthylalanine, 2-thienylalanine, tyrosine or
4-fluorophenylalanine); or a salt thereof.

(3) A metastatin derivative represented by formula:

10 XX0-XX2-XX3-XX4-XX5-XX6-AzaGly-XX8-XX9-XX10-NH₂

(wherein:

XX0 represents formyl, a C₁₋₂₀ alkanoyl, cyclopropanecarbonyl,
6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl,
15 6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl,
3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl,
acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl or
6-aminocaproyl;

XX2 represents Tyr, D-Tyr, D-Ala, D-Leu, D-Phe, D-Lys, D-Trp or a
20 chemical bond;

XX3 represents D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-NMeAla,
D-NMePhe, Aze(2), Pic(2), Pic(3), Hyp, Thz, NMeAla, Gly, Aib, Abz(2), Abz(3),
Sar, Leu, Lys, Glu, β -alanine, Pzc(2), Orn, His(3Me), Tyr(PO₃H₂), Pro(4NH₂) or
Hyp(Bzl);

25 XX4 represents Asn, 2-amino-3-ureidopropionic acid,
N ^{β} -formyl- β -diaminopropionic acid, N ^{β} -acetyl- β -diaminopropionic acid,
N ^{α} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine,
2,4-diaminobutanoic acid, His, Gln, Cit or D-Asn;

XX5 represents Ser, Thr, Val, NMeSer, Gly, Ala, Hyp, D-Ala or D-Thr;

30 XX6 represents Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), Phe(2F),
Phe(3F), Phe(4F), Phe(4Cl) or D-Phe;

AzaGly represents azaglycine;


XX8 represents Leu, Nva or Val;

XX9 represents Arg, Orn, Arg(Me) or Arg(asymMe₂); and,

35 XX10 represents Phe, Trp, 2-naphthylalanine, 2-thienylalanine, tyrosine or

4-fluorophenylalanine); or a salt thereof.

(4) A metastatin derivative represented by formula:



5

(wherein:

XX0 represents formyl, a C₁₋₁₂ alkanoyl, cyclopropanecarbonyl,

6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl,

6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl,

10 3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl, acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl or 6-aminocaproyl;

XX2 represents Tyr, D-Tyr, D-Ala, D-Leu, D-Phe, D-Lys, D-Trp or a chemical bond;

15 XX3 represents D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-NMeAla, D-NMePhe, Aze(2), Pic(2), Pic(3), Hyp, Thz, NMeAla, Gly, Aib, Abz(2), Abz(3), Sar, Leu, Lys, Glu, β -alanine, Pzc(2), Orn, His(3Me), Tyr(PO₃H₂), Pro(4NH₂) or Hyp(Bzl);

XX4 represents Asn, 2-amino-3-ureidopropionic acid,

20 N ^{β} -formyldiaminopropionic acid, N ^{β} -acetyldiaminopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine or 2,4-diaminobutanoic acid;

XX5 represents Ser, Thr or Val;

XX6 represents Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), Phe(2F),

25 Phe(3F), Phe(4F) or Phe(4Cl);

AzaGly represents azaglycine;

XX8 represents Leu, Nva or Val;

XX9 represents Arg, Orn, Arg(Me) or Arg(asymMe₂); and,

XX10 represents Phe, Trp, 2-naphthylalanine, 2-thienylalanine, tyrosine or

30 4-fluorophenylalanine); or a salt thereof.


(5) The metastatin derivative or a salt thereof according to (1), wherein:

XX0 represents a C₁₋₁₂ alkanoyl, 6-aminocaproyl, 6-acetylaminocaproyl, glycoloyl, 4-[bis-(2-pyridylmethyl)aminomethyl]benzoyl, 4-ureidobenzoyl, 3-(4-hydroxyphenyl)propionyl or pyroglutamyl;

35 XX2 represents D-Tyr, Tyr or a chemical bond;

- XX3 represents D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-Trp, D-Tyr, D-Pya(4) D-NMeAla, D-NMePhe, Aze(2), Aze(3), Pic(2), Pic(3), Hyp, Thz, Gly, Aib, Abz(2), Sar, Izc, Leu, Lys, Glu, Thr, Trp, Ser, Ala, NMeAla, β -alanine, DL-Ala(Pip), Pzc(2), Orn, His(3Me), Tyr(PO_3H_2), Pro(4NH_2), Hyp(Bzl), cisHyp, Pro(4F) or a chemical bond;
- XX4 represents Asn, 2-amino-3-ureidopropionic acid, N^ω -pentylasparagine, N^ω -cyclopropylasparagine, N^ω -benzylasparagine, 2,4-diaminobutanoic acid, 2,3-diaminopropionic acid, His, Gln, Gly, Arg, Cit, Nva, D-Asn or a chemical bond;
- XX5 represents Thr, NMeSer, Gly, Ala, Hyp, D-Ala, D-Thr, D-Pro or a chemical bond;
- XX6 represents Phe, Phe(2F), Phe(3F), Phe(4F), Phe(4Cl), α MePhe, Phe(2Me), Phe(3Me), Phe(4Me), threo-Ser(3Phenyl), erythro-Ser(3Phenyl) or D-Phe;
- XX8 represents Leu or Ala(cPr);
- XX9 represents Arg, Arg(Me) or D-Arg; and,
- XX10 represents Trp.

(6) The metastatin derivative according to (1), which is represented by formula:



(wherein:

- XX0 represents formyl, a C_{1-6} alkanoyl or glycoloyl;
- XX2 represents D-Tyr or a chemical bond;
- XX3 represents Aze(2), Hyp, Gly, Aib, Leu, Lys, Glu, His(3Me), Tyr(PO_3H_2), Hyp(Bzl), cisHyp or Pro(4F);
- XX4 represents Asn or 2-amino-3-ureidopropionic acid;
- XX5 represents Ser, Thr or Ala;
- XX6 represents Phe, Cha, Phe(2F), Phe(3F), Phe(4F), Phe(4Cl), Phe(2Me), Phe(3Me), Phe(4Me), threo-Ser(3Phenyl) or erythro-Ser(3Phenyl);
- AzaGly represents azaglycine;
- XX8 represents Leu or Ala(cPr);
- XX9 represents Arg or Arg(Me); and,
- XX10 represents Phe or Trp); or a salt thereof.

(7) Ac-D-Tyr-D-Trp-Asn-Thr-Phe-AzaGly-Leu-D-Arg-Trp- NH_2 (Compound No. 708),

- Ac-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 709),
Decanoyl-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 710),
Acp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 712),
Ac-Acp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 713),
5 Ac-D-Tyr-D-Trp-Asp(NHPen)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound
No. 714),
Ac-D-Tyr-D-Trp-Asp(NHcPr)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound
No. 715),
Ac-D-Tyr-D-Trp-Asp(NHBzl)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound
10 No. 716),
Ac-D-Tyr-D-Trp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
717),
Ac-D-Tyr-D-Pya(4)-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
718),
15 Ac-D-Tyr-D-Trp-Asn-D-Pro-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
719),
Ac-D-Tyr-Aze(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
720),
Ac-D-Tyr-Pic(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
20 721),
Ac-D-Tyr-Pic(3)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 722)
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 723),
Ac-D-Tyr-Thz-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 724),
Ac-D-Tyr-NMeAla-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
25 725).
Ac-D-Tyr-Gly-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 726)
Ac-D-Tyr-Aib-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 727),
Ac-D-Tyr-Abz(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
728)
30 Ac-D-Tyr-Aze(3)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
730)
Ac-D-Tyr-Sar-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 731)
Ac-D-Tyr-D-NMeAla-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
732),
35 Ac-D-Tyr-Izc-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 734),

- Ac-D-Tyr-D-Asp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 735),
Ac-D-Tyr-D-Dap-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 736),
5 Ac-D-Tyr-D-Ser-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 737),
Ac-D-Tyr-D-Gln-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 738),
Ac-D-Tyr-D-His-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 739),
10 Ac-D-Tyr-D-Trp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 740),
Ac-D-Tyr-Ala-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 742),
Ac-D-Tyr-Leu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 743),
15 Ac-D-Tyr-Ser-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 744),
Ac-D-Tyr-Lys-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 745),
Ac-D-Tyr-Glu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 746),
Ac-D-Tyr-β-Ala-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 747),
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 748),
20 Ac-D-Tyr-D-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 749),
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 750),
25 Ac-D-Tyr-Lys-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 754),
Ac-D-Tyr-Glu-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 755),
Ac-D-Tyr-Lys-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 756),
30 Ac-D-Tyr-Glu-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 757),
Ac-D-Tyr-Lys-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 758),
35 Ac-D-Tyr-Glu-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 759),

- 759),
Ac-D-Tyr-Pzc(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
760),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
5 763),
Ac-D-Tyr-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
764),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
765),
10 Ac-D-Tyr-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
766),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
767),
Ac-D-Tyr-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
15 768),
Ac-D-Tyr-Gly-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
769),
Ac-D-Tyr-Aib-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
770),
20 Ac-D-Tyr-Orn-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 771),
Ac-D-Tyr-Thr-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 772),
Ac-D-Tyr-His(3Me)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
773),
Ac-D-Tyr-DL-Ala(Pip)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound
25 No. 774),
Ac-D-Tyr-Tyr(PO₃H₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound
No. 775),
Glycoloyl-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
776)
30 Ac-D-Tyr-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 777),
Ac-D-Tyr-Pro(4NH₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
780),
Ac-D-Tyr-Hyp(Bzl)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
781),
35 Ac-D-Tyr-D-NMePhe-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.

- 782),
Ac-D-Tyr-Gly-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
783),
Ac-D-Tyr-Aib-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
5 784),
Ac-D-Tyr-Gly-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
785),
Ac-D-Tyr-Aib-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
786),
10 Ac-D-Tyr-Hyp-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
787),
Ac-D-Tyr-Glu-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
788),
Ac-D-Tyr-Lys-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
15 789),
Ac-D-Tyr-Gly-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
790),
Ac-D-Tyr-Aib-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
791),
20 Ac-D-Tyr-Hyp-Asn-Thr-D-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
794),
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg-Trp-NH₂ (Compound No. 797),
Ac-D-Tyr-Hyp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 800),
4-[Bis-(2-Pyridylmethyl)aminomethyl]benzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
25 (Compound No. 801),
Ac-D-Tyr-Hyp-Asn-NMeSer-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
809),
Ac-D-Tyr-Hyp-Asn-Hyp-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 810),
Ac-D-Tyr-Hyp-Asn-Gly-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 813),
30 Ac-D-Tyr-Hyp-Asn-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 814),
Ac-D-Tyr-Hyp-Asn-D-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
815),
Ac-D-Tyr-Hyp-His-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 816),
Ac-D-Tyr-Hyp-Gln-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 843),
35 Ac-D-Tyr-Hyp-D-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.

- 844),
Ac-D-Tyr-Hyp-Cit-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 845),
Ac-D-Tyr-Hyp-Asn-D-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 846),
5 Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Ala(cPr)-Arg(Me)-Trp-NH₂ (Compound No. 856),
4-Ureidobenzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 860),
Ac-D-Tyr-Hyp-Arg-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 861),
Ac-D-Tyr-Hyp-Gly-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 862),
10 Ac-D-Tyr-Hyp-Dap-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 863),
Ac-D-Tyr-Hyp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 864),
Ac-D-Tyr-Hyp-Asn-Thr- α MePhe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 868),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(2Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 870),
15 Ac-D-Tyr-Hyp-Asn-Thr-Phe(3Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 872),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 874),
20 Ac-D-Tyr-Hyp-Asn-Thr-threo-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
(Compound No. 877),
Ac-D-Tyr-Hyp-Asn-Thr-erythro-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
(Compound No. 882),
Ac-D-Tyr-Hyp-Nva-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 886),
25 Ac-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 887),
3-(p-Hydroxyphenyl)propionyl-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
(Compound No. 888),
pGlu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 889),
Ac-D-Tyr-cisHyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 896),
30 Ac-D-Tyr-Pro(4F)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 897),
Ac-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 899), or
a salt thereof.
35 (8) Ac-D-Tyr-D-Trp-Asp(NHPen)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

- (Compound No. 714),
Ac-D-Tyr-D-Trp-Asp(NHcPr)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 715),
Ac-D-Tyr-D-Trp-Asp(NHBzl)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 716),
5 Ac-D-Tyr-D-Trp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 717),
Ac-D-Tyr-D-Pya(4)-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 718),
10 Ac-D-Tyr-Aze(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 720),
Ac-D-Tyr-Pic(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 721),
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 723),
15 Ac-D-Tyr-Thz-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 724),
Ac-D-Tyr-Gly-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 726)
Ac-D-Tyr-Aib-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 727),
Ac-D-Tyr-D-NMeAla-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 732),
20 Ac-D-Tyr-D-Gln-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 738),
Ac-D-Tyr-D-His-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 739),
Ac-D-Tyr-D-Trp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 740),
25 Ac-D-Tyr-Ala-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 742),
Ac-D-Tyr-Leu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 743),
Ac-D-Tyr-Ser-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 744),
Ac-D-Tyr-Lys-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 745),
30 Ac-D-Tyr-Glu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 746),
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 748),
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 749),
35 Ac-D-Tyr-D-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.

- 750),
Ac-D-Tyr-Lys-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
754),
Ac-D-Tyr-Glu-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
5 755),
Ac-D-Tyr-Lys-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
756),
Ac-D-Tyr-Glu-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
757),
10 Ac-D-Tyr-Lys-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
758),
Ac-D-Tyr-Glu-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
759),
Ac-D-Tyr-Pzc(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
15 760),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
763),
Ac-D-Tyr-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
764),
20 Ac-D-Tyr-Hyp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
765),
Ac-D-Tyr-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
766),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
25 767),
Ac-D-Tyr-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
768),
Ac-D-Tyr-Gly-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
769),
30 Ac-D-Tyr-Aib-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
770),
Ac-D-Tyr-Orn-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 771),
Ac-D-Tyr-Thr-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 772),
Ac-D-Tyr-His(3Me)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
35 773),

- Ac-D-Tyr-Tyr(PO₃H₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 775),
Glycoloyl-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 776),
5 Ac-D-Tyr-Pro(4NH₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 780),
Ac-D-Tyr-Hyp(Bzl)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 781),
Ac-D-Tyr-D-NMePhe-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 782),
10 Ac-D-Tyr-Gly-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 783),
Ac-D-Tyr-Aib-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 784),
15 Ac-D-Tyr-Gly-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 785),
Ac-D-Tyr-Aib-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 786),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 787),
20 Ac-D-Tyr-Glu-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 788),
Ac-D-Tyr-Lys-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 789),
25 Ac-D-Tyr-Gly-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 790),
Ac-D-Tyr-Aib-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 791),
Ac-D-Tyr-Hyp-Asn-Thr-D-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 794),
30 Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg-Trp-NH₂ (Compound No. 797),
Ac-D-Tyr-Hyp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 800),
4-[Bis-(2-Pyridylmethyl)aminomethyl]benzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 801),
35 Ac-D-Tyr-Hyp-Asn-NMeSer-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.

- 809),
Ac-D-Tyr-Hyp-Asn-Hyp-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 810),
Ac-D-Tyr-Hyp-Asn-Gly-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 813),
Ac-D-Tyr-Hyp-Asn-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 814),
5 Ac-D-Tyr-Hyp-Asn-D-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 815),
Ac-D-Tyr-Hyp-His-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 816),
Ac-D-Tyr-Hyp-Gln-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 843),
Ac-D-Tyr-Hyp-D-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 844),
10 Ac-D-Tyr-Hyp-Cit-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 845),
Ac-D-Tyr-Hyp-Asn-D-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 846),
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Ala(cPr)-Arg(Me)-Trp-NH₂ (Compound No. 856),
15 4-Ureidobenzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 860),
Ac-D-Tyr-Hyp-Arg-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 861),
Ac-D-Tyr-Hyp-Gly-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 862),
Ac-D-Tyr-Hyp-Dap-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 863),
20 Ac-D-Tyr-Hyp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 864),
Ac-D-Tyr-Hyp-Asn-Thr- α MePhe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 868),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(2Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 870),
25 Ac-D-Tyr-Hyp-Asn-Thr-Phe(3Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 872),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 874),
Ac-D-Tyr-Hyp-Asn-Thr-threo-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
30 (Compound No. 877),
Ac-D-Tyr-Hyp-Asn-Thr-erythro-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
(Compound No. 882),
Ac-D-Tyr-Hyp-Nva-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 886),
3-(p-Hydroxyphenyl)propionyl-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
35 (Compound No. 888),

- Ac-D-Tyr-cisHyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 896),
Ac-D-Tyr-Pro(4F)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 897),
5 Ac-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 899), or a salt thereof.
(9) A prodrug of the metastin derivative according to (1), or a salt thereof.
(10) A prodrug of the metastin derivative according to (7), or a salt thereof.
(11) A medicament comprising the metastin derivative according to (1) through (8), a
10 salt thereof, or a prodrug thereof.
(12) The medicament according to (11), which is an agent for suppressing cancer metastasis or an agent for suppressing cancer growth.
(13) The medicament according to (11), which is an agent for preventing or treating cancer.
15 (14) The medicament according to (11), which is an agent for controlling placental function.
(15) The medicament according to (11), which is an agent for preventing or treating choriocarcinoma, hydatid mole, invasive mole, miscarriage, fetal hypoplasia, abnormal glucose metabolism, abnormal lipid metabolism or induction of delivery.
20 (16) The medicament according to (11), which is an agent for improving gonadal function.
(17) The medicament according to (11), which is an agent for preventing or treating hormone-dependent cancer, infertility, endometriosis, early puberty or myoma of the uterus.
25 (18) The medicament according to (11), which is an agent for inducing or stimulating ovulation.
(19) The medicament according to (11), which is a gonadotropic hormone secretagogue agent or a sex hormone secretagogue agent.
(20) The medicament according to (11), which is an agent for preventing or treating
30 Alzheimer's disease, Autism or moderate cognitive impairment.
(21) A method for suppressing cancer metastasis or cancer growth, which comprises administering to a mammal an effective dose of the metastin derivative according to (1) through (8), a salt thereof, or a prodrug thereof.
(22) A method of preventing or treating cancer, which comprises administering to a
35 mammal an effective dose of the metastin derivative according to (1) through (8) or a

salt thereof, or a prodrug thereof.

(23) A method for controlling placental function, which comprises administering to a mammal an effective dose of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof.

5 (24) A method for preventing or treating choriocarcinoma, hydatid mole, invasive mole, miscarriage, fetal hypoplasia, abnormal glucose metabolism, abnormal lipid metabolism or induction of delivery, which comprises administering to a mammal an effective dose of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof.

10 (25) A method for improving gonadal function, which comprises administering to a mammal an effective dose of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof.

(26) A method for preventing or treating hormone-dependent cancer, infertility, endometriosis, early puberty or myoma of the uterus, which comprises administering
15 to a mammal an effective dose of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof.

(27) A method for inducing or stimulating ovulation, which comprises administering to a mammal an effective dose of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof.

20 (28) A method for promoting gonadotropic hormone secretion or promoting sex hormone secretion, which comprises administering to a mammal an effective dose of the metastin derivative according to (1) or a salt thereof, or a prodrug thereof.

(29) A method for preventing or treating Alzheimer's disease, Autism or moderate cognitive impairment, which comprises administering to a mammal an effective dose
25 of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof.

(30) The medicament according to (11), which is a down-regulating agent for gonadotropic hormone or sex hormone.

(31) The medicament according to (11), which is a down-regulating agent for human
30 OT7T175 (metastin receptor) protein consisting of the amino acid sequence represented by SEQ ID NO: 9.

(32) The agent according to (30) or (31), which is an agent for preventing or treating hormone-dependent cancer.

(33) A method for down-regulating gonadotropic hormone or sex hormone, which
35 comprises administering to a mammal an effective dose of the metastin derivative

according to (1) through (8) or a salt thereof, or a prodrug thereof.

(34) A method for down-regulating human OT7T175 (metastin receptor) protein consisting of the amino acid sequence represented by SEQ ID NO: 9, which comprises administering to a mammal an effective dose of the metastin derivative

5 according to (1) through (8) or a salt thereof, or a prodrug thereof.

(35) A method for preventing or treating hormone-dependent cancer, which comprises administering to a mammal an effective dose of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof.

The present invention further provides the following features, and so on.

10 (36) Use of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof to manufacture an agent for suppressing cancer metastasis or an agent for suppressing cancer growth.

(37) Use of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof to manufacture an agent for preventing or treating cancer.

15 (38) Use of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof to manufacture an agent for controlling placental function.

(39) Use of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof to manufacture an agent for preventing or treating choriocarcinoma, hydatid mole, invasive mole, miscarriage, fetal hypoplasia, abnormal glucose
20 metabolism, abnormal lipid metabolism or induction of delivery.

(40) Use of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof to manufacture an agent for improving gonadal function.

(41) Use of the metastin derivative according to (1) through (8) or a salt thereof, or a prodrug thereof to manufacture an agent for preventing or treating

25 hormone-dependent cancer, infertility, endometriosis, early puberty or myoma of the uterus.

(42) Use of the metastin derivative according to (1) through (8), or a salt thereof, or a prodrug thereof to manufacture an agent for inducing or stimulating ovulation.

(43) Use of the metastin derivative according to (1) through (8), or a salt thereof, or a
30 prodrug thereof to manufacture a gonadotropic hormone secretagogue agent or a sex hormone secretagogue agent.

(44) Use of the metastin derivative according to (1) through (8), or a salt thereof, or a prodrug thereof to manufacture an agent for preventing or treating Alzheimer's disease, Autism or moderate cognitive impairment.

35

In the present invention, Tyr-Asn-Trp-Asn-Ser-Phe-Gly-Leu-Arg-Phe-NH₂ (SEQ ID NO: 16) is referred to as metastin 10 (Metastin 10), i.e., MS10.

Herein, the N-terminal Tyr and the C-terminal Phe in MS10 are counted as the 1- and 10-positions, respectively.

5

Tyr-Asn-Trp-Asn-Ser-Phe-Gly-Leu-Arg-Phe-NH₂
 1 2 3 4 5 6 7 8 9 10

For example, [Hph10]MS10 means a peptide wherein Phe at the C terminus (10-position) of MS10 is substituted with Hph.

For example, des(1)-MS10 means a peptide wherein Tyr at the N terminus (1-position) of MS10 is deleted.

For example, des(1-3)-Fmoc-MS10 means a peptide wherein Tyr-Asn-Trp at the N terminus (1 to 3-positions) is deleted and the amino group of Asn at the 4-position is modified with Fmoc.

For example,
 des(1)-Ac-[D-Tyr₂,D-Trp₃,Thr₅,AzaGly₇,D-Arg₉,Trp₁₀]MS10 of Compound No. 708 means a peptide wherein the amino end of MS10 is modified with Ac, Tyr at the N terminus (1-position) is deleted, Asn at the 2-position is substituted with D-Tyr, Trp at the 3-position is substituted with D-Trp, Ser at the 5-position is substituted with Thr, Gly at the 7-position is substituted with AzaGly, Arg at the 9-position is substituted with D-Arg and Phe at the 10-position is substituted with Trp, i.e., Ac-D-Tyr-D-Trp-Asn-Thr-Phe-AzaGly-Leu-D-Arg-Trp-NH₂.

In the formulae described above, XX0 represents a modifying group at the amino end; and each of XX2, XX3, XX4, XX5, XX6, XX8, XX9 and XX10 represents 2-, 3-, 4-, 5-, 6-, 8-, 9- and 10-positions of MS10 described above, respectively.

Each chemical bond "-" between XX0, XX2, XX3, XX4, XX5, XX6, AzaGly, XX8, XX9 and XX10 and NH₂ in the formula:
 "XX0-XX2-XX3-XX4-XX5-XX6-AzaGly-XX8-XX9-XX10-NH₂" represents the following meanings.

The chemical bond "-" in the formula "XX0-XX2" means a bond between the group represented by XX0 and the amino group (α -amino group) containing in XX2. More specifically, formula "XX0-XX2" represents the hydrogen atom in the amino group (NH₂) contained in XX2 is substituted with a group represented by

XX0.

The chemical bond "-" in the formula "XX2-XX3" means that the carboxyl group (α -carboxyl group) contained in XX2 is bound to the amino group (α -amino group) in XX3 through an amide bond. The chemical bonds "-" in the formulae
 5 "XX3-XX4," "XX4-XX5," "XX5-XX6," "XX8-XX9" and "XX9-XX10" have the same significance as described above.

The chemical bond "-" in the formula "XX6-AzaGly" means that the carboxyl group (α -carboxyl group) contained in XX6 is bound to the amino group (α -amino group) in AzaGly [azaglycine] through an amide bond.

10 The chemical bond "-" in the formula "AzaGly-XX8" means that the carboxyl group (α -carboxyl group) contained in AzaGly is bound to the amino group (α -amino group) in XX8 through an amide bond.

The chemical bond "-" in the formula "XX10-NH₂" represents the bond between the carboxyl group (α -carboxyl group) and -NH₂. More specifically, the
 15 formula "XX10-NH₂" indicates that -OH in the carboxyl group (-COOH) contained in XX10 is substituted with -NH₂.

Where XX2, XX3, XX4 or/and XX5 represent the chemical bonds "-", these chemical bonds "-" have the same significance as described above.

Specific examples of these chemical bonds include the bonds represented by
 20 the structural formulae shown in TABLE 1B later described, and so on.

In the formula described above, XX0 represents formyl, a C₁₋₂₀ alkanoyl (e.g., acetyl, propionyl, butyryl, hexanoyl, decanoyl, etc.; preferably represents a C₁₋₆ alkanoyl such as acetyl, propionyl, butyryl, etc.; more preferably, acetyl, etc.), cyclopropanecarbonyl, 6-(acetyl-D-arginylamino)caproyl,
 25 6-((R)-2,3-diaminopropionylamino)caproyl, 6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl, 3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl, acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl, 6-aminocaproyl, 6-acetylaminocaproyl, 4-[bis-(2-pyridylmethyl)aminomethyl]benzoyl or
 30 4-ureidobenzoyl, preferably, represents a C₁₋₁₂ alkanoyl, 6-aminocaproyl, 6-acetylaminocaproyl, glycoloyl, 4-[bis-(2-pyridylmethyl)aminomethyl]benzoyl, 4-ureidobenzoyl, 3-(4-hydroxyphenyl)propionyl or pyroglutamyl, more preferably, represents formyl, a C₁₋₆ alkanoyl or glycoloyl, much more preferably, represents a C₁₋₆ alkanoyl or glycoloyl, and most preferably, represents acetyl or glycoloyl. Also
 35 preferably, XX0 is formyl, a C₁₋₂₀ alkanoyl, cyclopropanecarbonyl,

6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl,
 6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl,
 3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl,
 acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl or
 5 6-aminocaproyl; also preferred are formyl, a C₁₋₁₂ alkanoyl, cyclopropanecarbonyl,
 6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl,
 6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl,
 3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl,
 acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl and
 10 6-aminocaproyl.

In the formula above, XX2 represents Tyr, D-Tyr, D-Ala, D-Leu, D-Phe,
 D-Lys, D-Trp or the chemical bond; preferably represents D-Tyr, Tyr or the chemical
 bond; more preferably represents D-Tyr or the chemical bond; and most preferably
 represents D-Tyr.

15 In the formula above, XX3 represents (i) an amino acid wherein α -amino
 group may optionally be methylated (an amino acid selected from the group
 consisting of Ala [alanine], Arg [arginine], Asn [asparagine], Asp [aspartic acid], Cys
 [cysteine], Gln [glutamine], Glu [glutamic acid], Gly [glycine], His [histidine], Ile
 [isoleucine], Leu [leucine], Lys [lysine], Met [methionine], Phe [phenylalanine], Ser
 20 [serine], Thr [threonine], Trp [tryptophan], Tyr [tyrosine] and Val [valine]), (ii) a
 cyclic amino acid (a cyclic amino acid selected from Pro[proline], Aze(2), Aze(3),
 Pic(2), Pic(3), Hyp, Thz, Abz(2), Abz(3), Pzc(2), Pro(4NH₂), Hyp(Bzl), cisHyp,
 Pro(4F) and Izc), (iii) an amino acid selected from D-Dap, D-Pya(4), DL-Ala(Pip),
 Orn, Aib and Tyr(PO₃H₂), or (iv) the chemical bond.

25 Herein, Aze(2) represents [azetidine-2-carboxylic acid], Aze(3) represents
 [azetidine-3-carboxylic acid], Pic(2) represents [pipecolinic acid], Pic(3) represents
 [3-piperidinecarboxylic acid], D-Dap represents [D-2,3-diaminopropionic acid],
 D-Pya(4) represents [4-pyridyl-D-alanine], Hyp represents [trans-4-hydroxyproline],
 Thz represents [thioprolin], Aib represents [α -aminoisobutanoic acid], Abz(2)
 30 represents [2-aminobenzoic acid], Abz(3) represents [3-aminobenzoic acid], Izc
 represents [imidazolidine-2-carboxylic acid], DL-Ala(Pip) represents
 [DL-(4-piperidin-1-yl)alanine], Pzc(2) represents [piperazine-2-carboxylic acid], Orn
 represents [ornithine], Tyr(PO₃H₂) represents [O-phosphotyrosine], Pro(4NH₂)
 represents [cis-4-aminoproline], Hyp(Bzl) represents [trans-4-benzoyloxyproline],
 35 cisHyp represents [cis-4-hydroxyproline], and Pro(4F) represents

[trans-4-fluoroproline].

Herein, the amino acid may be either an L-amino acid or a D-amino acid.

Alanine may be α -alanine or β -alanine unless otherwise indicated. Preferably, XX3 is D-Asp, D-Dap [D-2,3-diaminopropionic acid], D-Ser, D-Gln, D-His, D-Trp,

- 5 D-Tyr, D-Pya(4), D-NMeAla [D-N ^{α} -methylalanine], D-NMePhe [D-N ^{α} -methylphenylalanine], Aze(2), Aze(3) [azetidine-3-carboxylic acid], Pic(2), Pic(3), Hyp, Thz, NMeAla, Gly, Aib, Abz(2), Abz(3), Sar, Izc, Leu, Lys, Glu, Thr, Trp, Ser, Ala, NMeAla, β -alanine, Pzc(2), Orn, His(3Me) [3-methylhistidine], Tyr(PO₃H₂), Pro(4NH₂), Hyp(Bzl), cisHyp, Pro(4F) or the chemical bond; more
- 10 preferably represents D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-Trp, D-Tyr, D-Pya(4) D-NMeAla, D-NMePhe, Aze(2), Aze(3), Pic(2), Pic(3), Hyp, Thz, Gly, Aib, Abz(2), Sar, Izc, Leu, Lys, Glu, Thr, Trp, Ser, Ala, NMeAla, β -alanine, DL-Ala(Pip), Pzc(2), Orn, His(3Me), Tyr(PO₃H₂), Pro(4NH₂), Hyp(Bzl), cisHyp, Pro(4F) or the chemical bond; particularly preferably represents D-Gln, D-His, Aze(2), Pic(2), Hyp, Thz, Gly,
- 15 Aib, D-NMeAla, Leu, Lys, Glu, Orn, His(3Me), Tyr(PO₃H₂), Pro(4NH₂), D-NMePhe, Hyp(Bzl), cisHyp or Pro(4F), much more preferably represents Aze(2), Hyp, Gly, Aib, Leu, Lys, Glu, His(3Me), Tyr(PO₃H₂), Hyp(Bzl), cisHyp or Pro(4F), and most preferably represents Hyp, Glu, Hyp(Bzl) or Pro(4F). As XX3, D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-NMeAla, D-NMePhe, Aze(2), Pic(2), Pic(3), Hyp,
- 20 Thz, NMeAla, Gly, Aib, Abz(2), Abz(3), Sar, Leu, Lys, Glu, β -alanine, Pzc(2), Orn, His(3Me), Tyr(PO₃H₂), Pro(4NH₂) or Hyp(Bzl) is also preferred.

In the formula above, XX4 represents Asn, 2-amino-3-ureidopropionic acid, N ^{β} -formyl- β -diaminopropionic acid, N ^{β} -acetyl- β -diaminopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine,

- 25 2,4-diaminobutanoic acid, 2,3-diaminopropionic acid, His, Gln, Gly, Arg, Cit, Nva, D-Asn or the chemical bond; preferably represents Asn, 2-amino-3-ureidopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine, 2,4-diaminobutanoic acid, 2,3-diaminopropionic acid, His, Gln, Gly, Arg, Cit, Nva, D-Asn or the chemical bond; more preferably represents Asn or
- 30 2-amino-3-ureidopropionic acid. Also preferably, XX4 represents Asn, 2-amino-3-ureidopropionic acid, N ^{β} -formyl- β -diaminopropionic acid, N ^{β} -acetyl- β -diaminopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine, 2,4-diaminobutanoic acid, His, Gln, Cit or D-Asn; or, Asn, 2-amino-3-ureidopropionic acid, N ^{β} -formyldiaminopropionic acid,
- 35 N ^{β} -acetyldiaminopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine,

N^ω-benzylasparagine or 2,4-diaminobutanoic acid is also preferred.

In the formula described above, XX5 represents Ser, Thr, Val, NMeSer, Gly, Ala, Hyp, D-Ala, D-Thr, D-Pro or the chemical bond; preferably represents Thr, NMeSer, Gly, Ala, Hyp, D-Ala, D-Thr, D-Pro or the chemical bond; more preferably represents Ser, Thr or Ala, and most preferably represents Thr. Also preferably, XX5 is Ser, Thr, Val, NMeSer, Gly, Ala, Hyp, D-Ala or D-Thr; or, Ser, Thr or Val is also preferred.

In the formula described above, XX6 represents Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), threo-Ser(3Phenyl), erythro-Ser(3Phenyl) or an optionally substituted phenylalanine. Herein, the substituent in the optionally substituted phenylalanine includes, for example, oxo, a halogen atom (e.g., fluorine, chlorine, bromine, iodine, etc.), a C₁₋₃ alkylenedioxy (e.g., methylenedioxy, ethylenedioxy, etc.), nitro, cyano, an optionally substituted C₁₋₆ alkyl, an optionally substituted C₂₋₆ alkenyl, an optionally substituted C₂₋₆ alkynyl, an optionally substituted C₃₋₈ cycloalkyl, an optionally substituted C₆₋₁₄ aryl, an optionally substituted C₇₋₁₆ aralkyl, an optionally substituted C₁₋₆ alkoxy, hydroxy, an optionally substituted C₆₋₁₄ aryloxy, an optionally substituted C₇₋₁₆ aralkyloxy, mercapto, an optionally substituted C₁₋₆ alkylthio, an optionally substituted C₆₋₁₄ arylthio, an optionally substituted C₇₋₁₆ aralkylthio, an optionally substituted amino[amino, an optionally substituted mono- or di-C₁₋₆ alkyl-amino (e.g., methylamino, dimethylamino, ethylamino, diethylamino, propylamino, isopropylamino, etc.), an optionally substituted mono- or di-C₂₋₆ alkenyl-amino (e.g., vinylamino, propenylamino, isopropenylamino), an optionally substituted C₂₋₆ alkynyl-amino (e.g., 2-butyne-1-yl-amino, 4-pentyne-1-yl-amino, 5-hexyne-1-yl-amino), an optionally substituted mono- or di-C₃₋₈ cycloalkyl-amino (e.g., cyclopropylamino, cyclohexylamino), an optionally substituted C₆₋₁₄ aryl-amino (e.g., phenylamino, diphenylamino, naphthylamino), an optionally substituted C₁₋₆ alkoxy-amino (e.g., methoxyamino, ethoxyamino, propoxyamino, isopropoxyamino), formylamino, an optionally substituted C₁₋₆ alkyl-carbonylamino (e.g., acetylamino, propionylamino, pivaloylamino, etc.), an optionally substituted C₃₋₈ cycloalkyl-carbonylamino (e.g., cyclopropylcarbonylamino, cyclopentylcarbonylamino, cyclohexylcarbonylamino, etc.), an optionally substituted C₆₋₁₄ aryl-carbonylamino (e.g., benzoylamino, naphthoylamino, etc.), an optionally substituted C₁₋₆ alkoxy-carbonylamino (e.g., methoxycarbonylamino, ethoxycarbonylamino, propoxycarbonylamino, butoxycarbonylamino, etc.), an optionally substituted C₁₋₆ alkylsulfonylamino (e.g.,

methylsulfonylamino, ethylsulfonylamino, etc.), an optionally substituted C₆₋₁₄
 arylsulfonylamino (e.g., phenylsulfonylamino, 2-naphthylsulfonylamino,
 1-naphthylsulfonylamino, etc.)), formyl, carboxy, an optionally substituted C₁₋₆
 alkyl-carbonyl (e.g., acetyl, propionyl, pivaloyl, etc.), an optionally substituted C₃₋₈
 5 cycloalkyl-carbonyl (e.g., cyclopropylcarbonyl, cyclopentylcarbonyl,
 cyclohexylcarbonyl, 1-methylcyclohexyl-carbonyl, etc.), an optionally substituted
 C₆₋₁₄ aryl-carbonyl (e.g., benzoyl, 1-naphthoyl, 2-naphthoyl, etc.), an optionally
 substituted C₇₋₁₆ aralkyl-carbonyl (e.g., phenylacetyl, 3-phenylpropionyl, etc.), an
 optionally substituted 5- to 7-membered heterocyclic carbonyl containing 1 to 4
 10 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in
 addition to carbon atoms (e.g., nicotinoyl, isonicotinoyl, thenoyl, furoyl,
 morpholinocarbonyl, thiomorpholinocarbonyl, piperazin-1-ylcarbonyl,
 pyrrolidin-1-ylcarbonyl, etc.), an optionally esterified carboxyl, an optionally
 substituted carbamoyl group, an optionally substituted C₁₋₆ alkylsulfonyl (e.g.,
 15 methylsulfonyl, ethylsulfonyl, etc.), an optionally substituted C₁₋₆ alkylsulfinyl (e.g.,
 methylsulfinyl, ethylsulfinyl, etc.), an optionally substituted C₆₋₁₄ arylsulfonyl (e.g.,
 phenylsulfonyl, 1-naphthylsulfonyl, 2-naphthylsulfonyl, etc.), an optionally
 substituted C₆₋₁₄ arylsulfinyl (e.g., phenylsulfinyl, 1-naphthylsulfinyl,
 2-naphthylsulfinyl, etc.), an optionally substituted C₁₋₆ alkyl-carbonyloxy (e.g.,
 20 acetoxyl, propionyl, etc.), an optionally substituted C₆₋₁₄ aryl-carbonyloxy (e.g.,
 benzoyloxy, naphthylcarbonyloxy, etc.), an optionally substituted C₁₋₆
 alkoxy-carbonyloxy (e.g., methoxycarbonyloxy, ethoxycarbonyloxy,
 propoxycarbonyloxy, butoxycarbonyloxy, etc.), an optionally substituted mono-C₁₋₆
 alkylcarbamoyloxy (e.g., methylcarbamoyloxy, ethylcarbamoyloxy, etc.), an
 25 optionally substituted di-C₁₋₆ alkylcarbamoyloxy (e.g., dimethylcarbamoyloxy,
 diethylcarbamoyloxy, etc.), an optionally substituted mono- or di-C₆₋₁₄
 arylcarbamoyloxy (e.g., phenylcarbamoyloxy, naphthylcarbamoyloxy, etc.), an
 optionally substituted heterocyclic group, sulfo, sulfamoyl, sulfinamoyl,
 sulfenamoyl, or a group of 2 or more (e.g., 2 or 3) of these substituents combined,
 30 and the like (Substituent Group A). The number of the substituents is not
 particularly limited but these groups may have 1 to 5, preferably 1 to 3 substituents
 in substitutable positions, and when there are two or more substituents, each
 substituent may be the same or different.

The "optionally esterified carboxyl group" in the Substituent Group A
 35 includes, for example, an optionally substituted C₁₋₆ alkoxy-carbonyl (e.g.,

methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), an optionally substituted C₆₋₁₄ aryloxy-carbonyl (e.g., phenoxycarbonyl, etc.), an optionally substituted C₇₋₁₆ aralkyloxy-carbonyl (e.g., benzyloxycarbonyl, phenethyloxycarbonyl, etc.), and the like.

5 The "C₁₋₆ alkyl" in the "optionally substituted C₁₋₆ alkyl" in the Substituent Group A includes, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, hexyl, etc.

 The "C₂₋₆ alkenyl" in the "optionally substituted C₂₋₆ alkenyl" in the Substituent Group A includes, for example, vinyl, propenyl, isopropenyl,
10 2-buten-1-yl, 4-penten-1-yl, 5-hexen-1-yl, etc.

 The "C₂₋₆ alkynyl" in the "optionally substituted C₂₋₆ alkynyl" in the Substituent Group A includes, for example, 2-butyne-1-yl, 4-pentyne-1-yl, 5-hexyne-1-yl, etc.

 The "C₃₋₈ cycloalkyl" in the "optionally substituted C₃₋₈ cycloalkyl" in the
15 Substituent Group A includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.

 The "C₆₋₁₄ aryl" in the "optionally substituted C₆₋₁₄ aryl" in the Substituent Group A includes, for example, phenyl, 1-naphthyl, 2-naphthyl, 2-biphenyl, 3-biphenyl, 4-biphenyl, 2-anthryl, etc.

20 The "C₇₋₁₆ aralkyl" in the "optionally substituted C₇₋₁₆ aralkyl" in the Substituent Group A includes, for example, benzyl, phenethyl, diphenylmethyl, 1-naphthylmethyl, 2-naphthylmethyl, 2,2-diphenylethyl, 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, 2-biphenylmethyl, 3-biphenylmethyl, 4-biphenylmethyl), etc.

25 The "C₁₋₆ alkoxy" in the "optionally substituted C₁₋₆ alkoxy" in the Substituent Group A includes, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, pentyloxy, hexyloxy, etc.

 The "C₆₋₁₄ aryloxy" in the "optionally substituted C₆₋₁₄ aryloxy" in the Substituent Group A includes, for example, phenyloxy, 1-naphthyloxy, 2-naphthyloxy,
30 etc.

 The "C₇₋₁₆ aralkyloxy" in the "optionally substituted C₇₋₁₆ aralkyloxy" in the Substituent Group A includes, for example, benzyloxy, phenethyloxy, etc.

 The "C₁₋₆ alkylthio" in the "optionally substituted C₁₋₆ alkylthio" in the Substituent Group A includes, for example, methylthio, ethylthio, propylthio,
35 isopropylthio, butylthio, sec-butylthio, tert-butylthio, etc.

The "C₆₋₁₄ arylthio" in the "optionally substituted C₆₋₁₄ arylthio" in the Substituent Group A includes, for example, phenylthio, 1-naphthylthio, 2-naphthylthio, etc.

5 The "C₇₋₁₆ aralkylthio" in the "optionally substituted C₇₋₁₆ aralkylthio" in the Substituent Group A includes, for example, benzylthio, phenethylthio, etc.

The substituents for these "C₁₋₆ alkoxy-carbonyl," "C₁₋₆ alkyl group," "C₂₋₆ alkenyl," "C₂₋₆ alkynyl," "C₁₋₆ alkoxy," "C₁₋₆ alkylthio," "C₁₋₆ alkyl-amino," "C₂₋₆ alkenyl-amino," "C₂₋₆ alkynyl-amino," "C₁₋₆ alkoxy-amino," "C₁₋₆ alkyl-carbonyl," "C₁₋₆ alkylsulfonyl," "C₁₋₆ alkylsulfinyl," "C₁₋₆ alkyl-carbonylamino," "C₁₋₆ alkoxy-carbonylamino," "C₁₋₆ alkylsulfonylamino," "C₁₋₆ alkyl-carbonyloxy," "C₁₋₆ alkoxy-carbonyloxy," "mono-C₁₋₆ alkylcarbamoyloxy" and "di-C₁₋₆ alkylcarbamoyloxy" in the Substituent Group A include, for example, 1 to 5 substituents selected from, for example, a halogen atom (e.g., fluorine atom, chlorine atom, bromine atom, iodine atom), carboxy, hydroxy, amino, a mono- or di-C₁₋₆ alkylamino, a mono- or di-C₆₋₁₄ arylamino, a C₃₋₈ cycloalkyl, a C₁₋₆ alkoxy, a C₁₋₆ alkoxy-carbonyl, a C₁₋₆ alkylthio, a C₁₋₆ alkylsulfinyl, a C₁₋₆ alkylsulfonyl, the optionally esterified carboxyl described above, carbamoyl, thiocarbamoyl, a mono-C₁₋₆ alkylcarbamoyl (e.g., methylcarbamoyl, ethylcarbamoyl, etc.), a di-C₁₋₆ alkylcarbamoyl (e.g., dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl, etc.), a mono- or di-C₆₋₁₄ arylcarbamoyl (e.g., phenylcarbamoyl, 1-naphthylcarbamoyl, 2-naphthylcarbamoyl, etc.), a mono- or di-5- to 7-membered heterocyclic carbamoyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms (e.g., 2-pyridylcarbamoyl, 3-pyridylcarbamoyl, 4-pyridylcarbamoyl, 2-thienylcarbamoyl, 3-thienylcarbamoyl, etc.), and the like.

The substituents for the "C₆₋₁₄ aryloxy-carbonyl," "C₇₋₁₆ aralkyloxy-carbonyl," "C₃₋₈ cycloalkyl," "C₆₋₁₄ aryl," "C₇₋₁₆ aralkyl," "C₆₋₁₄ aryloxy," "C₇₋₁₆ aralkyloxy," "C₆₋₁₄ arylthio," "C₇₋₁₆ aralkylthio," "C₃₋₈ cycloalkyl-amino," "C₆₋₁₄ aryl-amino," "C₃₋₈ cycloalkyl-carbonyl," "C₆₋₁₄ aryl-carbonyl," "C₇₋₁₆ aralkyl-carbonyl," "5- to 7-membered heterocyclic carbonyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms," "C₆₋₁₄ arylsulfonyl," "C₆₋₁₄ arylsulfinyl," "C₃₋₈ cycloalkyl-carbonylamino," "C₆₋₁₄ aryl-carbonylamino," "C₆₋₁₄ arylsulfonylamino," "C₆₋₁₄ aryl-carbonyloxy" and "mono- or di-C₆₋₁₄ aryl-carbamoyloxy" in the Substituent Group A include, for example, 1 to 5 substituents selected from, for

example, a halogen atom, hydroxy, carboxy, nitro, cyano, the optionally substituted C₁₋₆ alkyl described above, the optionally substituted C₂₋₆ alkenyl described above, the optionally substituted C₂₋₆ alkynyl described above, the optionally substituted C₃₋₈ cycloalkyl described above, the optionally substituted C₁₋₆ alkoxy described
5 above, the optionally substituted C₁₋₆ alkylthio described above, the optionally substituted C₁₋₆ alkylsulfinyl described above, the optionally substituted C₁₋₆ alkylsulfonyl described above, the optionally esterified carboxyl described above, carbamoyl, thiocarbamoyl, a mono-C₁₋₆ alkylcarbamoyl, a di-C₁₋₆ alkylcarbamoyl, a mono- or di-C₆₋₁₄ arylcarbamoyl, a mono- or di-5- to 7-membered heterocyclic
10 carbamoyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms, and the like.

The "optionally substituted heterocyclic group" in the Substituent Group A includes, for example, a 5- to 14-membered (monocyclic, bicyclic or tricyclic) heterocyclic group containing 1 to 4 hetero atoms of 1 or 2 species selected from
15 nitrogen, sulfur and oxygen atoms in addition to carbon atoms, which may optionally be substituted with a halogen atom, hydroxy, carboxy, nitro, cyano, the optionally substituted C₁₋₆ alkyl described above, the optionally substituted C₂₋₆ alkenyl described above, the optionally substituted C₂₋₆ alkynyl described above, the optionally substituted C₃₋₈ cycloalkyl described above, the optionally substituted C₆₋₁₄ aryl described above, the optionally substituted C₁₋₆ alkoxy described above, the optionally substituted C₁₋₆ alkylthio described above, the optionally substituted C₆₋₁₄ arylthio described above, the optionally substituted C₇₋₁₆ aralkylthio described above, the optionally substituted C₁₋₆ alkylsulfinyl described above, the optionally substituted C₆₋₁₄ arylsulfinyl described above, the optionally substituted C₁₋₆
20 alkylsulfonyl described above, the optionally substituted C₆₋₁₄ arylsulfonyl described above, the optionally esterified carboxyl described above, carbamoyl, thiocarbamoyl, a mono-C₁₋₆ alkylcarbamoyl, a di-lower alkylcarbamoyl, a mono- or di-C₆₋₁₄ arylcarbamoyl, a mono- or di-5- or 7-membered heterocyclic carbamoyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms
25 in addition to carbon atoms, or the like; preferably (i) a 5- to 14-membered (preferably, 5- to 10-membered) aromatic heterocyclic group, (ii) a 5- to 10-membered non-aromatic heterocyclic group or (iii) a monovalent group formed by one optional hydrogen atom from 7- to 10-membered bridged-hetero ring, and the like, are employed; among them, preferably used is a 5-membered aromatic
30 heterocyclic group. Specifically used are an aromatic heterocyclic group such as

35

thienyl (e.g., 2-thienyl, 3-thienyl), furyl (e.g., 2-furyl, 3-furyl), pyridyl (e.g., 2-pyridyl, 3-pyridyl, 4-pyridyl), thiazolyl (e.g., 2-thiazolyl, 4-thiazolyl, 5-thiazolyl), oxazolyl (e.g., 2-oxazolyl, 4-oxazolyl), quinolyl (e.g., 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 8-quinolyl), isoquinolyl (e.g., 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl), pyrazinyl, pyrimidinyl (e.g., 2-pyrimidinyl, 4-pyrimidinyl), pyrrolyl (e.g., 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl), imidazolyl (e.g., 1-imidazolyl, 2-imidazolyl, 4-imidazolyl), pyrazolyl (e.g., 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl), pyridazinyl (e.g., 3-pyridazinyl, 4-pyridazinyl), isothiazolyl (e.g., 3-isothiazolyl), isoxazolyl (e.g., 3-isoxazolyl), indolyl (e.g., 1-indolyl, 2-indolyl, 3-indolyl), 2-benzothiazolyl, benzo[b]thienyl, (e.g., 2-benzo[b]thienyl, 3-benzo[b]thienyl), benzo[b]furanlyl (e.g., 2-benzo[b]furanlyl, 3-benzo[b]furanlyl), etc., a non-aromatic heterocyclic group such as pyrrolidinyl (e.g., 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl), oxazolidinyl (e.g., 2-oxazolidinyl), imidazolinyl (e.g., 1-imidazolinyl, 2-imidazolinyl, 4-imidazolinyl), piperidinyl (e.g., 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl), piperazinyl (e.g., 1-piperazinyl, 2-piperazinyl), morpholino, thiomorpholino, etc.

The "optionally substituted carbamoyl group" in the Substituent Group A includes a carbamoyl group, which may optionally be substituted with the optionally substituted C₁₋₆ alkyl, optionally substituted C₂₋₆ alkenyl, an optionally substituted C₂₋₆ alkynyl, an optionally substituted C₃₋₈ cycloalkyl, an optionally substituted C₆₋₁₄ aryl, an optionally substituted heterocyclic group described above, etc., and specific examples are carbamoyl, thiocarbamoyl, a mono-C₁₋₆ alkylcarbamoyl (e.g., methylcarbamoyl, ethylcarbamoyl, etc.), a di-C₁₋₆ alkylcarbamoyl (e.g., dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl, etc.), a C₁₋₆ alkyl(C₁₋₆ alkoxy)carbamoyl (e.g., methyl(methoxy)carbamoyl, ethyl(methoxy)carbamoyl), a mono- or di-C₆₋₁₄ arylcarbamoyl (e.g., phenylcarbamoyl, 1-naphthylcarbamoyl, 2-naphthylcarbamoyl, etc.), a mono- or di-5- to 7-membered heterocyclic carbamoyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms (e.g., 2-pyridylcarbamoyl, 3-pyridylcarbamoyl, 4-pyridylcarbamoyl, 2-thienylcarbamoyl, 3-thienylcarbamoyl, etc.), a 5- to 7-membered cyclic carbamoyl (e.g., 1-pyrrolidinylcarbonyl, 1-piperidinylcarbonyl, hexamethyleneiminocarbonyl), and the like.

The "optionally substituted amino" in the Substituent Group A includes an amino, which may optionally be substituted with 1 or 2 groups selected from the optionally substituted C₁₋₆ alkyl described above, the optionally substituted C₂₋₆

alkenyl described above, the optionally substituted C₂₋₆ alkynyl described above, the optionally substituted C₃₋₈ cycloalkyl described above, the optionally substituted C₆₋₁₄ aryl described above, the optionally substituted C₁₋₆ alkoxy described above, formyl, the optionally substituted C₁₋₆ alkyl-carbonyl described above, the optionally substituted C₃₋₈ cycloalkyl-carbonyl described above, the optionally substituted C₆₋₁₄ aryl-carbonyl described above, the optionally substituted C₁₋₆ alkoxy-carbonyl described above, the optionally substituted C₁₋₆ alkylsulfonyl described above, the optionally substituted C₆₋₁₄ arylsulfonyl, and the like.

More preferably, the substituents are a halogen atom, hydroxy, a C₁₋₆ alkoxy, an optionally halogenated C₁₋₆ alkyl, an optionally halogenated C₁₋₆ alkoxy, amino, nitro, cyano, etc.

XX6 preferably represents Phe, Tyr, Trp, Tyr(Me) [O-methyltyrosine], Thi [2-thienylalanine], Nal(2) [2-naphthylalanine], Cha [cyclohexylalanine], Pya(4) [4-pyridylalanine], Phe(2F) [2-fluorophenylalanine], Phe(3F) [3-fluorophenylalanine], Phe(4F) [4-fluorophenylalanine], Phe(4Cl) [4-chlorophenylalanine], α MePhe [α -methylphenylalanine], Phe(2Me), Phe(3Me), Phe(4Me), threo-Ser(3Phenyl), erythro-Ser(3Phenyl) or D-Phe, more preferably represents Phe, Cha, Phe(2F), Phe(3F), Phe(4F), Phe(4Cl), α MePhe, Phe(2Me), Phe(3Me), Phe(4Me), threo-Ser(3Phenyl), erythro-Ser(3Phenyl) or D-Phe, further more preferably represents Phe, Phe(2F), Phe(3F), Phe(4F), Phe(4Cl), α MePhe, Phe(2Me), Phe(3Me), Phe(4Me), threo-Ser(3Phenyl), erythro-Ser(3Phenyl) or D-Phe, much more preferably represents Phe, Cha, Phe(2F), Phe(3F), Phe(4F), Phe(4Cl), Phe(2Me), Phe(3Me), Phe(4Me), threo-Ser(3Phenyl) or erythro-Ser(3Phenyl), and most preferably represents Phe, Cha, Phe(3F) or Phe(4F). As XX6, Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), Phe(2F), Phe(3F), Phe(4F), Phe(4Cl) or D-Phe; or Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), Phe(2F), Phe(3F), Phe(4F) or Phe(4Cl) is also preferred.

In the formula described above, AzaGly represents azaglycine.

In the formula described above, XX8 represents Leu, Nva [norvaline], Val or Ala(cPr) [cyclopropylalanine], and preferably represents Leu or Ala(cPr). As XX8, Leu, Nva or Val is also preferred.

In the formula described above, XX9 represents an optionally substituted arginine, an optionally substituted lysine or an optionally substituted ornithine. Herein, substituents for the optionally substituted arginine, the optionally substituted lysine or the optionally substituted ornithine are 1 or substitutable number of C₁₋₆

alkyl (e.g., methyl, ethyl, propyl, isopropyl, butyl, etc.), C₁₋₆ acyl (e.g., acetyl, propionyl, etc.). XX9 preferably represents Arg, Orn [ornithine], Arg(Me) [N^ω-methylarginine], D-Arg or Arg (asymMe₂) [asymmetric-N^{ω,ω}-dimethylarginine], more preferably represents Arg, Arg(Me) or D-Arg, further more preferably
 5 represents Arg or Arg(Me). As XX9, Arg, Orn, Arg (Me) or Arg (asymMe₂) is also preferred.

In the formula described above, XX10 represents 2-naphthylalanine, 2-thienylalanine, tyrosine, an optionally substituted phenylalanine, or an optionally substituted tryptophan. Herein, the substituents in the optionally substituted
 10 phenylalanine and the optionally substituted tryptophan include, for example, oxo, a halogen atom (e.g., fluorine, chlorine, bromine, iodine, etc.), a C₁₋₃ alkylendioxy (e.g., methylenedioxy, ethylenedioxy, etc.), nitro, cyano, an optionally substituted C₁₋₆ alkyl, an optionally substituted C₂₋₆ alkenyl, an optionally substituted C₂₋₆ alkynyl, an optionally substituted C₃₋₈ cycloalkyl, an optionally substituted C₆₋₁₄ aryl,
 15 an optionally substituted C₇₋₁₆ aralkyl, an optionally substituted C₁₋₆ alkoxy, hydroxy, an optionally substituted C₆₋₁₄ aryloxy, an optionally substituted C₇₋₁₆ aralkyloxy, mercapto, an optionally substituted C₁₋₆ alkylthio, an optionally substituted C₆₋₁₄ arylthio, an optionally substituted C₇₋₁₆ aralkylthio, an optionally substituted amino [amino, an optionally substituted mono- or di-C₁₋₆ alkyl-amino (e.g., methylamino, dimethylamino, ethylamino, diethylamino, propylamino, isopropylamino, etc.), an
 20 optionally substituted mono- or di-C₂₋₆ alkenyl-amino (e.g., vinylamino, propenylamino, isopropenylamino), an optionally substituted C₂₋₆ alkynyl-amino (e.g., 2-butyne-1-yl-amino, 4-pentyne-1-yl-amino, 5-hexyne-1-yl-amino), an optionally substituted mono- or di-C₃₋₈ cycloalkyl-amino (e.g., cyclopropylamino, cyclohexylamino), an optionally substituted C₆₋₁₄ aryl-amino (e.g., phenylamino, diphenylamino, naphthylamino), an optionally substituted C₁₋₆ alkoxy-amino (e.g., methoxyamino, ethoxyamino, propoxyamino, isopropoxyamino), formylamino, an optionally substituted C₁₋₆ alkyl-carbonylamino (e.g., acetylamino, propionylamino, pivaloylamino, etc.), an optionally substituted C₃₋₈ cycloalkyl-carbonylamino (e.g., cyclopropylcarbonylamino, cyclopentylcarbonylamino, cyclohexylcarbonylamino, etc.), an optionally substituted C₆₋₁₄ aryl-carbonylamino (e.g., benzoylamino, naphthoylamino, etc.), an optionally substituted C₁₋₆ alkoxy-carbonylamino (e.g., methoxycarbonylamino, ethoxycarbonylamino, propoxycarbonylamino, butoxycarbonylamino, etc.), an optionally substituted C₁₋₆ alkylsulfonylamino (e.g.,
 35 methylsulfonylamino, ethylsulfonylamino, etc.), an optionally substituted C₆₋₁₄

arylsulfonylamino (e.g., phenylsulfonylamino, 2-naphthylsulfonylamino, 1-naphthylsulfonylamino, etc.)), formyl, carboxy, an optionally substituted C₁₋₆ alkyl-carbonyl (e.g., acetyl, propionyl, pivaloyl, etc.), an optionally substituted C₃₋₈ cycloalkyl-carbonyl (e.g., cyclopropylcarbonyl, cyclopentylcarbonyl, cyclohexylcarbonyl, 1-methylcyclohexyl-carbonyl, etc.), an optionally substituted C₆₋₁₄ aryl-carbonyl (e.g., benzoyl, 1-naphthoyl, 2-naphthoyl, etc.), an optionally substituted C₇₋₁₆ aralkyl-carbonyl (e.g., phenylacetyl, 3-phenylpropionyl, etc.), an optionally substituted 5- to 7-membered heterocyclic carbonyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms (e.g., nicotinoyl, isonicotinoyl, thenoyl, furoyl, morpholinocarbonyl, thiomorpholinocarbonyl, piperazin-1-ylcarbonyl, pyrrolidin-1-ylcarbonyl, etc.), an optionally esterified carboxyl, an optionally substituted carbamoyl group, an optionally substituted C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl, ethylsulfonyl, etc.), an optionally substituted C₁₋₆ alkylsulfinyl (e.g., methylsulfinyl, ethylsulfinyl, etc.), an optionally substituted C₆₋₁₄ arylsulfonyl (e.g., phenylsulfonyl, 1-naphthylsulfonyl, 2-naphthylsulfonyl, etc.), an optionally substituted C₆₋₁₄ arylsulfinyl (e.g., phenylsulfinyl, 1-naphthylsulfinyl, 2-naphthylsulfinyl, etc.), an optionally substituted C₁₋₆ alkyl-carbonyloxy (e.g., acetoxy, propionyloxy, etc.), an optionally substituted C₆₋₁₄ aryl-carbonyloxy (e.g., benzoyloxy, naphthylcarbonyloxy, etc.), an optionally substituted C₁₋₆ alkoxy-carbonyloxy (e.g., methoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, butoxycarbonyloxy, etc.), an optionally substituted mono-C₁₋₆ alkylcarbamoyloxy (e.g., methylcarbamoyloxy, ethylcarbamoyloxy, etc.), an optionally substituted di-C₁₋₆ alkylcarbamoyloxy (e.g., dimethylcarbamoyloxy, diethylcarbamoyloxy, etc.), an optionally substituted mono- or di-C₆₋₁₄ arylcarbamoyloxy (e.g., phenylcarbamoyloxy, naphthylcarbamoyloxy, etc.), an optionally substituted heterocyclic group, sulfo, sulfamoyl, sulfinamoyl, sulfenamoyl, or a group of 2 or more (e.g., 2 or 3) of these substituents combined, and the like (Substituent Group A). The number of the substituents is not particularly limited but these groups may have 1 to 5, preferably 1 to 3 substituents in substitutable positions, and when there are two or more substituents, each substituent may be the same or different.

The "optionally esterified carboxyl group" in the Substituent Group A includes, for example, an optionally substituted C₁₋₆ alkoxy-carbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), an

optionally substituted C₆₋₁₄ aryloxy-carbonyl (e.g., phenoxycarbonyl, etc.), an optionally substituted C₇₋₁₆ aralkyloxy-carbonyl (e.g., benzyloxycarbonyl, phenethyloxycarbonyl, etc.), and the like.

The "C₁₋₆ alkyl" in the "optionally substituted C₁₋₆ alkyl" in the Substituent Group A includes, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, hexyl, etc.

The "C₂₋₆ alkenyl" in the "optionally substituted C₂₋₆ alkenyl" in the Substituent Group A includes, for example, vinyl, propenyl, isopropenyl, 2-buten-1-yl, 4-penten-1-yl, 5-hexen-1-yl, etc.

The "C₂₋₆ alkynyl" in the "optionally substituted C₂₋₆ alkynyl" in the Substituent Group A includes, for example, 2-butyne-1-yl, 4-pentyne-1-yl, 5-hexyne-1-yl, etc.

The "C₃₋₈ cycloalkyl" in the "optionally substituted C₃₋₈ cycloalkyl" in the Substituent Group A includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.

The "C₆₋₁₄ aryl" in the optionally substituted C₆₋₁₄ aryl in the Substituent Group A includes, for example, phenyl, 1-naphthyl, 2-naphthyl, 2-biphenyl, 3-biphenyl, 4-biphenyl, 2-anthryl, etc.

The "C₇₋₁₆ aralkyl" in the "optionally substituted C₇₋₁₆ aralkyl" in the Substituent Group A includes, for example, benzyl, phenethyl, diphenylmethyl, 1-naphthylmethyl, 2-naphthylmethyl, 2,2-diphenylethyl, 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, 2-biphenylmethyl, 3-biphenylmethyl, 4-biphenylmethyl), etc.

The "C₁₋₆ alkoxy" in the "optionally substituted C₁₋₆ alkoxy" in the Substituent Group A includes, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, pentyloxy, hexyloxy, etc.

The "C₆₋₁₄ aryloxy" in the "optionally substituted C₆₋₁₄ aryloxy" in the Substituent Group A includes, for example, phenyloxy, 1-naphthyloxy, 2-naphthyloxy, etc.

The "C₇₋₁₆ aralkyloxy" in the "optionally substituted C₇₋₁₆ aralkyloxy" in the Substituent Group A includes, for example, benzyloxy, phenethyloxy, etc.

The "C₁₋₆ alkylthio" in the "optionally substituted C₁₋₆ alkylthio" in the Substituent Group A includes, for example, methylthio, ethylthio, propylthio, isopropylthio, butylthio, sec-butylthio, tert-butylthio, etc.

The "C₆₋₁₄ arylthio" in the "optionally substituted C₆₋₁₄ arylthio" in the

Substituent Group A includes, for example, phenylthio, 1-naphthylthio, 2-naphthylthio, etc.

The "C₇₋₁₆ aralkylthio" in the "optionally substituted C₇₋₁₆ aralkylthio" in the Substituent Group A includes, for example, benzylthio, phenethylthio, etc.

5 The substituents for these "C₁₋₆ alkoxy-carbonyl," "C₁₋₆ alkyl group," "C₂₋₆ alkenyl," "C₂₋₆ alkynyl," "C₁₋₆ alkoxy," "C₁₋₆ alkylthio," "C₁₋₆ alkyl-amino," "C₂₋₆ alkenyl-amino," "C₂₋₆ alkynyl-amino," "C₁₋₆ alkoxy-amino," "C₁₋₆ alkyl-carbonyl," "C₁₋₆ alkylsulfonyl," "C₁₋₆ alkylsulfinyl," "C₁₋₆ alkyl-carbonylamino," "C₁₋₆ alkoxy-carbonylamino," "C₁₋₆ alkylsulfonylamino," "C₁₋₆ alkyl-carbonyloxy," "C₁₋₆ alkoxy-carbonyloxy," "mono-C₁₋₆ alkylcarbamoyloxy" and "di-C₁₋₆ alkylcarbamoyloxy" include 1 to 5 substituents selected from, for example, a halogen atom (e.g., fluorine atom, chlorine atom, bromine atom, iodine atom), carboxy, hydroxy, amino, a mono- or di-C₁₋₆ alkylamino, a mono- or di-C₆₋₁₄ arylamino, a C₃₋₈ cycloalkyl, a C₁₋₆ alkoxy, a C₁₋₆ alkoxy-carbonyl, a C₁₋₆ alkylthio, a C₁₋₆ alkylsulfinyl, a C₁₋₆ alkylsulfonyl, the optionally esterified carboxyl described above, carbamoyl, thiocarbamoyl, a mono-C₁₋₆ alkylcarbamoyl (e.g., methylcarbamoyl, ethylcarbamoyl, etc.), a di-C₁₋₆ alkylcarbamoyl (e.g., dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl, etc.), a mono- or di-C₆₋₁₄ arylcarbamoyl (e.g., phenylcarbamoyl, 1-naphthylcarbamoyl, 2-naphthylcarbamoyl, etc.), a mono- or di-5- to 7-membered heterocyclic carbamoyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms (e.g., 2-pyridylcarbamoyl, 3-pyridylcarbamoyl, 4-pyridylcarbamoyl, 2-thienylcarbamoyl, 3-thienylcarbamoyl, etc.), and the like.

25 The substituents for the "C₆₋₁₄ aryloxy-carbonyl," "C₇₋₁₆ aralkyloxy-carbonyl," "C₃₋₈ cycloalkyl," "C₆₋₁₄ aryl," "C₇₋₁₆ aralkyl," "C₆₋₁₄ aryloxy," "C₇₋₁₆ aralkyloxy," "C₆₋₁₄ arylthio," "C₇₋₁₆ aralkylthio," "C₃₋₈ cycloalkyl-amino," "C₆₋₁₄ aryl-amino," "C₃₋₈ cycloalkyl-carbonyl," "C₆₋₁₄ aryl-carbonyl," "C₇₋₁₆ aralkyl-carbonyl," "5- to 7-membered heterocyclic carbonyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms," "C₆₋₁₄ arylsulfonyl," "C₆₋₁₄ arylsulfinyl," "C₃₋₈ cycloalkyl-carbonylamino," "C₆₋₁₄ aryl-carbonylamino," "C₆₋₁₄ arylsulfonylamino," "C₆₋₁₄ aryl-carbonyloxy" and "mono- or di-C₆₋₁₄ arylcarbamoyloxy" in the Substituent Group A include 1 to 5 substituents selected from, for example, a halogen atom, hydroxy, carboxy, nitro, cyano, the optionally substituted C₁₋₆ alkyl described above, the optionally substituted C₂₋₆ alkenyl described above, the optionally

substituted C₂₋₆ alkynyl described above, the optionally substituted C₃₋₈ cycloalkyl described above, the optionally substituted C₁₋₆ alkoxy described above, the optionally substituted C₁₋₆ alkylthio described above, the optionally substituted C₁₋₆ alkylsulfinyl described above, the optionally substituted C₁₋₆ alkylsulfonyl described above, the optionally esterified carboxyl described above, carbamoyl, thiocarbamoyl, a mono-C₁₋₆ alkylcarbamoyl, a di-C₁₋₆ alkylcarbamoyl, a mono- or di-C₆₋₁₄ arylcarbamoyl, a mono- or di-5- to 7-membered heterocyclic carbamoyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms, and the like.

The "optionally substituted heterocyclic group" in the Substituent Group A includes, for example, a 5- to 14-membered (monocyclic, bicyclic or tricyclic) heterocyclic group containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms, which may optionally be substituted with a halogen atom, hydroxy, carboxy, nitro, cyano, the optionally substituted C₁₋₆ alkyl described above, the optionally substituted C₂₋₆ alkenyl described above, the optionally substituted C₂₋₆ alkynyl described above, the optionally substituted C₃₋₈ cycloalkyl described above, the optionally substituted C₆₋₁₄ aryl described above, the optionally substituted C₁₋₆ alkoxy described above, the optionally substituted C₁₋₆ alkylthio described above, the optionally substituted C₆₋₁₄ arylthio described above, the optionally substituted C₇₋₁₆ aralkylthio described above, the optionally substituted C₁₋₆ alkylsulfinyl described above, the optionally substituted C₆₋₁₄ arylsulfinyl described above, the optionally substituted C₁₋₆ alkylsulfonyl described above, the optionally substituted C₆₋₁₄ arylsulfonyl described above, the optionally esterified carboxyl described above, carbamoyl, thiocarbamoyl, a mono-C₁₋₆ alkylcarbamoyl, a di-lower alkylcarbamoyl, a mono- or di-C₆₋₁₄ arylcarbamoyl, a mono- or di-5- or 7-membered heterocyclic carbamoyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms, or the like; preferably, (i) a 5- to 14-membered (preferably, 5- to 10-membered) aromatic heterocyclic group, (ii) a 5- to 10-membered non-aromatic heterocyclic group or (iii) a monovalent group formed by one optional hydrogen atom from 7- to 10-membered bridged-hetero ring, and among them, a 5-membered aromatic heterocyclic group is preferably used. Specifically used are, for example, an aromatic heterocyclic group such as thienyl (e.g., 2-thienyl, 3-thienyl), furyl (e.g., 2-furyl, 3-furyl), pyridyl (e.g., 2-pyridyl, 3-pyridyl, 4-pyridyl), thiazolyl (e.g., 2-thiazolyl, 4-thiazolyl, 5-thiazolyl), oxazolyl

(e.g., 2-oxazolyl, 4-oxazolyl), quinolyl (e.g., 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 8-quinolyl), isoquinolyl (e.g., 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl), pyrazinyl, pyrimidinyl (e.g., 2-pyrimidinyl, 4-pyrimidinyl), pyrrolyl (e.g., 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl), imidazolyl (e.g., 1-imidazolyl, 2-imidazolyl, 4-imidazolyl), pyrazolyl (e.g., 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl), pyridazinyl (e.g., 3-pyridazinyl, 4-pyridazinyl), isothiazolyl (e.g., 3-isothiazolyl), isoxazolyl (e.g., 3-isoxazolyl), indolyl (e.g., 1-indolyl, 2-indolyl, 3-indolyl), 2-benzothiazolyl, benzo[b]thienyl, (e.g., 2-benzo[b]thienyl, 3-benzo[b]thienyl), benzo[b]furanyl (e.g., 2-benzo[b]furanyl, 3-benzo[b]furanyl), etc.; a non-aromatic heterocyclic group such as pyrrolidinyl (e.g., 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl), oxazolidinyl (e.g., 2-oxazolidinyl), imidazolinyl (e.g., 1-imidazolinyl, 2-imidazolinyl, 4-imidazolinyl), piperidinyl (e.g., 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl), piperazinyl (e.g., 1-piperazinyl, 2-piperazinyl), morpholino, thiomorpholino, etc.

The "optionally substituted carbamoyl group" in the Substituent Group A includes a carbamoyl group, which may optionally be substituted with the optionally substituted C₁₋₆ alkyl, an optionally substituted C₂₋₆ alkenyl, an optionally substituted C₂₋₆ alkynyl, an optionally substituted C₃₋₈ cycloalkyl, an optionally substituted C₆₋₁₄ aryl, an optionally substituted heterocyclic group described above, etc., and specific examples include carbamoyl, thiocarbamoyl, a mono-C₁₋₆ alkylcarbamoyl (e.g., methylcarbamoyl, ethylcarbamoyl, etc.), di-C₁₋₆ alkylcarbamoyl (e.g., dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl, etc.), C₁₋₆ alkyl(C₁₋₆ alkoxy)carbamoyl (e.g., methyl(methoxy)carbamoyl, ethyl(methoxy)carbamoyl), a mono- or di-C₆₋₁₄ arylcarbamoyl (e.g., phenylcarbamoyl, 1-naphthylcarbamoyl, 2-naphthylcarbamoyl, etc.), a mono- or di-5- to 7-membered heterocyclic carbamoyl containing 1 to 4 hetero atoms of 1 or 2 species selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms (e.g., 2-pyridylcarbamoyl, 3-pyridylcarbamoyl, 4-pyridylcarbamoyl, 2-thienylcarbamoyl, 3-thienylcarbamoyl, etc.), a 5- to 7-membered cyclic carbamoyl (e.g., 1-pyrrolidinylcarbonyl, 1-piperidinylcarbonyl, hexamethyleneiminocarbonyl), and the like.

The "optionally substituted amino" in the Substituent Group A includes an amino, which may optionally be substituted with 1 or 2 groups selected from the optionally substituted C₁₋₆ alkyl described above, the optionally substituted C₂₋₆ alkenyl described above, the optionally substituted C₂₋₆ alkynyl described above, the optionally substituted C₃₋₈ cycloalkyl described above, the optionally substituted C₆₋₁₄ aryl described above, the optionally substituted C₁₋₆ alkoxy described above,

formyl, the optionally substituted C₁₋₆ alkyl-carbonyl described above, the optionally substituted C₃₋₈ cycloalkyl-carbonyl described above, the optionally substituted C₆₋₁₄ aryl-carbonyl described above, the optionally substituted C₁₋₆ alkoxy-carbonyl described above, the optionally substituted C₁₋₆ alkylsulfonyl described above, an
 5 optionally substituted C₆₋₁₄ arylsulfonyl), and the like.

More preferably, these substituents are a halogen atom, hydroxy, a C₁₋₆ alkoxy, an optionally halogenated C₁₋₆ alkyl, an optionally halogenated C₁₋₆ alkoxy, amino, nitro, cyano, etc.

XX10 preferably represents Phe, Trp, 2-naphthylalanine, 2-thienylalanine,
 10 tyrosine or 4-fluorophenylalanine, more preferably represents Phe or Trp, and most preferably represents Trp.

In a combination of these groups, there is preferably provided a metastatin derivative represented by formula:

15 XX0-XX2-XX3-XX4-XX5-XX6-AzaGly-XX8-XX9-XX10-NH₂

(wherein:

XX0 represents formyl, a C₁₋₆ alkanoyl or glycoloyl;

XX2 represents D-Tyr or the chemical bond;

20 XX3 represents Aze(2), Hyp, Gly, Aib, Leu, Lys, Glu, His(3Me), Tyr(PO₃H₂), Hyp(Bzl) or Pro(4F);

XX4 represents Asn or 2-amino-3-ureidopropionic acid;

XX5 represents Ser, Thr or Ala;

XX6 represents Phe, Cha, Phe(2F), Phe(3F), Phe(4F) or Phe(4Cl);

25 AzaGly represents azaglycine;

XX8 represents Leu or Ala(cPr);

XX9 represents Arg or Arg(Me); and,

XX10 represents Phe or Trp); or a salt thereof.

In a combination of these groups, there is more preferably provided a
 30 metastatin derivative represented by formula:

XX0-XX2-XX3-XX4-XX5-XX6-AzaGly-XX8-XX9-XX10-NH₂

(wherein:

35 XX0 represents acetyl or glycoloyl (preferably acetyl);

XX2 represents D-Tyr;

XX3 represents Hyp, Glu, Hyp(Bzl) or Pro(4F);

XX4 represents Asn or 2-amino-3-ureidopropionic acid;

XX5 represents Thr;

5 XX6 represents Phe, Cha, Phe(3F) or Phe(4F);

AzaGly represents azaglycine;

XX8 represents Leu or Ala(cPr);

XX9 represents Arg or Arg(Me); and,

XX10 represents Trp), or a salt thereof.

10 In the metastatin derivatives, all compounds wherein the groups shown by the respective symbols described above are optionally combined are preferably used. The compounds shown by the following compound numbers are also preferably used.

Compound No. 708: des(1)-Ac-[D-Tyr2,D-Trp3,Thr5,AzaGly7,D-Arg9,Trp10]MS10

15 Ac-D-Tyr-D-Trp-Asn-Thr-Phe-AzaGly-Leu-D-Arg-Trp-NH₂

Compound No. 709: des(1-3)-Ac-[Thr5,AzaGly7,Arg(Me)9,Trp10]MS10

Ac-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 710: des(1-3)-Decanoyl-[Thr5,AzaGly7,Arg(Me)9,Trp10]MS10

Decanoyl-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

20 Compound No. 712: des(1-2)-[Acp3, Thr5,AzaGly7,Arg(Me)9,Trp10]MS10

Acp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 713: des(1-2)-Ac-[Acp3, Thr5,AzaGly7,Arg(Me)9,Trp10]MS10

Ac-Acp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 714:

25 des(1)-Ac-[D-Tyr2,D-Trp3,Asp(NHPen)4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10

Ac-D-Tyr-D-Trp-Asp(NHPen)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 715:

des(1)-Ac-[D-Tyr2,D-Trp3,Asp(NHcPr)4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10

Ac-D-Tyr-D-Trp-Asp(NHcPr)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

30 Compound No. 716:

des(1)-Ac-[D-Tyr2,D-Trp3,Asp(NHBzl)4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10

Ac-D-Tyr-D-Trp-Asp(NHBzl)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 717:

des(1)-Ac-[D-Tyr2,D-Trp3,Alb4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10

35 Ac-D-Tyr-D-Trp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

- Compound No. 718:
des(1)-Ac-[D-Tyr2,D-Pya(4)3,Alb4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-D-Pya(4)-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 719:
5 des(1)-Ac-[D-Tyr2,D-Trp3,D-Pro5,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-D-Trp-Asn-D-Pro-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 720:
des(1)-Ac-[D-Tyr2,Aze(2)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Aze(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 10 Compound No. 721:
des(1)-Ac-[D-Tyr2,Pic(2)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Pic(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 722:
des(1)-Ac-[D-Tyr2,Pic(3)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
15 Ac-D-Tyr-Pic(3)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 723:
des(1)-Ac-[D-Tyr2,Hyp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 724: des(1)-Ac-[D-Tyr2,Thz3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
20 Ac-D-Tyr-Thz-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 725:
des(1)-Ac-[D-Tyr2,NMeAla3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-NMeAla-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 726: des(1)-Ac-[D-Tyr2,Gly3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
25 Ac-D-Tyr-Gly-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 727: des(1)-Ac-[D-Tyr2,Aib3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Aib-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 728:
des(1)-Ac-[D-Tyr2,Abz(2)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
30 Ac-D-Tyr-Abz(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 730:
des(1)-Ac-[D-Tyr2,Aze(3)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Aze(3)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 731: des(1)-Ac-[D-Tyr2,Sar3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
35 Ac-D-Tyr-Sar-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

- Compound No. 732:
 des(1)-Ac-[D-Tyr2,D-NMeAla3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-D-NMeAla-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 734: des(1)-Ac-[D-Tyr2,Izc3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 5 Ac-D-Tyr-Izc-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 735:
 des(1)-Ac-[D-Tyr2,D-Asp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-D-Asp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 736:
 10 des(1)-Ac-[D-Tyr2,D-Dap3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-D-Dap-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 737:
 des(1)-Ac-[D-Tyr2,D-Ser3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-D-Ser-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 15 Compound No. 738:
 des(1)-Ac-[D-Tyr2,D-Gln3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-D-Gln-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 739:
 des(1)-Ac-[D-Tyr2,D-His3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 20 Ac-D-Tyr-D-His-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 740:
 des(1)-Ac-[D-Tyr2,D-Trp3,Dab4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-D-Trp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 742: des(1)-Ac-[D-Tyr2,Ala3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 25 Ac-D-Tyr-Ala-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 743: des(1)-Ac-[D-Tyr2,Leu3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Leu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 744: des(1)-Ac-[D-Tyr2,Ser3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Ser-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 30 Compound No. 745: des(1)-Ac-[D-Tyr2,Lys3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Lys-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 746: des(1)-Ac-[D-Tyr2,Glu3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Glu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 747: des(1)-Ac-[D-Tyr2, β-Ala3,Thr5,AzaGly7,Arg(Me)9,Trp10]
 35 MS10

- Ac-D-Tyr-β-Ala-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
Compound No. 748:
des(1)-Ac-[D-Tyr2,D-Trp3,Thr5,Phe(4Cl)6,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 5 Compound No. 749:
des(1)-Ac-[D-Tyr2,D-Trp3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
Compound No. 750:
des(1)-Ac-[D-Tyr2,D-Trp3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10
- 10 Ac-D-Tyr-D-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
Compound No. 754:
des(1)-Ac-[D-Tyr2,Lys3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Lys-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
Compound No. 755:
- 15 des(1)-Ac-[D-Tyr2,Glu3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Glu-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
Compound No. 756:
des(1)-Ac-[D-Tyr2,Lys3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Lys-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 20 Compound No. 757:
des(1)-Ac-[D-Tyr2,Glu3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Glu-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
Compound No. 758:
des(1)-Ac-[D-Tyr2,Lys3,Thr5,Phe(4Cl)6,AzaGly7,Arg(Me)9,Trp10]MS10
- 25 Ac-D-Tyr-Lys-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
Compound No. 759:
des(1)-Ac-[D-Tyr2,Glu3,Thr5,Phe(4Cl)6,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Glu-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
Compound No. 760:
- 30 des(1)-Ac-[D-Tyr2,Pzc(2)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Pzc(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
Compound No. 763:
des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10
Ac-D-Tyr-Hyp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 35 Compound No. 764:

- des(1)-Ac-[D-Tyr2,Trp3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 765:
- des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 5 Ac-D-Tyr-Hyp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 766:
- des(1)-Ac-[D-Tyr2,Trp3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 767:
- 10 des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(4Cl)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 768:
- des(1)-Ac-[D-Tyr2,Trp3,Thr5,Phe(4Cl)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 15 Compound No. 769:
- des(1)-Ac-[D-Tyr2,Gly3,Thr5,Phe(4Cl)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Gly-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 770:
- des(1)-Ac-[D-Tyr2,Aib3,Thr5,Phe(4Cl)6,AzaGly7,Arg(Me)9,Trp10]MS10
 20 Ac-D-Tyr-Aib-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 771: des(1)-Ac-[D-Tyr2,Orn3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Orn-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 772: des(1)-Ac-[D-Tyr2,Thr3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Thr-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 25 Compound No. 773:
- des(1)-Ac-[D-Tyr2,His(3Me)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-His(3Me)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 774:
- des(1)-Ac-[D-Tyr2,DL-Ala(Pip)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 30 Ac-D-Tyr-DL-Ala(Pip)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 775:
- des(1)-Ac-[D-Tyr2,Tyr(PO₃H₂)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Tyr(PO₃H₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 776:
- 35 des(1)-Glycoloyl-[D-Tyr2,Hyp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10

- Glycoloyl-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 777: des(1-2)-Ac-[D-Tyr3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 780:
 5 des(1)-Ac-[D-Tyr2,Pro(4NH₂)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Pro(4NH₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 781:
 des(1)-Ac-[D-Tyr2,Hyp(Bzl)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp(Bzl)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 10 Compound No. 782:
 des(1)-Ac-[D-Tyr2,D-NMePhe3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-D-NMePhe-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 783:
 des(1)-Ac-[D-Tyr2,Gly3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 15 Ac-D-Tyr-Gly-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 784:
 des(1)-Ac-[D-Tyr2,Aib3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Aib-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 785:
 20 des(1)-Ac-[D-Tyr2,Gly3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Gly-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 786:
 des(1)-Ac-[D-Tyr2,Aib3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Aib-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 25 Compound No. 787:
 des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(4F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 788:
 des(1)-Ac-[D-Tyr2,Glu3,Thr5,Phe(4F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 30 Ac-D-Tyr-Glu-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 789:
 des(1)-Ac-[D-Tyr2,Lys3,Thr5,Phe(4F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Lys-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 790:
 35 des(1)-Ac-[D-Tyr2,Gly3,Thr5,Phe(4F)6,AzaGly7,Arg(Me)9,Trp10]MS10

- Ac-D-Tyr-Gly-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 791:
 des(1)-Ac-[D-Tyr2,Aib3,Thr5,Phe(4F)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Aib-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 5 Compound No. 794:
 des(1)-Ac-[D-Tyr2,Hyp3,Thr5,D-Phe6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-D-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 797: des(1)-Ac-[D-Tyr2,Hyp3,Thr5,AzaGly7,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg-Trp-NH₂
- 10 Compound No. 800:
 des(1)-Ac-[D-Tyr2,Hyp3,Alb4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 801:
 des(1-5)-4-[Bis-(2-Pyridylmethyl)aminomethyl]benzoyl-[AzaGly7,Arg(Me)9,Trp10]
- 15 MS10
 4-[Bis-(2-Pyridylmethyl)aminomethyl]benzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 809:
 des(1)-Ac-[D-Tyr2,Hyp3,NMeSer5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-NMeSer-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 20 Compound No. 810:
 des(1)-Ac-[D-Tyr2,Hyp3,Hyp5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Hyp-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 813:
 des(1)-Ac-[D-Tyr2,Hyp3,Gly5,AzaGly7,Arg(Me)9,Trp10]MS10
- 25 Ac-D-Tyr-Hyp-Asn-Gly-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 814:
 des(1)-Ac-[D-Tyr2,Hyp3,Ala5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 815:
 des(1)-Ac-[D-Tyr2,Hyp3,D-Ala5,AzaGly7,Arg(Me)9,Trp10]MS10
- 30 Ac-D-Tyr-Hyp-Asn-D-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 816:
 des(1)-Ac-[D-Tyr2,Hyp3,His4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-His-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 35 Compound No. 843:

- des(1)-Ac-[D-Tyr2,Hyp3,Gln4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Gln-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 844:
- des(1)-Ac-[D-Tyr2,Hyp3,D-Asn4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 5 Ac-D-Tyr-Hyp-D-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 845:
- des(1)-Ac-[D-Tyr2,Hyp3,Cit4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Cit-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 846:
- 10 des(1)-Ac-[D-Tyr2,Hyp3,D-Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-D-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 856:
- des(1)-Ac-[D-Tyr2,Hyp3,Thr5,AzaGly7,Ala(cPr)8,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Ala(cPr)-Arg(Me)-Trp-NH₂
- 15 Compound No. 860: des(1-5)-4-Ureidobenzoyl-[AzaGly7,Arg(Me)9,Trp10]MS10
 4-Ureidobenzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 861:
- des(1)-Ac-[D-Tyr2,Hyp3,Arg4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Arg-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 20 Compound No. 862:
- des(1)-Ac-[D-Tyr2,Hyp3,Gly4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Gly-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 863:
- des(1)-Ac-[D-Tyr2,Hyp3,Dap4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 25 Ac-D-Tyr-Hyp-Dap-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 864:
- des(1)-Ac-[D-Tyr2,Hyp3,Dab4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
- Compound No. 868: des(1)-Ac-[D-Tyr2,Hyp3,Thr5, αMePhe6,AzaGly7,Arg(Me)9,
 30 Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-αMePhe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 870:
- des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(2Me)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-Phe(2Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂
- 35 Compound No. 872:

- des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(3Me)6,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-Phe(3Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 874:
- des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(4Me)6,AzaGly7,Arg(Me)9,Trp10]MS10
 5 Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 877:
- des(1)-Ac-[D-Tyr2,Hyp3,Thr5,threo-Ser(3Phenyl)6,AzaGly7,Arg(Me)9,Trp10]MS10
 0 Ac-D-Tyr-Hyp-Asn-Thr-threo-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 10 Compound No. 882:
- des(1)-Ac-[D-Tyr2,Hyp3,Thr5,erythro-Ser(3Phenyl)6,AzaGly7,Arg(Me)9,Trp10]MS10
 10 Ac-D-Tyr-Hyp-Asn-Thr-erythro-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 886:
- 15 des(1)-Ac-[D-Tyr2,Hyp3,Nva4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-Hyp-Nva-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 887: des(1-2)-Ac-[Hyp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 888:
- 20 des(1-2)-3-(p-Hydroxyphenyl)propionyl-[Hyp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 10 3-(p-Hydroxyphenyl)propionyl-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 889: des(1-2)-[pGlu3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 pGlu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 25 Compound No. 896:
- des(1)-Ac-[D-Tyr2,cisHyp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-D-Tyr-cisHyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 897:
- des(1)-Ac-[D-Tyr2,Pro(4F)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 30 Ac-D-Tyr-Pro(4F)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 899: des(1)-Ac-[Tyr2,Hyp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10
 Ac-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

In particular, the compounds represented by the following compound numbers are preferred as the metastatin derivatives.

- 35 Ac-D-Tyr-D-Trp-Asp(NH₂Pen)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound

- No. 714),
Ac-D-Tyr-D-Trp-Asp(NHcPr)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 715),
Ac-D-Tyr-D-Trp-Asp(NHBzl)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 716),
5 Ac-D-Tyr-D-Trp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 717),
Ac-D-Tyr-D-Pya(4)-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 718),
10 Ac-D-Tyr-Aze(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 720),
Ac-D-Tyr-Pic(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 721),
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 723),
15 Ac-D-Tyr-Thz-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 724),
Ac-D-Tyr-Gly-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 726)
Ac-D-Tyr-Aib-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 727),
Ac-D-Tyr-D-NMeAla-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 732),
20 Ac-D-Tyr-D-Gln-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 738),
Ac-D-Tyr-D-His-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 739),
Ac-D-Tyr-D-Trp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 740),
25 Ac-D-Tyr-Ala-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 742),
Ac-D-Tyr-Leu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 743),
Ac-D-Tyr-Ser-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 744),
Ac-D-Tyr-Lys-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 745),
30 Ac-D-Tyr-Glu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 746),
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 748),
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 749),
35 Ac-D-Tyr-D-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.

- 750),
Ac-D-Tyr-Lys-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 754),
Ac-D-Tyr-Glu-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 755),
5 Ac-D-Tyr-Lys-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 756),
Ac-D-Tyr-Glu-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 757),
10 Ac-D-Tyr-Lys-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 758),
Ac-D-Tyr-Glu-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 759),
Ac-D-Tyr-Pzc(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 760),
15 Ac-D-Tyr-Hyp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 763),
Ac-D-Tyr-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 764),
20 Ac-D-Tyr-Hyp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 765),
Ac-D-Tyr-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 766),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 767),
25 Ac-D-Tyr-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 768),
Ac-D-Tyr-Gly-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 769),
30 Ac-D-Tyr-Aib-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 770),
Ac-D-Tyr-Orn-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 771),
Ac-D-Tyr-Thr-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 772),
Ac-D-Tyr-His(3Me)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 773),
35

- Ac-D-Tyr-Tyr(PO₃H₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 775),
Glycoloyl-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 776),
5 Ac-D-Tyr-Pro(4NH₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 780),
Ac-D-Tyr-Hyp(Bzl)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 781),
Ac-D-Tyr-D-NMePhe-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 782),
10 Ac-D-Tyr-Gly-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 783),
Ac-D-Tyr-Aib-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 784),
15 Ac-D-Tyr-Gly-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 785),
Ac-D-Tyr-Aib-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 786),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 787),
20 Ac-D-Tyr-Glu-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 788),
Ac-D-Tyr-Lys-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 789),
25 Ac-D-Tyr-Gly-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 790),
Ac-D-Tyr-Aib-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 791),
Ac-D-Tyr-Hyp-Asn-Thr-D-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 794),
30 Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg-Trp-NH₂ (Compound No. 797),
Ac-D-Tyr-Hyp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 800),
4-[Bis-(2-Pyridylmethyl)aminomethyl]benzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 801),
35 Ac-D-Tyr-Hyp-Asn-NMeSer-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.

- 809),
 Ac-D-Tyr-Hyp-Asn-Hyp-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 810),
 Ac-D-Tyr-Hyp-Asn-Gly-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 813),
 Ac-D-Tyr-Hyp-Asn-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 814),
 5 Ac-D-Tyr-Hyp-Asn-D-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 815),
 Ac-D-Tyr-Hyp-His-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 816),
 Ac-D-Tyr-Hyp-Gln-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 843),
 Ac-D-Tyr-Hyp-D-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.
 10 844),
 Ac-D-Tyr-Hyp-Cit-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 845),
 Ac-D-Tyr-Hyp-Asn-D-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 846),
 Compound No. 856:
 15 des(1)-Ac-[D-Tyr₂,Hyp₃,Thr₅,AzaGly₇,Ala(cPr)₈,Arg(Me)₉,Trp₁₀]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Ala(cPr)-Arg(Me)-Trp-NH₂
 Compound No. 860: des(1-5)-4-Ureidobenzoyl-[AzaGly₇,Arg(Me)₉,Trp₁₀]MS10
 4-Ureidobenzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 861:
 20 des(1)-Ac-[D-Tyr₂,Hyp₃,Arg₄,Thr₅,AzaGly₇,Arg(Me)₉,Trp₁₀]MS10
 Ac-D-Tyr-Hyp-Arg-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 862:
 des(1)-Ac-[D-Tyr₂,Hyp₃,Gly₄,Thr₅,AzaGly₇,Arg(Me)₉,Trp₁₀]MS10
 Ac-D-Tyr-Hyp-Gly-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 25 Compound No. 863:
 des(1)-Ac-[D-Tyr₂,Hyp₃,Dap₄,Thr₅,AzaGly₇,Arg(Me)₉,Trp₁₀]MS10
 Ac-D-Tyr-Hyp-Dap-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 864:
 des(1)-Ac-[D-Tyr₂,Hyp₃,Dab₄,Thr₅,AzaGly₇,Arg(Me)₉,Trp₁₀]MS10
 30 Ac-D-Tyr-Hyp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 868: des(1)-Ac-[D-Tyr₂,Hyp₃,Thr₅,αMePhe₆,AzaGly₇,Arg(Me)₉,
 Trp₁₀]MS10
 Ac-D-Tyr-Hyp-Asn-Thr-αMePhe-AzaGly-Leu-Arg(Me)-Trp-NH₂
 Compound No. 870:
 35 des(1)-Ac-[D-Tyr₂,Hyp₃,Thr₅,Phe(2Me)₆,AzaGly₇,Arg(Me)₉,Trp₁₀]MS10

Ac-D-Tyr-Hyp-Asn-Thr-Phe(2Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 872:

des(1)-Ac-[D-Tyr₂,Hyp₃,Thr₅,Phe(3Me)₆,AzaGly₇,Arg(Me)₉,Trp₁₀]MS₁₀

Ac-D-Tyr-Hyp-Asn-Thr-Phe(3Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂

5 Compound No. 874:

des(1)-Ac-[D-Tyr₂,Hyp₃,Thr₅,Phe(4Me)₆,AzaGly₇,Arg(Me)₉,Trp₁₀]MS₁₀

Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 877:

des(1)-Ac-[D-Tyr₂,Hyp₃,Thr₅,threo-Ser(3Phenyl)₆,AzaGly₇,Arg(Me)₉,Trp₁₀]MS₁₀

10 0

Ac-D-Tyr-Hyp-Asn-Thr-threo-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 882:

des(1)-Ac-[D-Tyr₂,Hyp₃,Thr₅,erythro-Ser(3Phenyl)₆,AzaGly₇,Arg(Me)₉,Trp₁₀]MS₁₀

10

15 Ac-D-Tyr-Hyp-Asn-Thr-erythro-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 886:

des(1)-Ac-[D-Tyr₂,Hyp₃,Nva₄,Thr₅,AzaGly₇,Arg(Me)₉,Trp₁₀]MS₁₀

Ac-D-Tyr-Hyp-Nva-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 888:

20 des(1-2)-3-(p-Hydroxyphenyl)propionyl-[Hyp₃,Thr₅,AzaGly₇,Arg(Me)₉,Trp₁₀]MS₁₀

10

3-(p-Hydroxyphenyl)propionyl-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 896:

des(1)-Ac-[D-Tyr₂,cisHyp₃,Thr₅,AzaGly₇,Arg(Me)₉,Trp₁₀]MS₁₀

25 Ac-D-Tyr-cisHyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

Compound No. 897:

des(1)-Ac-[D-Tyr₂,Pro(4F)₃,Thr₅,AzaGly₇,Arg(Me)₉,Trp₁₀]MS₁₀

Ac-D-Tyr-Pro(4F)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂

or

30 Compound No. 899: des(1)-Ac-[Tyr₂,Hyp₃,Thr₅,AzaGly₇,Arg(Me)₉,Trp₁₀]MS₁₀

Ac-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂.

The metastin derivatives of the present invention or their salts or prodrugs have excellent blood stability, solubility, etc., in addition to the excellent effects of suppressing cancer metastasis and cancer growth, and are useful as agents for

preventing or treating cancers (for example, lung cancer, gastric cancer, liver cancer, pancreatic cancer, colorectal cancer, rectal cancer, colonic cancer, prostate cancer, ovarian cancer, cervical cancer, breast cancer, etc.). The metastin derivatives of the present invention or their salts or prodrugs have the effect of controlling pancreatic function and are useful as agents for preventing or treating pancreatic diseases (e.g.,
5 acute or chronic pancreatitis, pancreatic cancer, etc.). The metastin derivatives of the present invention or their salts or prodrugs have the effect of controlling placental function and are useful as agents for preventing or treating choriocarcinoma, hydatid mole, invasive mole, miscarriage, fetal hypoplasia, abnormal glucose metabolism,
10 abnormal lipid metabolism or induction of delivery.

Also, the metastin derivatives of the present invention or their salts or prodrugs have the effects of increasing sugar level, promoting pancreatic glucagon secretion and promoting urine formation, and are useful as agents for preventing or treating obesity, hyperlipemia, type II diabetes mellitus, hypoglycemia, hypertension,
15 diabetic neuropathy, diabetic nephropathy, diabetic retinopathy, edema, urinary disturbances, insulin resistance, unstable diabetes mellitus, fatty atrophy, insulin allergy, insulinoma, arteriosclerosis, thrombotic disorders or lipotoxicity.

In addition, the metastin derivatives of the present invention or their salts or prodrugs have excellent activities of promoting gonadotropic hormone secretion, promoting sex hormone secretion, inducing ovulation or stimulating ovulation, and are useful as low toxic and stable agents, e.g., agents for improving gonadal function, agents for preventing or treating hormone-dependent cancer (e.g., prostate cancer, breast cancer, etc.), infertility, endometriosis, early puberty, myoma of the uterus, etc., agents for inducing or stimulating ovulation, gonadotropic hormone secretagogue
25 agents, contraceptives, sex hormone secretagogue agents, or the like.

Furthermore, the metastin derivatives of the present invention or their salts or prodrugs are useful as agents for preventing or treating Alzheimer's disease, Autism, moderate cognitive impairment, etc.

The metastin derivatives of the present invention or their salts or prodrugs
30 are useful as agents for suppressing gonadotropic hormone secretion or sex hormone secretion; down-regulating agents for gonadotropic hormone or sex hormone; down-regulating agents for human OT7T175 (metastin receptor) protein consisting of the amino acid sequence represented by SEQ ID NO: 9; agents for preventing or treating hormone-dependent cancers (e.g., prostate cancer, breast cancer, etc.;
35 particularly, hormone-sensitive prostate cancer, hormone-sensitive prostate cancer,

etc.); agents for preventing or treating endometriosis; agents for inhibiting ovarian follicular maturation; menstrual cycle-suspending agents; agents for treating myoma of the uterus; agents for treating early puberty; contraceptives, etc.

In addition, the metastin derivatives of the present invention or their salts or prodrugs are useful as agents for potentiating immunity (eg., prophylactic agents for infection after bone-marrow transplant, agents for potentiating immunity intended for cancer, etc); immunostimulator (eg., regeneration of the thymus, regrowth of the thymus, enhancement of T cell development, etc); agents for preventing or treating bulbospinal muscular atrophy; agents for protecting ovary; agents for preventing or treating benign prostate hypertrophy (BPH); agents for preventing or treating gender identity disorder; or agents for in vitro fertilization (IVF). In addition, they are also useful as agents for preventing or treating infertility, hypogonadism, oligospermia, azoospermia, aspermia, asthenospermia, or necrospermia. Further, they are useful for hormone-dependent diseases (e.g., sex hormone dependent cancer such as prostate cancer, uterine cancer, breast cancer, hypophysial tumor, etc.), prostate gland enlargement, endometriosis, uterine fibroid, early puberty, dysmenorrhea, amenorrhea, menstrual syndrome, multilocular ovary syndrome, postoperative relapse of the above-mentioned cancers, metastasis of the above-mentioned cancers, hypopituitarism, dwarfism (the case where the secretion of growth hormone was compromised with hyposecretion of pituitary hormone, etc.), menopausal disorder, indefinite complaint, sex hormone dependent disorders such as calcium phosphor bone metabolic disorders. It is also applicable for contraception (or infertility when rebound effects after cessation of the drug are utilized), etc.

Furthermore, metastin per se or DNA encoding metastin, etc. are also useful as agents for suppressing gonadotropic hormone secretion or sex hormone secretion; down-regulating agents for gonadotropic hormone or sex hormone; down-regulating agents for human OT7T175 (metastin receptor) protein consisting of the amino acid sequence represented by SEQ ID NO: 9; agents for preventing or treating hormone-dependent cancers (e.g., prostate cancer, breast cancer, etc.; particularly, hormone-sensitive prostate cancer, hormone-sensitive breast cancer, etc.); agents for preventing or treating endometriosis; agents for inhibiting ovarian follicular maturation; menstrual cycle-suspending agents; agents for treating myoma of the uterus; agents for treating early puberty; contraceptives, etc.

The metastin derivatives of the present invention can be prepared by

publicly known methods for peptide synthesis. As the methods for peptide synthesis, for example, either solid phase synthesis or liquid phase synthesis may be used. That is, the partial peptide or amino acids that can constitute the peptide of the present invention are repeatedly condensed with the remaining part to give the product having a desired sequence. Where the product has protecting groups, these protecting groups are removed to give the desired peptide. Publicly known methods for condensation and removal of the protecting groups are described in (1) to (5) below.

(1) M. Bodanszky & M.A. Ondetti: Peptide Synthesis, Interscience Publishers, New York (1966)

(2) Schroeder & Luebke: The Peptide, Academic Press, New York (1965)

(3) Nobuo Izumiya, et al.: *Peptide Gosei-no-Kiso to Jikken* (Basics and experiments of peptide synthesis), published by Maruzen Co. (1975)

(4) Haruaki Yajima & Shunpei Sakakibara: *Seikagaku Jikken Kozà* (Biochemical Experiment) 1, *Tanpakushitsu no Kagaku* (Chemistry of Proteins) IV, 205 (1977)

(5) Haruaki Yajima, ed.: *Zoku Iyakuhin no Kaihatsu* (A sequel to Development of Pharmaceuticals), Vol. 14, Peptide Synthesis, published by Hirokawa Shoten

After completion of the reaction, the product may be purified and isolated by a combination of conventional purification methods such as solvent extraction, distillation, column chromatography, liquid chromatography and recrystallization to give the peptide of the present invention. When the peptide obtained by the above methods is in a free form, the peptide can be converted into an appropriate salt by a publicly known method; conversely when the peptide is obtained in a salt form, it can be converted into its free form by publicly known methods.

For condensation of the protected amino acids or peptides, a variety of activation reagents for peptide synthesis may be used, but trisphosphonium salts, tetramethyluronium salts, carbodiimides, etc. are particularly preferred. Examples of trisphosphonium salts include benzotriazol-1-yloxytris(pyrrolidino)phosphonium hexafluorophosphate (PyBOP), bromotris(pyrrolidino) phosphonium hexafluorophosphate (PyBroP) and

7-azabenzotriazol-1-yloxytris(pyrrolidino)phosphonium hexafluorophosphate (PyAOP), examples of tetramethyluronium salts include

2-(1H-benzotriazol-1-yl)-1,1,3,3-hexafluorophosphate (HBTU),

2-(7-azabenzotriazol-1-yl)-1,1,3,3-hexafluorophosphate (HATU),

2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TBTU),

2-(5-norbornene-2,3-dicarboxyimido)-1,1,3,3-tetramethyluronium tetrafluoroborate (TNTU) and O-(N-succimidyl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TSTU); examples of carbodiimides include DCC, N,N'-diisopropylcarbodiimide (DIPCDI) and N-ethyl-N'-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDCI.HCl); etc. For condensation using these reagents, the addition of racemization inhibitors (e.g., HONB, HOBt, HOAt, HOObt, etc.) is preferred. Solvents used in condensation may be appropriately chosen from solvents that are known to be usable for condensation. For example, acid amides such as anhydrous or hydrous N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone, etc., halogenated hydrocarbons such as methylene chloride, chloroform, etc., alcohols such as trifluoroethanol, phenol, etc., sulfoxides such as dimethyl sulfoxide, etc., tertiary amines such as pyridine, etc., ethers such as dioxane, tetrahydrofuran, etc., nitriles such as acetonitrile, propionitrile, etc., esters such as methyl acetate, ethyl acetate, etc., or suitable mixtures thereof, etc. are used. The reaction temperature is appropriately chosen from the range known to be applicable to peptide binding reactions and is normally suitably chosen from the range of about -20°C to 50°C. The activated amino acid derivatives are used generally in 1.5 to 6 times excess. In the case of solid phase synthesis, the condensation is examined using the ninhydrin reaction; when the condensation is insufficient, the condensation can be completed by repeating the condensation reaction without removal of the protecting groups. When the condensation is yet insufficient even after repeating the reaction, the unreacted amino acids are acylated with acetic anhydride or acetylimidazole to cancel any adverse effect on the subsequent reaction.

Examples of the protecting groups used to protect amino groups in the starting amino acids include Z, Boc, tert-pentyloxycarbonyl, isobornyloxycarbonyl, 4-methoxybenzyloxycarbonyl, Cl-Z, Br-Z, adamantyloxycarbonyl, trifluoroacetyl, phthaloyl, formyl, 2-nitrophenylsulphenyl, diphenylphosphinothioyl, Fmoc, trityl, etc. Examples of protecting groups for a carboxyl group include, in addition to the C₁₋₆ alkyl group, C₃₋₈ cycloalkyl group and C₇₋₁₄ aralkyl group for R described above, allyl, 2-adamantyl, 4-nitrobenzyl, 4-methoxybenzyl, 4-chlorobenzyl, phenacyl group, benzyloxycarbonylhydrazide, tert-butoxycarbonylhydrazide, tritylhydrazide, etc.

The hydroxyl group of serine and threonine can be protected, for example, by esterification or etherification. Examples of groups suitable for this esterification include a group derived from organic acid such as a lower (C₂₋₄) alkanoyl group such as acetyl group, an aroyl group such as benzoyl group, etc. Examples of a group

suitable for the etherification include benzyl group, tetrahydropyranyl group, tert-butyl group, trytyl group (Trt), etc.

Examples of groups for protecting the phenolic hydroxyl group of tyrosine include Bzl, 2,6-dichlorobenzyl (Cl₂-Bzl), 2-nitrobenzyl, Br-Z, tert-butyl, etc.

5 Examples of groups used to protect the imidazole moiety of histidine include Tos, 4-methoxy-2,3,6-trimethylbenzenesulfonyl (Mtr), DNP, Bom, Bum, Boc, Trt, Fmoc, etc.

10 Examples of protecting groups for the guanidino group in arginine include Tos, Z, 4-methoxy-2,3,6-trimethylbenzenesulfonyl (Mtr), p-methoxybenzenesulfonyl (MBS), 2,2,5,7,8-pentamethylchroman-6-sulfonyl (Pmc), mesitylene-2-sulfonyl (Mts), 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl (Pbf), Boc, Z, NO₂, etc.

 Examples of protecting groups for side chain amino group of lysine include Z, Cl-Z, trifluoroacetyl, Boc, Fmoc, Trt, Mtr, 4,4-dimethyl-2,6-dioxocyclohexylideneyl (Dde), etc.

15 Examples of protecting groups for the indolyl of tryptophan include formyl (For), Z, Boc, Mts, Mtr, etc.

 A protecting group for asparagine and glutamine include Trt, xanthyl (Xan), 4,4'-dimethoxybenzhydryl (Mbh), 2,4,6-trimethoxybenzyl (Tmob), etc.

20 Examples of the activated carboxyl groups in the starting material include the corresponding acid anhydrides, azides, activated esters [esters with alcohols (e.g., pentachlorophenol, 2,4,5-trichlorophenol, 2,4-dinitrophenol, cyanomethyl alcohol, p-nitrophenol, HONB, N-hydroxysuccinimide, 1-hydroxybenzotriazole (HOBt) or 1-hydroxy-7-azabenzotriazole (HOAt)], etc. As the amino acids in which the amino groups in the starting material are activated, the corresponding phosphorous amides
25 are employed.

 To eliminate (split off) the protecting groups, there are used catalytic reduction under hydrogen gas flow in the presence of a catalyst such as Pd-black or Pd-carbon; an acid treatment with anhydrous hydrogen fluoride, methanesulfonic acid, trifluoromethanesulfonic acid, trifluoroacetic acid, trimethylsilane bromide
30 (TMSBr), trimethylsilyl trifluoromethanesulfonate, tetrafluoroboric acid, tris(trifluoro)boron, boron tribromide or a mixed solution thereof, a base treatment with diisopropylethylamine, triethylamine, piperidine, piperazine, etc., and reduction with sodium in liquid ammonia. The elimination of protecting groups by the acid treatment described above is carried out generally at a temperature of approximately
35 -20°C to 40°C. In the acid treatment, it is efficient to add a cation scavenger such

as anisole, phenol, thioanisole, m-cresol, p-cresol, etc., dimethylsulfide, 1,4-butanedithiol, 1,2-ethanedithiol, etc. Furthermore, 2,4-dinitrophenyl group used as the protecting group for the imidazole of histidine is removed by a treatment with thiophenol. Formyl group used as the protecting group of the indole of tryptophan
5 is removed by the aforesaid acid treatment in the presence of 1,2-ethanedithiol, 1,4-butanedithiol, etc. as well as by a treatment with an alkali such as a dilute sodium hydroxide solution, dilute ammonia, etc.

Protection of functional groups that should not be involved in the reaction of the starting materials, protecting groups, removal of the protecting groups and
10 activation of functional groups involved in the reaction may be appropriately chosen from publicly known groups and publicly known means.

Methods for obtaining the amide of the peptide include, for example, solid phase synthesis using resins for the formation of peptide amide. In another method for obtaining the amides of the peptide, for example, the α -carboxyl group of the
15 carboxy terminal amino acid is first protected by amidation; the peptide chain is then extended from the amino group side to a desired length. Thereafter, a peptide in which only the protecting group of the N-terminal α -amino group in the peptide chain has been removed from the peptide and a peptide (or an amino acid) in which only the protecting group of the C-terminal carboxyl group has been eliminated are
20 prepared. The two peptides are condensed in a mixture of the solvents described above. The details of the condensation reaction are the same as described above. After the protected peptide obtained by the condensation is purified, all the protecting groups are removed by the method described above to give the desired crude peptide. This crude peptide is purified by various known purification means.
25 Lyophilization of the major fraction gives the amide of the desired peptide.

When the metastin derivative of the present invention is present in the form of a configurational isomer, a diastereomer, a conformer, or the like, each can be isolated by the separating and purifying means described above, if desired. In
30 addition, when the compound of the present invention is racemic, it can be separated into an S isomer and an R isomer by the conventional optical resolving means.

When steric isomers exist for the metastin derivative of the present invention, the present invention includes both of these isomers alone and the isomers present as a mixture thereof.

In addition, the metastin derivative of the present invention may also be
35 hydrated or non-hydrated.

The metastin derivative of the present invention may also be labeled with an isotope (e.g., ^3H , ^{14}C , ^{35}S), etc.

Throughout the specification, the peptides are represented in accordance with the conventional way of describing peptides, that is, the N-terminus (amino terminus) at the left hand and the C-terminus (carboxyl terminus) at the right hand. In the peptides, the C-terminus is usually in the form of an amide ($-\text{CONH}_2$), a carboxyl group ($-\text{COOH}$), a carboxylate ($-\text{COO}^-$), an alkylamide ($-\text{CONHR}$) or an ester ($-\text{COOR}$) and the amide ($-\text{CONH}_2$) is particularly preferred. Examples of R in the ester or alkylamide include a C_{1-6} alkyl group such as methyl, ethyl, n-propyl, isopropyl, n-butyl, etc.; a C_{3-8} cycloalkyl group such as cyclopentyl, cyclohexyl, etc.; a C_{6-12} aryl group such as phenyl, α -naphthyl, etc.; a C_{7-14} aralkyl group such as a phenyl- C_{1-2} -alkyl group, e.g., benzyl, phenethyl, etc., or an α -naphthyl- C_{1-2} -alkyl group such as α -naphthylmethyl, etc.; pivaloyloxymethyl group, which are widely used as an ester for oral use, and the like.

Examples of a salt of the metastin derivative of the present invention include a metal salt, an ammonium salt, a salt with an organic base, a salt with inorganic acid, a salt with organic acid, a salt with basic or acidic amino acid, and the like. Preferred examples of the metal salt include alkali metal salts such as sodium salts, potassium salts, etc.; alkaline earth metal salts such as calcium salts, magnesium salts, barium salts, etc.; aluminum salts; and the like. Preferred examples of the salts with organic bases include salts with trimethylamine, triethylamine, pyridine, picoline, 2,6-lutidine, ethanolamine, diethanolamine, triethanolamine, cyclohexylamine, dicyclohexylamine, N,N'-dibenzylethylenediamine, etc. Preferred examples of the salts with inorganic acids include salts with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, etc. Preferred examples of salts with organic acids include salts with formic acid, acetic acid, trifluoroacetic acid, phthalic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, etc. Preferred examples of salts with basic amino acids include salts with arginine, lysine, ornithine, etc., and preferred examples of salts with acidic amino acids include salts with aspartic acid, glutamic acid, etc.

Among them, pharmaceutically acceptable salts are preferable. For example, when the compound has an acidic functional group, inorganic salts such as alkali metal salts (e.g., sodium salt, potassium salt, etc.), alkaline earth metal salts (e.g., calcium salt, magnesium salt, barium salt, etc.), ammonium salts, etc. are

preferable. When the compound has a basic functional group, salts with inorganic acids with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, etc., and salts with organic acids such as acetic acid, phthalic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, methanesulfonic acid, p-toluenesulfonic acid, etc. are preferable.

The prodrug of the metastin derivative or a salt thereof (hereinafter sometimes simply referred to as the metastin derivative of the present invention) refers to such a metastin derivative that is converted into the metastin derivative of the present invention under physiological conditions or with a reaction due to an enzyme, a gastric acid, etc., in the living body. In other words, the prodrug of the present invention is the metastin derivative that undergoes enzymatic oxidation, reduction, hydrolysis, etc. to be converted into the metastin derivative of the present invention, or the metastin derivative that undergoes hydrolysis, etc. by gastric acid, etc. to be converted into the metastin derivative of the present invention.

Examples of the prodrug of the metastin derivative of the present invention include a metastin derivative wherein an amino group of the metastin derivative of the present invention is substituted with an acyl, an alkyl, phosphoric acid, etc. (e.g., metastin derivatives wherein an amino group of the metastin derivative of the present invention is substituted with eicosanoyl, alanyl, pentylaminocarbonyl (5-methyl-2-oxo-1,3-dioxolen-4-yl)methoxycarbonyl, tetrahydrofuranyl, pyrrolidylmethyl, pivaloyloxymethyl, tert-butyl, etc); metastin derivatives wherein a hydroxy group of the metastin derivative of the present invention is substituted with an acyl, an alkyl, phosphoric acid, boric acid, etc. (e.g., metastin derivatives wherein an hydroxy group of the metastin derivative of the present invention is substituted with acetyl, palmitoyl, propanoyl, pivaloyl, succinyl, fumaryl, alanyl, dimethylaminomethylcarbonyl, etc.); and metastin derivatives wherein a carboxy group of the metastin derivative of the present invention is substituted with ester, amide, etc. (e.g., metastin derivatives wherein the carboxy group of the metastin derivative of the present invention is converted into the ethyl ester, phenyl ester, carboxymethyl ester, dimethylaminomethyl ester, pivaloyloxymethyl ester, ethoxycarbonyloxyethyl ester, phthalidyl ester, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl ester, cyclohexyloxycarbonylethyl ester, methylamide, etc); and the like. These metastin derivatives can be produced from the metastin derivatives of the present invention by per se known methods.

The prodrugs of the metastins derivative of the present invention may be

those that are converted into the metastin derivatives of the present invention under the physiological conditions as described in "Pharmaceutical Research and Development", Vol. 7 (Drug Design), pages 163-198, published 1990 by Hirokawa Publishing Co.

5 The metastin derivatives of the present invention or their salts or prodrugs (hereinafter sometimes simply referred to as the compound of the present invention) possess the cancer metastasis suppressing activity or the cancer growth suppressing activity. Thus, the metastin derivatives are useful for medicaments such as agents for preventing or treating all cancers (e.g., lung cancer, gastric cancer, liver cancer,
10 pancreas cancer, colorectal cancer, rectal cancer, colonic cancer, prostate cancer, ovarian cancer, cervical cancer, breast cancer, etc.), and the like.

 The compound of the present invention also possesses the effect of controlling pancreatic function and is thus useful as a medicament such as an agent for preventing or treating various pancreatic diseases (e.g., acute or chronic
15 pancreatitis, pancreatic cancer, etc.).

 Furthermore, the compound of the present invention possesses the effect of controlling placental function and is thus useful as a medicament such as an agent for preventing or treating choriocarcinoma, hydatid mole, invasive mole, miscarriage, fetal hypoplasia, abnormal glucose metabolism, abnormal lipid metabolism or
20 induction of delivery, etc.

 Moreover, the compound of the present invention possesses the effects of increasing sugar level, promoting pancreatic glucagon secretion and promoting urine formation and is thus useful as a medicament such as hyperglycemic agents, pancreatic glucagon secretagogue agents or agents for promoting urine formation,
25 which are useful for agents for preventing or treating obesity, hyperlipemia, type II diabetes mellitus, hypoglycemia, hypertension, diabetic neuropathy, diabetic nephropathy, diabetic retinopathy, edema, urinary disturbances, insulin resistance, unstable diabetes mellitus, fatty atrophy, insulin allergy, insulinoma, arteriosclerosis, thrombotic disorders or lipotoxicity; and the like.

30 In addition, the compound of the present invention also possesses the effects of promoting gonadotropic hormone (e.g., FSH, LH, etc.) secretion, promoting sex hormone [e.g., androgens (e.g., testosterone, androstenedione, etc.), estrogens (e.g., estradiol, estrone, etc.), progesterones, etc.] secretion, improving gonadal function and inducing or stimulating ovulation, as well as a sexual maturation effect, etc., and
35 hence, can be used as an agent for improving gonadal function, an agent for inducing

or stimulating ovulation, a gonadotropic hormone secretagogue agent or a sex hormone secretagogue agent, or an agent for preventing or treating hormone-dependent cancers [e.g., prostate cancer, breast cancer, etc.], infertility [e.g., irregular menstruation, dysmenorrhea, amenorrhea, weight loss-induced amenorrhea, secondary amenorrhea, anovulation, hypoovarianism, hypogonadism, spermatogenetic failure, hypogonadism (e.g., impotence, etc.), genital atrophy, testicular atrophy, testicular function disorder, azoospermia, hypoandrogenemia, etc.], endometriosis, early puberty, myoma of the uterus, etc.

Furthermore, the prodrug of the metastin derivative of the present invention or its salt is useful as an agent for preventing or treating Alzheimer's disease, Autism, moderate cognitive impairment, etc.

Moreover, the compound of the present invention has excellent blood stability, solubility and solution stability, as compared to native metastin such as metastin 54 (1-54) or metastin 10 (45-54).

The metastin derivative of the present invention or its salt or prodrug, metastin per se, or DNA encoding metastin, etc. is useful as an agent for suppressing gonadotropic hormone (e.g., FSH, LH) secretion or sex hormone [e.g., androgen (e.g., testosterone, androstenedione), estrogen (e.g., estradiol, estrone), progesterone] secretion; a down-regulating agent for gonadotropic hormone or sex hormone; in particular, it is useful for suppressing gonadotropic hormone secretion or sex hormone secretion via down-regulation of gonadotropic hormone or sex hormone (wherein, the down-regulation of gonadotropic hormone or sex hormone may be pulse loss of LHRH or depletion of LHRH) or down-regulation of human OT7T175 (metastin receptor) protein consisting of the amino acid sequence represented by SEQ ID NO: 9; particularly useful as an agent for preventing or treating hormone-dependent cancers (e.g., prostate cancer, breast cancer, etc.; especially a prostate cancer, hormone-sensitive prostate cancer, etc.); an agent for preventing or treating endometriosis; an agent for inhibiting ovarian follicular maturation; a menstrual cycle-suspending agent; an agent for treating myoma of the uterus; an agent for treating early puberty; or as a contraceptive, etc. Where the metastin derivative of the present invention or its salt or prodrug, metastin per se, or DNA encoding metastin, etc. has normal agonist activity, an effective dose of the metastin derivative sufficient to suppress the secretion of gonadotropic hormone or sex hormone is administered at the site or tissue where the therapeutic effects are to be exerted, so that the metastin derivative is present in a dose more than required (i.e.,

the metastin derivative is administered in an excess over the normal effective dose, at which the metastin derivative exerts the effects of suppressing cancer metastasis, suppressing cancer growth, etc.; or the effect of promoting gonadotropic hormone secretion, the effect of promoting sex hormone secretion, etc.) to exhibit the effects of suppressing gonadotropic hormone secretion or sex hormone secretion. Specific examples include sustained or continuous administration of the normal effective dose (including an administration technique to gradually release the pharmaceutical ingredients by bolus administration); and the like. Further when the metastin derivative of the present invention or its salt or the prodrug thereof, etc. has a sufficient agonist activity more than required (a super-agonist activity), it becomes possible to sustain the activities more than exhibited by the necessary dose at the site or tissue where the therapeutic effect are to be exhibited. It is therefore sufficient even by normal effective dose administration to suppress the secretion of gonadotropic hormone or sex hormone, whereby the effect of suppressing gonadotropic hormone secretion or sex hormone secretion is exhibited.

In other words, an effective dose of the metastin derivative of the present invention or its salt or prodrug, metastin per se, or DNA encoding metastin, etc. sufficient to suppress the secretion of gonadotropic hormone or sex hormone is administered so that the metastin derivative is present in a dose more than required at the site or tissue where the therapeutic effects are to be exerted, or its activities can be sustained more than required, which enables to exhibit the effects of suppressing gonadotropic hormone secretion or sex hormone secretion.

The pharmaceutical comprising the compound of the present invention is low toxic. Therefore, the compound of the present invention can be safely administered either directly as it is or as a mixture with pharmacologically acceptable carriers, orally or parenterally (e.g., topically, rectally, intravenously, etc.), in the form of pharmaceutical preparations such as tablets (including dragees and film-coated tablets), powdery dosage forms, granules, capsules (including soft capsules), liquid dosage forms, injections, suppositories, sustained release dosage forms, etc., in accordance with publicly known means generally used in process for producing pharmaceutical preparations.

The compound of the present invention is contained in the pharmaceutical preparation of the present invention in about 0.01 to about 100 wt%, based on the total weight of the preparation.

A dose of the compound of the present invention may vary depending upon

subject to be administered, target organ, conditions, route of administration, etc., and in oral administration, the compound is generally administered to the patient with cancer (as 60 kg body weight) in a daily dose of about 0.01 to about 100 mg, preferably about 0.1 to about 50 mg and more preferably about 0.1 to about 20 mg.

- 5 In parenteral administration, a single dose of the compound may vary depending upon subject to be administered, target organ, conditions, route of administration, etc., and in the form of an injectable dosage form, it is advantageous to administer the compound to the patient with cancer (as 60 kg body weight) generally in a daily dose of about 0.001 to about 30 mg, preferably about 0.01 to about 20 mg, and more
10 preferably about 0.01 to about 10 mg. For other animal species, the corresponding dose as converted per 60 kg weight can be administered.

- Pharmacologically acceptable carriers, which may be used in manufacturing the pharmaceutical preparation of the present invention, include various organic or inorganic carrier substances conventionally used as materials for pharmaceutical
15 preparations. These substances include, e.g., an excipient, a lubricant, a binder and a disintegrating agent in a solid dosage form, and a solvent, a dissolution aid, a suspending agent, an isotonicizing agent, a buffer, a soothing agent, etc. in a liquid dosage form. In addition, conventional additives such as a preservative, an antioxidant, a colorant, a sweetener, an adsorbent, a wetting agent, etc. can be
20 appropriately used in suitable amounts, if necessary.

Examples of excipients include lactose, saccharose, D-mannitol, starch, cornstarch, crystalline cellulose, light anhydrous silicic acid, etc.

Examples of lubricants include magnesium stearate, calcium stearate, talc, colloidal silica, etc.

- 25 Examples of binders include crystalline cellulose, saccharose, D-mannitol, dextrin, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, starch, sucrose, gelatin, methylcellulose, sodium carboxymethylcellulose, etc.

- Examples of disintegrating agents include starch, carboxymethylcellulose, carboxymethylcellulose calcium, sodium carboxymethyl starch,
30 L-hydroxypropylcellulose, etc.

Examples of solvents include water for injection, alcohol, propylene glycol, Macrogol, sesame oil, corn oil, olive oil, etc.

- Examples of dissolution aids include polyethylene glycol, propylene glycol,
35 D-mannitol, benzyl benzoate, ethanol, trisaminomethane, cholesterol,

triethanolamine, sodium carbonate, sodium citrate, etc.

Examples of suspending agents include surfactants such as stearyl triethanolamine, sodium laurylsulfate, lauryl aminopropionate, lecithin, benzalkonium chloride, benzethonium chloride, glycerine monostearate, etc.;

5 hydrophilic polymers such as polyvinyl alcohol, polyvinyl pyrrolidone, sodium carboxymethylcellulose, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, etc.

Examples of isotonizing agents include glucose, D-sorbitol, sodium chloride, glycerine, D-mannitol, etc.

10 Examples of buffers include buffering solutions of a phosphate, acetate, carbonate, citrate, etc.

Examples of soothing agents include benzyl alcohol, etc.

Examples of preservatives include p-hydroxybenzoates, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, sorbic acid, etc.

15 Examples of antioxidants include a sulfite, ascorbic acid, α -tocopherol, etc.

Furthermore, the compound of the present invention can be used in combination with drugs other than the compound of the present invention.

Examples of the drugs, which can be used in combination with the compound of the present invention (hereinafter sometimes simply referred to as
20 concomitant drugs), include chemotherapeutic agents for treating cancer, hormonal therapeutic agents, immunotherapeutic agents, etc. (hereinafter simply referred to as concomitant agents).

Examples of "chemotherapeutic agents" include alkylating agents, antimetabolites, anticancer antibiotics, anticancer agents derived from plants, etc.

25 Examples of "alkylating agents" include nitrogen mustard, nitrogen mustard-N-oxide hydrochloride, chlorambutyl, cyclophosphamide, ifosfamide, thiotepa, carboquone, improsulfan tosylate, busulfan, nimustine hydrochloride, mitobronitol, melphalan, dacarbazine, ranimustine, estramustine sodium phosphate, triethylenemelamine, carmustine, lomustine, streptozocin, pipobroman, etoglucid,
30 carboplatin, cisplatin, miboplatin, nedaplatin, oxaliplatin, altretamine, ambamustine, dibrospidium hydrochloride, fotemustine, prednimustine, pumitepa, ribomustin, temozolomide, treosulphan, trophosphamide, zinostatin stimalamer, carboquone, adozelesin, cystemustine, bizelesin, etc.

Examples of "antimetabolites" include mercaptopurine, 6-mercaptopurine
35 riboside, thioinosine, methotrexate, enocitabine, cytarabine, cytarabine ocfosfate,

ancitabine hydrochloride, 5-FU drugs (e.g., fluorouracil, tegafur, UFT, doxifluridine, carmofur, gallocitabine, emmitofur, etc.), aminopterin, leucovorin calcium, tabloid, butocine, folinate calcium, levofolinate calcium, cladribine, emitefur, fludarabine, gemcitabine, hydroxycarbamide, pentostatin, piritrexim, idoxuridine, mitoguazone, thiazophrine, ambamustine, etc.

Examples of "anticancer antibiotics" include actinomycin D, actinomycin C, mitomycin C, chromomycin A3, bleomycin hydrochloride, bleomycin sulfate, peplomycin sulfate, daunorubicin hydrochloride, doxorubicin hydrochloride, aclarubicin hydrochloride, pirarubicin hydrochloride, epirubicin hydrochloride, neocarzinostatin, mithramycin, sarcomycin, carzinophilin, mitotane, zorubicin hydrochloride, mitoxantrone hydrochloride, idarubicin hydrochloride, etc.

Examples of "anticancer agents derived from plants" include etoposide, etoposide phosphate, vinblastine sulfate, vincristine sulfate, vindesine sulfate, teniposide, paclitaxel, docetaxel, vinorelbine, etc.

Examples of "hormonal therapeutic agents" include fosfestrol, diethylstilbestrol, chlorotrianisene, medroxyprogesterone acetate, megestrol acetate, chlormadinone acetate, cyproterone acetate, danazol, allylestrenol, gestrinone, mepartricin, raloxifene, ormeloxifene, levormeloxifene, anti-estrogens (e.g., tamoxifen citrate, toremifene citrate, etc.), pill dosage forms, mepitiostane, testrolactone, aminoglutethimide, LH-RH agonists (e.g., goserelin acetate, buserelin, Leuprorelin, etc.), droloxifen, epitiostanol, ethinylestradiol sulfonate, aromatase inhibitors (e.g., fadrozole hydrochloride, anastrozole, retrozole, exemestane, vorozole, formestane, etc.), anti-androgens (e.g., flutamide, bicartamide, nilutamide, etc.), 5 α -reductase inhibitors (e.g., finasteride, epristeride, etc.), adrenocorticothormone drugs (e.g., dexamethasone, prednisolone, betamethasone, triamcinolone, etc.), androgen synthesis inhibitors (e.g., abiraterone, etc.), retinoid and drugs that retard retinoid metabolism (e.g., liarozole, etc.), and among them, LH-RH agonists (e.g., goserelin acetate, buserelin, Leuprorelin, etc.) are preferable.

Examples of "immunotherapeutic agents (BRM)" include picibanil, krestin, sizofiran, lentinan, ubenimex, interferons, interleukins, macrophage colony-stimulating factor, granulocyte colony-stimulating factor, erythropoietin, lymphotoxin, BCG vaccine, Corynebacterium parvum, levamisole, polysaccharide K, procodazole, etc.

The combined use of the compound of the present invention and a concomitant drug exhibits the following excellent effects.

(1) The dose can be reduced as compared to the dose when the compound of the present invention or a concomitant drug is administered alone.

(2) A drug concomitantly administered with the compound of the present invention can be chosen depending on the condition (mild, severe, etc.) of a patient.

5 (3) A concomitant drug, whose functional mechanism is different from that of the compound of the present invention, can be chosen so that a treatment period can be set longer.

(4) A concomitant drug, whose functional mechanism is different from that of the compound of the present invention, can be chosen so that sustained therapeutic
10 effects can be achieved.

(5) A synergistic effect can be obtained by the concomitant use of the compound of the present invention and a concomitant drug.

In addition, the compound of the present invention can reduce values of testosterone to emascuate level immediately after medication. Thus when the
15 concomitant drug such as LH-RH agonist (e.g., goserelin acetate, buserelin, Leuprorelin etc.; preferably Leuprorelin) is used in combination with the compound of the present invention, the values of testosterone can be reduced to emascuate level immediately after medication of the compound of the present invention. Further, since the combined use of the concomitant drug such as LH-RH agonist (e.g.,
20 goserelin acetate, buserelin, Leuprorelin, etc.; preferably Leuprorelin) and the compound of the present invention results in prolonged preservation of hormone-dependent period, it can advantageously be used.

Hereinafter, the combined use of the compound of the present invention and the concomitant drug is referred to as "the combined preparation of the present
25 invention."

When the combined preparation of the present invention is used, a dosing period of the compound of the present invention and the concomitant drug is not restricted; the compound of the present invention or its pharmaceutical composition and the concomitant drug or its pharmaceutical composition may be administered to
30 the subject to be administered either simultaneously or at certain time intervals. The dose of the concomitant drug may be modified according to the dose used clinically and may be appropriately chosen depending upon subject to be administered, route for administration, disease, combination, etc.

A mode for administration of the combined preparation of the present
35 invention is not particularly limited, but it is sufficient that the compound of the

present invention is used in combination with the concomitant drug at the time of administration. For such mode of administration, there are, for example, (1) administration of a single dosage form obtained by mixing the compound of the present invention and the concomitant drug together at the same time, (2) simultaneous administration of two dosage forms prepared separately from the compound of the present invention and the concomitant drug through the same route for administration, (3) administration of two dosage forms prepared separately from the compound of the present invention and the concomitant drug at certain time intervals through the same route for administration, (4) simultaneous administration of two dosage forms prepared separately from the compound of the present invention and the concomitant drug through different routes for administration, (5) administration of two dosage forms prepared separately from the compound of the present invention and the concomitant drug at certain time intervals (e.g., administration of the compound of the present invention and the concomitant drug in this order, or administration in a reversed order) through different routes for administration, etc.

The combined preparation of the present invention is low toxic and thus can be safely administered orally or parenterally (e.g., topically, rectally, intravenously, etc.) either directly as they are or in the form of pharmaceutical preparations such as tablets (including dragees and film-coated tablets), powdery dosage forms, granules, capsules (including soft capsules), liquid dosage forms, injections, suppositories, sustained release dosage forms, etc., which are obtained by mixing the compound of the present invention or (and) the concomitant drug described above with pharmacologically acceptable carriers by publicly known methods. Injectable dosage forms can be administered intravenously, intramuscularly or subcutaneously, into the organ, or directly at the focus.

Pharmacologically acceptable carriers, which may be used to manufacture the combined preparation of the present invention, include various organic or inorganic carrier substances conventionally used as materials for pharmaceutical preparations. These substances include, e.g., an excipient, a lubricant, a binder and a disintegrating agent in a solid dosage form, and a solvent, a dissolution aid, a suspending agent, an isotonicizing agent, a buffer, a soothing agent, etc. in a liquid dosage form. In addition, conventional additives such as a preservative, an antioxidant, a colorant, a sweetener, an adsorbent, a wetting agent, etc. can be appropriately used in suitable amounts, if necessary.

Examples of excipients include lactose, saccharose, D-mannitol, starch, cornstarch, crystalline cellulose, light anhydrous silicic acid, etc.

Examples of lubricants include magnesium stearate, calcium stearate, talc, colloidal silica, etc.

5 Examples of binders include crystalline cellulose, saccharose, D-mannitol, dextrin, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, starch, sucrose, gelatin, methylcellulose, sodium carboxymethylcellulose, etc.

10 Examples of disintegrating agents include starch, carboxymethylcellulose, carboxymethylcellulose calcium, sodium carboxymethyl starch, L-hydroxypropylcellulose, etc.

Examples of solvents include water for injection, alcohol, propylene glycol, Macrogol, sesame oil, corn oil, olive oil, etc.

15 Examples of dissolution aids include polyethylene glycol, propylene glycol, D-mannitol, benzyl benzoate, ethanol, trisaminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate, etc.

20 Examples of suspending agents include surfactants such as stearyltriethanolamine, sodium laurylsulfate, lauryl aminopropionate, lecithin, benzalkonium chloride, benzethonium chloride, glycerine monostearate, etc.; hydrophilic polymers such as polyvinyl alcohol, polyvinyl pyrrolidone, sodium carboxymethylcellulose, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, etc.

Examples of isotonizing agents include glucose, D-sorbitol, sodium chloride, glycerine, D-mannitol, etc.

25 Examples of buffers include buffering solutions of a phosphate, acetate, carbonate, citrate, etc.

Examples of soothing agents include benzyl alcohol, etc.

Examples of preservatives include p-hydroxybenzoates, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, sorbic acid, etc.

30 Examples of antioxidants include a sulfite, ascorbic acid, α -tocopherol, etc.

In the combined preparation of the present invention, a ratio of the compound of the present invention to the concomitant drug can be appropriately chosen depending upon subject to be administered, route for administration, disease, etc.

35 For example, the amount of the compound of the present invention

contained in the combined preparation of the present invention varies depending on the dosage form of the preparation, but is usually about 0.01 to 100% by weight, preferably about 0.1 to 50% by weight, and more preferably about 0.5 to 20% by weight, based on the total weight of the preparation.

5 The amount of the concomitant drug contained in the combined preparation of the present invention varies depending on the dosage form of the preparation, but is usually about 0.01 to 100% by weight, preferably about 0.1 to 50% by weight, and more preferably about 0.5 to 20% by weight, based on the total weight of the preparation.

10 The amount of additives such as a carrier, etc. contained in the combined preparation of the present invention varies depending on the dosage form of the preparation, and is usually about 1 to 99.99% by weight, preferably about 10 to 90% by weight, based on the total weight of the preparation.

 These amounts may be the same, also when the compound of the present
15 invention and the concomitant drug are separately prepared, respectively.

 These preparations can be manufactured by per se publicly known methods conventionally used in general.

 For example, the compound of the present invention or the concomitant
20 drug can be prepared into an injectable dosage form by formulating with a dispersing agent (e.g., Tween 80 (manufactured by Atlas Powder Company, USA), HCO 60 (manufactured by Nikko Chemicals Co., Ltd.), polyethylene glycol, carboxymethyl cellulose, sodium alginate, hydroxypropylmethyl cellulose, dextrin, etc.), a stabilizer (e.g., ascorbic acid, sodium pyrosulfite), a surfactant (e.g., polysorbate 80, macrogol, etc.), a solubilizing agent (e.g., glycerin, ethanol, etc.), a buffering agent (e.g.,
25 phosphoric acid or its alkali metal salt, citric acid or its alkali metal salt, etc.), an isotonizing agent (e.g., sodium chloride, potassium chloride, mannitol, sorbitol, glucose, etc.), a pH adjusting agent (e.g., hydrochloric acid, sodium hydroxide, etc.), a preservative (e.g., ethyl p-oxybenzoate, benzoic acid, methylparabene, propylparabene, benzyl alcohol, etc.), a solubilizer (e.g., concentrated glycerin,
30 meglumine, etc.), a dissolution aid (e.g., propylene glycol, saccharose, etc.), a soothing agent (e.g., glucose, benzyl alcohol, etc.), to prepare into aqueous injection; or by dissolving, suspending, or emulsifying with a vegetable oil such as olive oil, sesame oil, cottonseed oil, corn oil, etc., a dissolution aid such as propylene glycol or the like to prepare into an oily injection.

35 An oral dosage form can be produced in a conventional manner by adding to

the compound of the present invention or the concomitant drug, for example, an excipient (e.g., lactose, saccharose, starch, etc.), a disintegrating agent (e.g., starch, calcium carbonate, etc.), a binder (e.g., starch, gum arabic, carboxymethyl cellulose, polyvinylpyrrolidone, hydroxypropyl cellulose, etc.), a lubricant (e.g., talc, magnesium stearate, polyethylene glycol 6000, etc.) and other additives, compressing the resulting mixture and, if necessary, coating the compressed product for the purpose of taste masking, enteric degradation or sustained release by techniques per se publicly known. Coating agents for this purpose include, for example, hydroxypropylmethyl cellulose, ethyl cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, polyoxyethylene glycol, Tween 80, Prulonic F68, cellulose acetate phthalate, hydroxypropylmethyl cellulose phthalate, hydroxymethyl cellulose acetate succinate, Eudragit (manufactured by Rohm Company, Germany, methacrylic acid/acrylic acid copolymer) and dyes (e.g., iron oxide, titanium dioxide). The oral dosage form may be either an immediate release dosage form or a sustained release dosage form.

For example, in a suppository, the compound of the present invention or the concomitant drug is prepared into an oily or aqueous solid, semi-solid or liquid composition by techniques per se publicly known. Oily bases used for the composition described above include glycerides of higher fatty acids [e.g., cacao butter, uitepsols (manufactured by Dynamite Nobel Company, Germany), etc.], moderate fatty acids [e.g., miglyols (manufactured by Dynamite Nobel Company, Germany), etc.], vegetable oils (e.g., sesame oil, soybean oil, cottonseed oil, etc.), and the like. Aqueous bases include, for example, polyethylene glycols and propylene glycol. Bases for aqueous gels include, for example, natural rubbers, cellulose derivatives, vinyl polymers, acrylic polymers, etc.

Examples of the sustained release dosage form above include sustained release microcapsules, and the like.

Sustained release microcapsules can be obtained by per se publicly known methods, and are preferably prepared in the form of, e.g., a sustained release dosage form by the method [2] shown below and administered.

Preferably, the compound of the present invention is prepared into a dosage form for oral administration such as a solid dosage form (e.g., powdery dosage form, granules, tablets, capsules) or into a dosage form for rectal administration such as a suppository, etc. A dosage form for oral administration is particularly preferred.

The concomitant drug can be prepared into the dosage form described above,

depending on the kind of drug.

Hereinafter, [1] an injectable preparation of the compound of the present invention or the concomitant drug and its production, [2] a sustained release or immediate release preparation of the compound of the present invention or the concomitant drug and its production and [3] a sublingual, buccal or rapid oral disintegrating preparations of the compound of the present invention or the concomitant drug and its production will be specifically described.

[1] Injectable Preparation and its Production

10 An injectable preparation obtained by dissolving the compound of the present invention or the concomitant drug in water is preferred. The injectable preparation may contain a benzoate and/or a salicylate.

The injectable preparation is obtained by dissolving the compound of the present invention or the concomitant drug and optionally a benzoate and/or a salicylate in water.

Examples of the benzoate and/or salicylate described above include an alkali metal salt such as sodium and potassium salts, etc., an alkaline earth metal salt such as calcium and magnesium salts, etc., an ammonium salt, a meglumine salt, a salt of an organic acid such as trometamol, and the like.

20 The concentration of the compound of the present invention or the concomitant drug in the injectable preparation is about 0.05 to 50 w/v %, preferably about 0.3 to 20 w/v %. The concentration of the benzoate and/or salicylate is 0.5 to 50 w/v %, preferably 3 to 20 w/v %.

Furthermore, additives generally used in an injectable preparation such as a stabilizer (ascorbic acid, sodium pyrosulfite, etc.), a surfactant (polysorbate 80, macrogol, etc.), a solubilizing agent (glycerin, ethanol, etc.), a buffering agent (phosphoric acid and its alkali metal salt, citric acid and its alkali metal salt, etc.), an isotonicizing agent (sodium chloride, potassium chloride, etc.), a dispersing agent (hydroxypropylmethyl cellulose, dextrin), a pH adjusting agent (hydrochloric acid, sodium hydroxide, etc.), a preservative (ethyl p-oxybenzoate, benzoic acid, etc.), a solubilizer (concentrated glycerin, meglumine, etc.), a dissolution aid (propylene glycol, saccharose, etc.), a soothing agent (glucose, benzyl alcohol, etc.) are appropriately added to the preparation. Any of these additives is added in an amount generally used in an injectable preparation.

35 The injectable preparation is adjusted to pH of 2 to 12, preferably 2.5 to 8.0

by adding a pH adjusting agent.

The injectable preparation is obtained by dissolving both the compound of the present invention or the concomitant drug and optionally a benzoate and/or salicylate, and, if necessary, the above additives in water. These components may
5 be dissolved in any order according to the same manner as in a conventional injectable preparation.

An aqueous solution for injection is preferably warmed, and used as an injectable preparation after sterilization by filtration or autoclaved as in a conventional injectable preparation to provide for an injectable preparation.

10 An aqueous injectable preparation is preferably autoclaved, e.g., at 100 to 121°C for 5 to 30 minutes.

Moreover, the preparation may be in a solution form to which antibacterial activity is imparted to be usable as a multiple dosage form in divided dosing.

15 [2] Sustained Release or Immediate Release Preparation and its Production

A preferred sustained release preparation comprises a core comprising the compound of the present invention or the concomitant drug, which is optionally coated with a water-insoluble material or a swelling polymer. For example, a sustained release preparation for oral administration of a once-daily dosage form is
20 preferred.

Examples of the water-insoluble material used for the coating agent include cellulose ethers such as ethyl cellulose, butyl cellulose, etc., cellulose esters such as cellulose acetate, cellulose propionate, etc., polyvinyl esters such as polyvinyl acetate, polyvinyl butyrate, etc., acrylic acid polymers such as an acrylic acid/methacrylic acid copolymer, a methyl methacrylate copolymer, an ethoxyethyl
25 methacrylate/cinnamoethyl methacrylate/aminoalkyl methacrylate copolymer, a polyacrylic acid, a polymethacrylic acid, a methacrylic acid alkylamide copolymer, a poly(methyl methacrylate), a polymethacrylate, an aminoalkyl methacrylate copolymer, a poly(methacrylic anhydride), a glycidyl methacrylate copolymer, in particular, a series of Eudragits (Rohm & Pharma) such as Eudragit RS-100, RL-100, RS-30D, RL-30D, RL-PO and RS-PO (ethyl acrylate/methyl
30 methacrylate/chlorotrimethyl methacrylate/ethyl ammonium copolymer) and Eudragit NE-30D (methyl methacrylate/ethyl acrylate copolymer), etc., hydrogenated oils such as hydrogenated castor oil (e.g., LUBRI WAX (Freund Industrial Co., Ltd.), etc.), waxes such as carnauba wax, a fatty acid glycerin ester,
35

paraffin, etc., polyglycerin fatty acid esters, etc.

The swelling polymer is preferably a polymer having an acidic removable group and exhibiting pH-dependent swelling, and a polymer having an acidic removable group, which undergoes a less swelling at an acidic pH such as in the stomach but is swollen extensively at a neutral pH such as in the small and large intestines, is preferred.

Examples of such a polymer having an acidic removable group and exhibiting pH-dependent swelling include a crosslinked polyacrylic acid polymer such as Carbomers 934P, 940, 941, 974P, 980, 1342, etc., polycarbophil and calcium polycarbophil (all manufactured by BF Goodrich Chemicals), Hivis Wakos 103, 104, 105 and 304 (all manufactured by Wako Pure Chemical Industries, Ltd.), etc.

The coating agent used in the sustained release preparation may further contain a hydrophilic material.

Examples of the hydrophilic material include a polysaccharide which may have a sulfate group, such as pullulan, dextrin, alkali metal alginates, etc., a polysaccharide having a hydroxyalkyl group or a carboxyalkyl group such as hydroxypropyl cellulose, hydroxypropylmethyl cellulose, sodium carboxymethylcellulose, etc., methyl cellulose, polyvinyl pyrrolidone, polyvinyl alcohol, polyethylene glycol, etc.

The amount of the water-insoluble material contained in the coating agent of the sustained release preparation is about 30 to about 90% (w/w), preferably about 35 to about 80% (w/w), more preferably about 40 to about 75% (w/w), and the swelling polymer content is about 3 to about 30% (w/w), preferably about 3 to about 15% (w/w). The coating agent may further contain a hydrophilic material, and the amount of the hydrophilic material contained in the coating agent is about 50% (w/w) or less, preferably about 5 to about 40% (w/w), more preferably about 5 to about 35% (w/w). As used herein, the % (w/w) above is used to mean a % by weight based on the coating agent composition, which is the remainder of the coating agent solution after removing any solvent (e.g., water, a lower alcohol such as methanol, ethanol, etc.).

The sustained release preparation is manufactured by preparing a core containing a drug as illustrated below, followed by coating the resulting core with a coating agent solution obtained by heat-melting a water-insoluble material or a swelling polymer or by dissolving or dispersing such a material in a solvent.

I. Production of Drug-Containing Core

The shape of a core containing a drug to be coated with a coating agent (hereinafter sometimes simply referred to as a core) is not specifically limited but preferably prepared into a particulate shape such as granules, fine granules, or the like.

5 When the core is granules or fine granules, they have a mean particle size of preferably about 150 to about 2,000 μm , more preferably about 500 to about 1,400 μm .

 The core can be prepared in a conventional manner. For example, a drug is mixed with a suitable excipient, binder, disintegrating agent, lubricant, stabilizer, etc.,
10 and then subjected to wet extrusion granulation, fluidized bed granulation, or the like.

 The drug content in the core is about 0.5 to about 95% (w/w), preferably about 5.0 to about 80% (w/w), more preferably about 30 to about 70% (w/w).

 Examples of the excipient contained in the core include a saccharide such as
15 saccharose, lactose, mannitol, glucose, etc., starch, crystalline cellulose, calcium phosphate, cornstarch, etc. Among them, crystalline cellulose and cornstarch are preferred.

 Examples of the binder used include polyvinyl alcohol, hydroxypropyl cellulose, polyethylene glycol, polyvinyl pyrrolidone, Pluronic F68, gum arabic,
20 gelatin, starch, etc. Examples of the disintegrating agent include calcium carboxymethyl cellulose (ECG505), sodium croscarmellose (Ac-Di-Sol), crosslinked polyvinyl pyrrolidone (crospovidone), low substituted hydroxypropyl cellulose (L-HPC), etc. Among them, hydroxypropyl cellulose, polyvinyl pyrrolidone and low substituted hydroxypropyl cellulose are preferred. Examples of the lubricant
25 and the anticoagulant include talc, magnesium stearate and its inorganic salts, and examples of the lubricant include polyethylene glycol, etc. Examples of the stabilizer include an acid such as tartaric acid, citric acid, succinic acid, fumaric acid, maleic acid, etc.

 In addition to the technique described above, the core can be prepared by
30 using other techniques such as a tumbling granulation technique, a pan coating technique, a fluidized bed coating technique and a melt granulation technique, wherein a drug or a mixture of the drug with an excipient, a lubricant, etc. is portionwise added to inert carrier particles as seeds for the core with spraying a binder dissolved in a suitable solvent such as water, a lower alcohol (e.g., methanol,
35 ethanol, etc.) or the like. Examples of the inert carrier particles include those

prepared from saccharose, lactose, starch, crystalline cellulose and waxes, and, preferably, these carriers have a mean particle size of about 100 μm to about 1,500 μm .

In order to separate the drug contained in the core from a coating agent, the surface of the core may be covered with a protective material. Examples of the protective material include the hydrophilic material described above and water-insoluble material. The preferred protective material is polyethylene glycol or a polysaccharide having a hydroxyalkyl group or a carboxyalkyl group, more preferably, hydroxypropylmethyl cellulose and hydroxypropyl cellulose. The protective material may contain, as a stabilizer, an acid such as tartaric acid, citric acid, succinic acid, fumaric acid, maleic acid, etc., and a lubricant such as talc. When the protective material is used, the amount thereof to be coated is about 1 to about 15% (w/w), preferably about 1 to about 10% (w/w), more preferably about 2 to about 8% (w/w) based on the core.

The protective material can be coated by a conventional coating method and specifically, the core is spray-coated with the protective material by a fluidized bed coating technique, a pan coating technique, etc.

II. Coating of Core with Coating Agent

The core obtained in I above is coated with a coating agent solution prepared by melt-heating the water-insoluble material and pH-dependent swelling polymer described above and a hydrophilic material or by dissolving or dispersing them in a solvent to obtain a sustained release preparation.

As a coating method of the core with the coating agent solution, there are, for example, spray-coating, etc.

The composition ratio of the water-insoluble material, swelling polymer and hydrophilic material in the coating agent solution can be appropriately chosen to be within the amounts of the respective components contained in the coating.

The amount of the coating agent is about 1 to about 90% (w/w), preferably about 5 to about 50% (w/w), more preferably about 5 to about 35% (w/w) based on the core (excluding the protective material coating).

As the solvent for the coating agent solution, water and an organic solvent can be used singly or as a mixture thereof. When a mixture is used, the ratio of water and the organic solvent (water/organic solvent: a weight ratio) may vary with the range of 1 to 100%, and is preferably 1 to about 30%. The organic solvent is not particularly limited so far as it can dissolve the water-insoluble material, and

examples of the solvent include a lower alcohol such as methyl alcohol, ethyl alcohol, isopropyl alcohol, n-butyl alcohol, etc., a lower alkanone such as acetone, acetonitrile, chloroform, methylene chloride, etc. Among them, a lower alcohol is preferred, with ethyl alcohol and isopropyl alcohol being more preferred. Water
5 and a mixture of water and an organic solvent are used preferably as solvents for the coating agent solution. In this case, an acid such as tartaric acid, citric acid, succinic acid, fumaric acid, maleic acid, etc. may be added to the coating agent solution, if necessary, for the purpose of stabilizing the coating agent solution.

To carry out the coating through spray coating, the coating can be made
10 using a conventional coating method. Specifically, the core is sprayed with a coating agent solution by a fluidized bed coating technique, a pan coating technique, or the like. At this time, a lubricant such as talc, titanium oxide, magnesium stearate, calcium stearate, light silicic anhydride, etc., and a plasticizer such as glycerin fatty ester, hardened castor oil, triethyl citrate, cetyl alcohol, stearyl alcohol,
15 etc. may also be added.

After coating with a coating agent, an antistatic agent such as talc may also be admixed, if necessary.

The immediate release preparation may be a liquid (solution, suspension, emulsion, etc.) or a solid (particles, pills, tablets, etc.). An oral preparation and a
20 parenteral preparation such as an injectable preparation may be used, and an oral preparation is preferred.

The immediate release preparation may usually contain a carrier, additives and an excipient (hereinafter sometimes abbreviated as excipients) which are conventionally used in the pharmaceutical field, in addition to a drug which is an
25 active ingredient. The pharmaceutical excipients are not specifically limited so long as they are excipients conventionally used in the pharmaceutical field. Examples of the excipient for an oral solid preparation include lactose, starch, corn starch, crystalline cellulose (Avicel PH101, manufactured by Asahi Kasei Corporation, etc.), powdered sugar, granulated sugar, mannitol, light silicic
30 anhydride, magnesium carbonate, calcium carbonate, L-cysteine, etc., with corn starch and mannitol being preferred. Any of these excipients may be employed alone or in combination with each other. The amounts of the excipients are, for example, about 4.5 to about 99.4 w/w %, preferably about 20 to about 98.5 w/w %, more preferably about 30 to about 97 w/w %, based on the total weight of the immediate
35 release preparation.

The content of drug in the immediate release preparation may appropriately be selected from the range of about 0.5% to about 95%, preferably about 1% to about 60% to the whole amount of the immediate release preparation.

When the immediate release preparation is an oral solid preparation, the preparation contains a disintegrating agent in addition to the components described above. Examples of the disintegrating agent include calcium carboxymethylcellulose (ECG505 manufactured by GOTOKU CHEMICAL Co., Ltd.), sodium croscarmellose (for example, Ac-Di-Sol manufactured by Asahi Kasei Corporation), crospovidone (for example, COLIDON CL manufactured by BASF), low-substituted hydroxypropyl cellulose (Shin-Etsu chemical Co., Ltd.), carboxymethyl starch (MATSUTANI CHEMICAL INDUSTRY Co., Ltd.), sodium carboxymethyl starch (EXORITAB manufactured by KIMURA SANGYO), partial α starch (PCS manufactured by Asahi Kasei Corporation), etc. For example, the disintegrating agent that disintegrates granules by water absorption or swelling upon contact with water, or forming a channel between the active component comprising the core and an excipient can be used. Any of these disintegrating agents can be used alone or in combination with each other. The amount of the disintegrating agent used may be appropriately chosen depending upon the type and the amount of the drug used or a particular preparation design for the intended release performance. For example, the amount is about 0.05 to about 30 w/w %, preferably about 0.5 to about 15 w/w % based on the total weight of the immediate release preparation.

When the immediate release preparation is an oral solid preparation, the oral solid preparation may optionally contain additives conventionally used in a solid preparation, in addition to the components described above. Examples of the additives include binders (for example, sucrose, gelatin, powdery gum arabic, methyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, carboxymethylcellulose, polyvinyl pyrrolidone, pullran, dextrin, etc.), lubricants (polyethylene glycol, magnesium stearate, talc, light silicic anhydride (for example, aerosil (NIPPON AEROSIL)), surfactants (for example, anionic surfactants such as sodium alkylsulfate, nonionic surfactants such as polyoxyethylene fatty ester, polyoxyethylene sorbitan fatty ester, polyoxyethylene castor oil derivatives, etc.), colorants (for example, tar colorants, caramel, colcothar, titanium oxide, riboflavins), if necessary, corrigents (for example, sweeteners, flavors, etc.), adsorbents, preservatives, wetting agents, antistatic agents, etc. Furthermore, an organic acid such as tartaric acid, citric acid, succinic acid, fumaric acid or the like can also be

added as a stabilizer.

As the binder above, hydroxypropyl cellulose, polyethylene glycol and polyvinyl pyrrolidone, etc. are preferably used.

The immediate release preparation can be prepared by mixing the
5 components described above and kneading the mixture, if necessary, and then molding according to a conventional technique for making pharmaceutical preparations. The mixing above can be carried out in a conventional manner, e.g., by mixing, kneading, etc. Specifically, where the immediate release preparation is in the form of particles, the preparation can be prepared by mixing components with
10 a vertical granulator, a multi-purpose kneader (HATA IRON WORKS CO., LTD), a fluidized bed granulator FD-5S (POWREX CORPORATION) or the like, and then granulating the resulting by wet extrusion granulation or fluidized bed granulation by a technique similar to that for preparing the core of the sustained release preparation described above.

15 The immediate release preparation and the sustained release preparation thus obtained can be compounded, as they are, or, together with appropriate pharmaceutical excipients, in pharmaceutical preparations separately in a conventional manner to prepare respective preparations for administering in combination with each other simultaneously or at certain time intervals.
20 Alternatively, both preparations may be compounded in a single dosage form for oral administration (e.g., granules, fine granules, tablets, capsules) as they are, or, together with appropriate pharmaceutical excipients. Both preparations in the form of granules or fine granules may also be filled in a single capsule for oral administration.

25

[3] Sublingual, Buccal or Rapid Oral Disintegrating Preparation and its Production

A sublingual, buccal or rapid oral disintegrating preparation may be in the form of a solid preparation such as a tablet, or may be in the form of an oral mucosal patch (film) or oral disintegrating film.

30 The sublingual, buccal or rapid oral disintegrating preparation is preferably a preparation containing the compound of the present invention or the concomitant drug and an excipient. The preparation may also contain auxiliary agents such as a lubricant, an isotonicizing agent, a hydrophilic carrier, a water-dispersible polymer, a stabilizer, etc. Further for the purpose of promoting the absorption and enhancing
35 the bioavailability, the preparation may also contain β -cyclodextrin or β -cyclodextrin

derivatives (e.g., hydroxypropyl- β -cyclodextrin, etc.), and the like.

Examples of the above excipient include lactose, saccharose, D-mannitol, starch, crystalline cellulose, light silicic anhydride, etc. Examples of the lubricant include magnesium stearate, calcium stearate, talc, colloidal silica, etc., with
5 magnesium stearate and colloidal silica being preferred. Examples of the isotonizing agent include sodium chloride, glucose, fructose, mannitol, sorbitol, lactose, saccharose, glycerin and urea, with mannitol being particularly preferred. As the hydrophilic carrier, there are, for example, a swelling hydrophilic carrier such as crystalline cellulose, ethyl cellulose, crosslinked polyvinyl pyrrolidone, light
10 silicic anhydride, silicic acid, dicalcium phosphate, calcium carbonate, etc., with crystalline cellulose (e.g., microcrystalline cellulose, etc.) being preferred. As the water-dispersible polymer, there are, for example, a gum (e.g., tragacanth gum, acacia gum, guar gum), alginate (e.g., sodium alginate), cellulose derivatives (e.g., methyl cellulose, carboxymethylcellulose, hydroxymethyl cellulose, hydroxypropyl
15 cellulose, hydroxypropylmethyl cellulose), gelatin, water-soluble starch, polyacrylic acid (e.g., carbomer), polymethacrylic acid, polyvinyl alcohol, polyethylene glycol, polyvinyl pyrrolidone, polycarbophil, ascorbate palmitate salt, etc., with hydroxypropylmethyl cellulose, polyacrylic acid, alginate, gelatin, carboxymethylcellulose, polyvinyl pyrrolidone and polyethylene glycol being
20 preferred. Hydroxypropylmethyl cellulose is particularly preferred. As the stabilizer, there are, for example, cysteine, thiosorbitol, tartatic acid, citric acid, sodium carbonate, ascorbic acid, glycine, sodium sulfite, etc., with citric acid and ascorbic acid being particularly preferred.

The sublingual, buccal or rapid oral disintegrating preparation can be
25 prepared by mixing the compound of the present invention or the concomitant drug and an excipient by a method per se known. Furthermore, if desired, the auxiliary agents described above, such as the lubricant, isotonizing agent, hydrophilic carrier, water-dispersible polymer, stabilizer, colorant, sweetener, preservative, etc. may also be admixed. After mixing the components described above simultaneously or at
30 certain time intervals, the mixture is compressed into tablets to obtain the sublingual, buccal or rapid oral disintegration tablet. In order to obtain a suitable hardness, a solvent such as water, an alcohol, etc. can be used to moisturize or wet the components before or after tableting, followed by drying.

In preparing the oral mucosal patch (film), the compound of the present
35 invention or the concomitant drug and the water-dispersible polymer (preferably,

hydroxypropyl cellulose, hydroxypropylmethyl cellulose), excipient, etc. described above are dissolved in a solvent such as water, etc. and then the resulting solution is cast into a film. In addition, additives such as a plasticizer, a stabilizer, an antioxidant, a preservative, a colorant, a buffering agent, a sweeteners, etc. may be added to the preparation. A glycol such as polyethylene glycol, propylene glycol, etc. may be added to impart an appropriate elasticity to a film, and a bioadhesive polymer (e.g., polycarbophile, carbopol) may also be added to enhance the adhesion of the film to the oral mucosal lining. The casting can be carried out by pouring a solution onto a non-adhesive surface, spreading the solution using a coater such as a doctor blade in a uniform thickness (preferably, approximately 10 to 1000 microns), and then drying the solution to form a film. The film thus formed is dried at room temperature or while warming, and then cut into pieces each having a desired surface area.

A preferred rapid oral disintegrating preparation is, for example, a rapid diffusion preparation in a solid network form, which comprises the compound of the present invention or the concomitant drug and a water-soluble or water-diffusible carrier inert to the compound of the present invention or the concomitant drug. The network is formed by sublimating a solvent from the solid composition comprising a solution of the compound of the present invention or the concomitant drug in a suitable solvent.

In addition to the compound of the present invention or the concomitant drug, the composition of the rapid oral disintegrating preparation may preferably contain a matrix-forming agent and secondary components.

Examples of the matrix-forming agent include gelatins, dextrans and animal or vegetable proteins from soybean, wheat, psyllium seed, etc.; gummy materials such as gum arabic, guar gum, agar, xanthane gum, etc.; polysaccharides; alginates; carboxymethylcelluloses; carrageenans; dextrans; pectins; synthetic polymers such as polyvinyl pyrrolidones; materials derived from gelatin-gum arabic complexes, etc. The matrix-forming agent further includes saccharides such as mannitol, dextrose, lactose, galactose, trehalose, etc.; cyclic saccharides such as cyclodextrins, etc.; inorganic salts such as sodium phosphate, sodium chloride, aluminum silicate, etc.; amino acids having 2 to 12 carbon atoms such as glycine, L-alanine, L-aspartic acid, L-glutamic acid, L-hydroxyproline, L-isoleucine, L-leucine, L-phenylalanine, etc.

One or more matrix-forming agents can be incorporated into a solution or suspension before solidification. The matrix-forming agents may be present in

addition to a surfactant, or may be present in the absence of a surfactant. The matrix-forming agents serve not only to form a matrix itself, but also assist to maintain diffusion of the compound of the present invention or the concomitant drug in the solution or suspension.

5 The composition may contain a secondary component such as a preservative, an antioxidant, a surfactant, a thickening agent, a colorant, pH adjusting agent, a flavor, a sweetener, a taste masking agent, etc. As the suitable colorant, there are, for example, iron oxide red, black and yellow, FD & C dyes available from ERIS & EVERALD such as FD & C Blue No. 2 and FD & C Red No. 40, etc. Examples of
10 the suitable flavor include mint, raspberry, licorice, orange, lemon, grape fruit, caramel, vanilla, cherry, grape flavor and a combination thereof. Examples of the suitable pH adjusting agent include citric acid, tartaric acid, phosphoric acid, hydrochloric acid and maleic acid. Examples of the suitable sweetener include aspartame, acesulfame K and thaumatococin. Examples of the suitable taste masking
15 agent include sodium bicarbonate, ion exchange resins, cyclodextrin inclusion compounds, adsorbents and microencapsulated apomorphine.

 The preparation generally contains the compound of the present invention or the concomitant drug in an amount of about 0.1 to about 50% by weight, preferably about 0.1 to about 30% by weight and, preferably, the preparation (the sublingual
20 tablet, buccal, etc. described above) allows 90% or more of the compound of the present invention or the concomitant drug to be dissolved (in water) within a time period of about 1 to about 60 minutes, preferably about 1 minute to about 15 minutes, more preferably about 2 minutes to about 5 minutes, or is a rapid oral disintegrating preparation which disintegrates within about 1 to about 60 seconds, preferably about
25 1 to about 30 seconds, more preferably about 1 to about 10 seconds, after being placed in the oral cavity.

 The amount of the above excipient is about 10 to about 99% by weight, preferably about 30 to about 90% by weight based on the total weight of the preparation. The amount of β -cyclodextrin or β -cyclodextrin derivative is about 0
30 to about 30% by weight based on the total weight of the preparation. The amount of the lubricant is about 0.01 to about 10% by weight, preferably about 1 to about 5% by weight based on the total weight of the preparation. The amount of the isotonicizing agent is about 0.1 to about 90% by weight, preferably about 10 to about 70% by weight based on the total weight of the preparation. The amount of the
35 hydrophilic carrier is about 0.1 to about 50% by weight, preferably about 10 to about

30% by weight based on the total weight of the preparation. The amount of the water-dispersible polymer is about 0.1 to about 30% by weight, preferably about 10 to about 25% by weight based on the total weight of the preparation. The amount of the stabilizer is about 0.1 to about 10% by weight, preferably about 1 to about 5% by weight based on the total weight of the preparation. If necessary, the preparation described above may further contain additives such as a colorant, a sweetener, a preservative, etc.

A dose of the combined preparations of the present invention varies depending upon kind of the compound of the present invention, age, body weight, conditions, dosage form, route for administration, dosing period, etc.

A dose of the compound of the present invention may vary depending upon subject to be administered, target organ, conditions, route of administration, etc., and in oral administration, the compound is generally administered to the patient with cancer (as 60 kg body weight) in a daily dose of about 0.01 to about 100 mg, preferably about 0.1 to about 50 mg and more preferably about 0.1 to about 20 mg. In parenteral administration, a single dose of the compound may vary depending upon subject to be administered, target organ, conditions, route of administration, etc., and in the form of an injectable dosage form, it is advantageous to intravenously administer the compound to the patient with cancer (as 60 kg body weight) generally in a daily dose of about 0.001 to about 30 mg, preferably about 0.01 to about 20 mg, and more preferably about 0.01 to about 10 mg. For other animal species, the corresponding dose as converted per 60 kg weight can be administered. Of course, the dose may vary depending on individual conditions as described above; in such a case, a dose less than the dose given above may be sufficient, or a dose higher than the range above may be used.

It is possible to set any range of a dose for the concomitant drug, so long as it causes no adverse side effects. A daily dose of the concomitant drug may vary depending on the severity of disease, the age, sex, body weight and susceptibility of the subject, dosing period and intervals, characteristics, formulation, type and active components of the pharmaceutical preparation, etc. and is not particularly limited. For example, in oral administration, the dose is about 0.001 to 2000 mg, preferably about 0.01 to 500 mg, and more preferably about 0.1 to 100 mg per kg body weight of mammals in terms of a drug; usually, this dose is administered by dividing 1 to 4 times per day.

When the pharmaceutical preparations of the present invention are

administered, they may be administered concomitantly. Alternatively, the concomitant drug is first administered and then the compound of the present invention is administered, or the compound of the present invention is first administered and then the concomitant drug is administered. When they are
5 administered at certain time intervals, the intervals vary depending on the active component to be administered, dosage form and route of administration; for example, when the concomitant drug is first administered, the compound of the present invention may be administered within 1 minute to 3 days, preferably 10 minutes to 1 day, more preferably 15 minutes to 1 hour after the administration of the concomitant
10 drug. When the compound of the present invention is first administered, the concomitant drug may be administered within 1 minute to 1 day, preferably 10 minutes to 6 hours, more preferably 15 minutes to 1 hour after the administration of the compound of the present invention.

As a preferred method of administration, for example, about 0.001 to 200
15 mg/kg of the concomitant drug in the form of an oral dosage preparation is administered orally and, after about 15 minutes, about 0.005 to 0.5 mg/kg of the compound of the present invention in the form of a parenteral preparation is administered parenterally as a daily dose.

As the metastins, there are used, for example, human metastin described in
20 WO 00/24890, mouse or rat metastin described in WO 01/75104, etc.

Specific examples of human metastin include a peptide comprising the N-terminal 47th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 and consisting of 8 to 54 amino acid residues, and the like.

The "peptide comprising the N-terminal 47th-54th amino acid sequence in
25 the amino acid sequence represented by SEQ ID NO: 1 and consisting of 8 to 54 amino acid residues" may be any peptide, as far as it is a peptide comprising the N-terminal 47th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 and consisting of 8 to 54 amino acid residues, but means that these peptides have substantially the same physiological activity (e.g., a receptor binding
30 activity, a signal transduction action, a sugar level elevating action, a pancreatic glucagon secretion promoting action, a urine formation promoting action, etc.). Specifically, there are used (i) a peptide having the amino acid sequence represented by SEQ ID NO: 1, (ii) a peptide comprising the N-terminal 47th-54th amino acid sequence at the C terminus in the amino acid sequence represented by SEQ ID NO: 1
35 and consisting of 8 to 15 amino acid residues, etc.

More specifically, human metastin used includes (i) a peptide consisting of the amino acid sequence represented by SEQ ID NO: 1 (human metastin 54 (1-54)), (ii) a peptide consisting of the N-terminal 40th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 (human metastin 15 (40-54); SEQ ID NO: 15), (iii) a peptide consisting of the N-terminal 45th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 (human metastin 10 (45-54); SEQ ID NO: 16), (iv) a peptide consisting of the N-terminal 46th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 (human metastin 9 (46-54); SEQ ID NO: 17), (v) a peptide consisting of the N-terminal 47th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 (human metastin 8 (47-54); SEQ ID NO: 18), etc.

As mouse metastin (A), there are used, for example, (i) a peptide comprising the N-terminal 134th-141st amino acid sequence in the amino acid sequence represented by SEQ ID NO: 3 and consisting of 8 to 52 amino acid residues. Specific examples of mouse metastin (A) used include (i) a peptide consisting of the N-terminal 90th-141st amino acid sequence in the amino acid sequence represented by SEQ ID NO: 3, (ii) a peptide consisting of the N-terminal 132nd-141st amino acid sequence in the amino acid sequence represented by SEQ ID NO: 3, (iii) a peptide consisting of the N-terminal 127th-141st amino acid sequence in the amino acid sequence represented by SEQ ID NO: 3, and the like.

As mouse metastin (B), there are used, for example, (i) a peptide comprising the N-terminal 138th-145th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 5 and consisting of 8 to 52 amino acid residues. Specific examples of mouse metastin (B) used include a peptide consisting of the N-terminal 94th-145th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 5, and the like.

As rat metastin, there are used, for example, (i) a peptide comprising the N-terminal 112th-119th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 7 and consisting of 8 to 52 amino acid residues. Specific examples of rat metastin used include (i) a peptide consisting of the N-terminal 68th-119th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 7, (ii) a peptide consisting of the N-terminal 110th-119th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 7, (iii) a peptide consisting of the N-terminal 105th-119th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 7, and the like.

Throughout the specification, the metastins are represented in accordance with the conventional way of describing peptides, that is, the N-terminus (amino terminus) at the left hand and the C-terminus (carboxyl terminus) at the right hand. In the peptide represented by SEQ ID NO: 1, the C-terminus may be in any form of a carboxyl group (-COOH), a carboxylate (-COO⁻), an amide (-CONH₂) and an ester (-COOR). Herein, examples of R of the ester group or alkyl amide include a C₁₋₆ alkyl group such as methyl, ethyl, n-propyl, isopropyl, n-butyl, etc.; a C₃₋₈ cycloalkyl group such as cyclopentyl, cyclohexyl, etc.; a C₆₋₁₂ aryl group such as phenyl, α-naphthyl, etc.; a C₇₋₁₄ aralkyl such as a phenyl-C₁₋₂ alkyl group, e.g., benzyl, phenethyl, benzhydryl, etc., or an α-naphthyl-C₁₋₂ alkyl group such as α-naphthylmethyl, etc.; pivaloyloxymethyl group, which are widely used as an ester for oral use, and the like.

Furthermore, the metastins include peptides, wherein the amino group at the N-terminal methionine residue is protected with a protecting group (e.g., a C₁₋₆ acyl group such as a C₂₋₆ alkanoyl group, e.g., formyl group, acetyl group, etc.); those wherein the N-terminal region is cleaved in vivo and the glutamyl group thus formed is pyroglutaminated; those wherein a substituent (e.g., -OH, -SH, amino group, imidazole group, indole group, guanidino group, etc.) on the side chain of an amino acid in the molecule is protected with a suitable protecting group (e.g., a C₁₋₆ acyl group such as a C₂₋₆ alkanoyl group, e.g., formyl group, acetyl group, etc.), or conjugated peptides such as glycopeptides bound to sugar chains.

For salts of the metastin of the present invention, preferred are salts with physiologically acceptable bases (e.g., alkali metal salts) or acids (e.g., organic acids or inorganic acids), etc., especially preferred are physiologically acceptable acid addition salts. Examples of such salts include salts with, for example, inorganic acids (e.g., hydrochloric acid, phosphoric acid, hydrobromic acid, sulfuric acid); salts with organic acids (e.g., acetic acid, formic acid, propionic acid, fumaric acid, maleic acid, succinic acid, tartaric acid, citric acid, malic acid, oxalic acid, benzoic acid, methanesulfonic acid, benzenesulfonic acid) and the like.

As the DNAs encoding metastins, there are used, for example, DNAs encoding human metastin described in WO 00/24890, DNAs encoding mouse or rat metastin described in WO 01/75104, etc.

The DNAs encoding the metastins may be any of genomic DNA, genomic DNA library, cDNA derived from the cells and tissues described above, cDNA library derived from the cells and tissues described above and synthetic DNA. The

vector to be used for the library may be any of bacteriophage, plasmid, cosmid and phagemid. The DNA may also be directly amplified by reverse transcriptase polymerase chain reaction (hereinafter abbreviated as RT-PCR) using the total RNA or mRNA fraction prepared from the cells and tissues described above.

5 The DNA encoding human metastin, mouse metastin precursor (A), mouse metastin precursor (B) or rat metastin precursor may be any DNA, so long as each is a DNA containing a base sequence represented by SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8, or a DNA having a base sequence hybridizable to the base sequence represented by any base sequence represented by SEQ ID NO: 2,
10 SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8 under highly stringent conditions and encoding the human metastin, mouse metastin (A), mouse metastin (B) or rat metastin described above.

 Specific examples of the DNA hybridizable to the base sequence represented by any of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO:
15 8 include DNAs containing a base sequence having at least about 70% homology, preferably at least about 80% homology, more preferably at least about 90% homology and the most preferably at least about 95% homology, to the base sequence represented by any of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8.

20 Homology in the base sequence can be measured under the following conditions (an expectation value = 10; gaps are allowed; filtering = ON; match score = 1; mismatch score = -3) using the homology scoring algorithm NCBI BLAST (National Center for Biotechnology Information Basic Local Alignment Search Tool).

25 The hybridization can be carried out by per se publicly known methods or by modifications of these methods, for example, according to the method described in Molecular Cloning, 2nd (J. Sambrook et al., Cold Spring Harbor Lab. Press, 1989). A commercially available library may also be used according to the instructions of the attached manufacturer's protocol. Preferably, the hybridization can be carried
30 out under highly stringent conditions.

 The highly stringent conditions used herein are, for example, those in a sodium concentration at about 19 to 40 mM, preferably about 19 to 20 mM at a temperature of about 50 to 70°C, preferably about 60 to 65°C. In particular, hybridization conditions in a sodium concentration of about 19 mM at a temperature
35 of about 65°C are most preferred.

Specifically, as the DNA encoding the human metastin consisting of the amino acid sequence represented by SEQ ID NO: 1, the DNA consisting of the base sequence represented by SEQ ID NO: 2 is used. Accordingly, for the base sequence encoding the human metastin consisting of the various amino acid sequences described above, a base sequence corresponding to each of the partial amino acid sequences in the amino acid sequence represented by SEQ ID NO: 1 may be chosen from the base sequence represented by SEQ ID NO: 2.

As the DNA encoding the mouse metastin precursor (A) comprising the amino acid sequence represented by SEQ ID NO: 3, there are employed a DNA consisting of the base sequence represented by SEQ ID NO: 4, and the like. Accordingly, for the base sequence encoding the mouse metastin precursor (A) consisting of the various amino acid sequences described above, a base sequence corresponding to each of the partial amino acid sequences in the amino acid sequence represented by SEQ ID NO: 3 may be chosen from the base sequence represented by SEQ ID NO: 4.

As the DNA encoding the mouse metastin precursor (B) comprising the amino acid sequence represented by SEQ ID NO: 5, there are employed a DNA consisting of the base sequence represented by SEQ ID NO: 6, and the like. Accordingly, for the base sequence encoding the mouse metastin precursor (B) comprising of the various amino acid sequences described above, a base sequence corresponding to each of the partial amino acid sequences in the amino acid sequence represented by SEQ ID NO: 5 may be chosen from the base sequence represented by SEQ ID NO: 6.

As the DNA encoding the rat metastin comprising the amino acid sequence represented by SEQ ID NO: 7, there are employed a DNA consisting of the base sequence represented by SEQ ID NO: 8, and the like. Accordingly, for the base sequence encoding the rat metastin consisting of the various amino acid sequences described above, a base sequence corresponding to each of the partial amino acid sequences in the amino acid sequence represented by SEQ ID NO: 7 may be chosen from the base sequence represented by SEQ ID NO: 8.

More specifically, for the peptide consisting of the amino acid sequence represented by SEQ ID NO: 1 (human metastin 54 (1-54)), a DNA containing the base sequence represented by SEQ ID NO: 2, etc. is used.

For the peptide consisting of the N-terminal 40th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 (human metastin 15

(40-54); SEQ ID NO: 15); a DNA containing the base sequence represented by SEQ ID NO: 19, etc. is used.

For the peptide consisting of the N-terminal 45th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 (human metastin 10 (45-54); represented by SEQ ID NO: 16), a DNA containing the base sequence represented by SEQ ID NO: 20, etc. is used.

For the peptide consisting of the N-terminal 46th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 (human metastin 9 (46-54); represented by SEQ ID NO: 17), a DNA containing the base sequence represented by SEQ ID NO: 21, etc. is used.

For the peptide consisting of the N-terminal 47th-54th amino acid sequence in the amino acid sequence represented by SEQ ID NO: 1 (human metastin 8 (47-54); represented by SEQ ID NO: 18), a DNA containing the base sequence represented by SEQ ID NO: 22, etc. is used.

As the metastin receptor, its partial peptides or salts thereof, there are used, for example, a human metastin receptor, its partial peptides or salts thereof described in WO 00/24890, a mouse or rat human metastin receptor, its partial peptides or salts thereof described in WO 01/75104, etc.

Specifically, a protein comprising the same or substantially the same amino acid sequence as the amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13, etc. is used as the metastin receptor.

The amino acid sequence which is substantially the same amino acid sequence as the amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13 includes, for example, an amino acid sequence having at least about 70% homology, preferably at least about 80% homology, more preferably at least about 90% homology, and most preferably at least about 95% homology, to the amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13.

Homology of the amino acid sequences can be determined under the following conditions (an expectation value = 10; gaps are allowed; matrix = BLOSUM62; filtering = OFF) using a homology scoring algorithm NCBI BLAST (National Center for Biotechnology Information Basic Local Alignment Search Tool).

As the protein comprising substantially the same amino acid sequence as the amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO:

13, preferred is a protein having substantially the same amino acid sequence as the amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13 and having the activity of the same nature as that of a protein having the amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13,
5 etc.

As the activity of substantially the same nature, there are, for example, a ligand binding activity, a signal transduction activity, and the like. The "substantially the same nature" is used to mean that the nature of these activities is equivalent in terms of quality. Thus, the activities such as a ligand binding activity,
10 a signal transduction activity, etc. are preferably equivalent (e.g., about 0.01 to 100 times, preferably about 0.5 to 10 times, more preferably 0.5 to 2 times), but differences in quantitative factors such as a level of these activities, or such as a molecular weight of the protein may be present and allowable.

The activities such as a ligand binding activity, a signal transduction activity,
15 etc. can be assayed by per se publicly known method with modifications and may be determined according to methods of determining a ligand or screening methods described in, e.g., WO 00/24890 or WO 01/75104.

Examples of the metastin receptor used include proteins comprising (i) the amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO:
20 13, of which at least 1 or 2 (preferably about 1 to about 30, more preferably about 1 to about 10 and most preferably several (1 or 2)) amino acids are deleted; (ii) the amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13, to which at least 1 or 2 (preferably about 1 to about 30, more preferably about 1 to about 10 and most preferably several (1 or 2)) amino acids are added; (iii) the
25 amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13, in which at least 1 or 2 (preferably about 1 to about 30, more preferably about 1 to about 10 and most preferably several (1 or 2)) amino acids are substituted by other amino acids; or (iv) a combination of these amino acid sequences; and the like.

Throughout the specification, the metastin receptors are represented in
30 accordance with the conventional way of describing peptides, that is, the N-terminus (amino terminus) at the left hand and the C-terminus (carboxyl terminus) at the right hand. In the metastin receptors including the metastin receptor represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13, the C-terminus may be in any form of a carboxyl group (-COOH), a carboxylate (-COO-), an amide (-CONH₂) and an ester
35 (-COOR). Herein, examples of R of the ester group include a C₁₋₆ alkyl group such as

methyl, ethyl, n-propyl, isopropyl, n-butyl, etc.; a C₃₋₈ cycloalkyl group such as cyclopentyl, cyclohexyl, etc.; a C₆₋₁₂ aryl group such as phenyl, α -naphthyl, etc.; a C₇₋₁₄ aralkyl such as a phenyl-C₁₋₂ alkyl group, e.g., benzyl, phenethyl, etc., or an α -naphthyl-C₁₋₂ alkyl group such as α -naphthylmethyl, etc.; and pivaloyloxymethyl
5 group, which are widely used as an ester for oral use, and the like.

Where the metastin receptors contain a carboxyl group (or a carboxylate) at a position other than the C-terminus, the carboxyl group may be amidated or esterified and such amides or esters are also included within the receptor protein of the present invention. In this case, the ester group used may be the same group as
10 the C-terminal esters described above.

Furthermore, the metastin receptors include those wherein the amino group at the N-terminal methionine residue is protected with a protecting group (e.g., a C₁₋₆ acyl group such as a C₂₋₆ alkanoyl group, e.g., formyl group, acetyl group, etc.); those wherein the N-terminal region is cleaved in vivo and the glutamyl group thus
15 formed is pyroglutaminated; those wherein a substituent (e.g., -OH, -SH, amino group, imidazole group, indole group, guanidino group, etc.) on the side chain of an amino acid in the molecule is protected with a suitable protecting group (e.g., a C₁₋₆ acyl group such as a C₂₋₆ alkanoyl group, e.g., formyl group, acetyl group, etc.), or conjugated proteins such as glycoproteins bound to sugar chains.

Specific examples of the metastin receptors include human metastin receptor consisting of the amino acid sequence represented by SEQ ID NO: 9, rat metastin
20 receptor consisting of the amino acid sequence represented by SEQ ID NO: 11, mouse metastin receptor consisting of the amino acid sequence represented by SEQ ID NO: 13, etc.

The partial peptides of the metastin receptor (hereinafter sometimes simply referred to as the partial peptide) may be any peptide, so long as they are partial
25 peptides of the metastin receptor described above; there are used those such as protein molecules of the metastin receptor, which are the sites exposed outside the cell membrane, and having a ligand binding activity.

Specifically, the partial peptide of the metastin receptor consisting of the amino acid sequence represented by SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13 is a peptide containing the parts analyzed to be extracellular domains (hydrophilic
30 domains) in the hydrophobic plotting analysis. A peptide containing a hydrophobic domain in part can be used as well. In addition, the peptide may contain each
35 domain separately or a plurality of domains together.

In the metastin receptor, preferred partial peptides are those having the number of amino acids of at least 20, preferably at least 50, and more preferably at least 100, in the amino acid sequence described above, which constitutes the metastin receptor.

5 The partial peptide may be a peptide having the amino acid sequence described above, of which at least 1 or 2 (preferably about 1 to about 10 and more preferably several (1 or 2)) amino acids are deleted; to which at least 1 or 2 (preferably about 1 to about 10 and more preferably several (1 or 2)) amino acids are added; or, in which at least 1 or 2 (preferably about 1 to about 10 and more
10 preferably several (1 or 2)) amino acids are substituted by other amino acids.

In the partial peptide, the C terminus may be any form of a carboxyl group (-COOH), a carboxylate (-COO⁻), an amide (-CONH₂) and an ester (-COOR), as in the metastin receptor described above.

Furthermore, the partial peptides include peptides, wherein the amino group
15 at the N-terminal methionine residue is protected with a protecting group; those wherein the N-terminal region is cleaved in vivo and the glutamyl group thus formed is pyroglutaminated; those wherein a substituent on the side chain of an amino acid in the molecule is protected with a suitable protecting group, or conjugated peptides such as glycopeptides bound to sugar chains, as in the metastin receptors described
20 above.

For salts of the metastin receptor or the partial peptide, preferred are salts with physiologically acceptable acids, especially physiologically acceptable acid addition salts. Examples of the salts include salts with, for example, inorganic acids (e.g., hydrochloric acid, phosphoric acid, hydrobromic acid, sulfuric acid); salts with
25 organic acids (e.g., acetic acid, formic acid, propionic acid, fumaric acid, maleic acid, succinic acid, tartaric acid, citric acid, malic acid, oxalic acid, benzoic acid, methanesulfonic acid, benzenesulfonic acid) and the like.

As the DNA encoding the metastin receptor or its partial peptides, there are used, for example, a DNA encoding the human metastin receptor or its partial
30 peptides described in WO 00/24890, a DNA encoding the mouse or rat metastin receptor or its partial peptides described in WO 01/75104, etc.

The DNAs encoding the metastin receptor or its partial peptides may be any of genomic DNA, genomic DNA library, cDNA derived from the cells and tissues described above, cDNA library derived from the cells and tissues described above
35 and synthetic DNA. The vector to be used for the library may be any of

bacteriophage, plasmid, cosmid and phagemid. The DNA may also be directly amplified by reverse transcriptase polymerase chain reaction (hereinafter abbreviated as RT-PCR) using the total RNA or mRNA fraction prepared from the cells and tissues described above.

5 Specifically, the DNA encoding human metastin receptor, mouse metastin receptor or rat metastin receptor may be any DNA, so long as it is a DNA comprising each base sequence represented by SEQ ID NO: 10, SEQ ID NO: 12 or SEQ ID NO: 14, or a DNA comprising a base sequence hybridizable to the base sequence represented by SEQ ID NO: 10, SEQ ID NO: 12 or SEQ ID NO: 14 under highly
10 stringent conditions and encoding a receptor having the activity of substantially the same nature (e.g., a ligand binding activity, a signal transduction activity, etc.) as that of the human metastin receptor, mouse metastin receptor or rat metastin receptor consisting of the amino acid sequence represented by SEQ ID NO: 10, SEQ ID NO: 12 or SEQ ID NO: 14.

15 Examples of the DNA hybridizable to the base sequence represented by any of SEQ ID NO: 10, SEQ ID NO: 12 or SEQ ID NO: 14 include DNAs comprising a base sequence having at least about 70% homology, preferably at least about 80% homology, more preferably at least about 90% homology and the most preferably at least about 95% homology, to the base sequence represented by any of SEQ ID NO:
20 10, SEQ ID NO: 12 or SEQ ID NO: 14.

Homology in the base sequence can be measured under the following conditions (an expectation value = 10; gaps are allowed; filtering = ON; match score = 1; mismatch score = -3) using the homology scoring algorithm NCBI BLAST (National Center for Biotechnology Information Basic Local Alignment Search
25 Tool).

The hybridization can be carried out by per se publicly known methods or by modifications of these methods, for example, according to the method described in Molecular Cloning, 2nd (J. Sambrook et al., Cold Spring Harbor Lab. Press, 1989), etc. A commercially available library may also be used according to the
30 instructions of the attached manufacturer's protocol. Preferably, the hybridization can be carried out under highly stringent conditions.

The highly stringent conditions used herein are, for example, those in a sodium concentration at about 19 to 40 mM, preferably about 19 to 20 mM at a temperature of about 50 to 70°C, preferably about 60 to 65°C. In particular,
35 hybridization conditions in a sodium concentration of about 19 mM at a temperature

of about 65°C are most preferred.

More specifically, as the DNA encoding the human metastin receptor consisting of the amino acid sequence represented by SEQ ID NO: 9, the DNA consisting of the base sequence represented by SEQ ID NO: 10 is used.

5 As the DNA encoding the rat metastin receptor consisting of the amino acid sequence represented by SEQ ID NO: 11, the DNA consisting of the base sequence represented by SEQ ID NO: 12 is used.

As the DNA encoding the mouse metastin receptor consisting of the amino acid sequence represented by SEQ ID NO: 13, the DNA consisting of the base
10 sequence represented by SEQ ID NO: 14 is used.

The metastin receptors, their partial peptides or salts thereof and the DNAs encoding the metastin receptors or their partial peptides can be obtained or produced by the methods described in WO 00/24890 or WO 01/75104.

15 The present invention will be described in detail by referring to EXAMPLES, FORMULATION EXAMPLES AND TEST EXAMPLES, but is not deemed to be limited thereto, and any modification may be made without departing from the scope of the present invention.

In the following EXAMPLES, the term "room temperature" normally means
20 a temperature of about 10°C to about 35°C. In percentages, the yield is shown by mol/mol% and the solvent used in chromatography by vol%, and the remaining by wt%. In proton NMR spectra, data on OH, NH protons, etc. that are broad and unidentified are not shown.

The other abbreviations used in the specification mean as follows.

25	Abbreviation	Description
	10Ψ,CSNH	: C-terminal-CONH ₂ at the 10-position is substituted with -CSNH ₂ .
	1Ψ2,CH ₂ NH	: The -CONH- bond between the 1- and 2-positions is substituted with the -CH ₂ NH-bond.
	2Ψ3,CH ₂ NH	: The -CONH- bond between the 2- and 3-positions is substituted with the -CH ₂ NH-bond.
30	3Ψ4,CH ₂ NH	: The -CONH- bond between the 3- and 4-positions is substituted with the -CH ₂ NH- bond.
	4Ψ5,CH ₂ NH	: The -CONH- bond between the 4- and 5-positions is substituted with the -CH ₂ NH- bond.
35	6Ψ7,CSNH	: The -CONH- bond between the 6- and 7-positions is substituted

- with the -CSNH- bond.
- 6Ψ7,NHCO : The -CONH- bond between the 6- and 7-positions is substituted with the -NHCO- bond.
- 6Ψ7,CH₂NH : The -CONH- bond between the 6- and 7-positions is substituted with the -CH₂NH- bond.
- 5 6Ψ7,CH₂O : The -CONH- bond between the 6- and 7-positions is substituted with the -CH₂O- bond.
- 7Ψ8,CH₂NH : The -CONH- bond between the 7- and 8-positions is substituted with the -CH₂NH- bond.
- 10 8Ψ9,CH₂NH : The -CONH- bond between the 8- and 9-positions is substituted with the -CH₂NH- bond.
- 9Ψ10,CH₂NH : The -CONH- bond between the 9- and 10-positions is substituted with the -CH₂NH- bond.
- Abu : 2-aminobutanoic acid
- 15 Abz(2) : 2-aminobenzoic acid
- Abz(3) : 3-aminobenzoic acid
- Ac : acetyl
- AcONB : N-acetoxy-5-norbornene-2,3-dicarboximide
- Acp : 6-aminocaproic acid
- 20 AcOEt : ethyl acetate
- AcOH : acetic acid
- Aib : α-aminoisobutanoic acid
- Ala(2-Qui) : 2-quinolylalanine
- Ala(3-Bzt) : 3-benzothienylalanine
- 25 Ala(cPr) : cyclopropylalanine
- Ala(Pip) : (4-piperidin-1-yl)alanine
- Alb : Albizziin 2-amino-3-ureidopropionic acid
- Ambz(4) : 4-aminomethylbenzoyl
- Arg(Ac) : N^ω-acetylarginine
- 30 Arg(Boc₂,Me) : N^{ω,ω'}-bis-tert-butoxycarbonyl-N^ω-methylarginine
- Arg(Et) : N^ω-ethylarginine
- Arg(Me) : N^ω-methylarginine
- Arg(asyMe₂) or Arg(Me₂)asym : asymmetric-N^{ω,ω'}-dimethylarginine
- Arg(symMe₂) or Arg(Me₂)sym : symmetric-N^{ω,ω'}-dimethylarginine
- 35 Arg(NO₂) : N^ω-nitroarginine

	Arg(Pbf)	: N ^ω -2,2,4,6,7-pentamethyldihydrobenzofuransulfonylarginine
	Arg(n-Pr)	: N ^ω -propylarginine
	Arg(Tos)	: N ^ω -tosylarginine
	Asp(NHMe)	: N ^ω -methylasparagine
5	Asp(NMe ₂)	: N ^{ω,ω} -dimethylasparagine
	Asp(NHPen)	: N ^ω -pentylasparagine
	Asp(NHcPr)	: N ^ω -cyclopropylasparagine
	Asp(NHBzl)	: N ^ω -benzylasparagine
	AzaGly	: azaglycine
10	AzaPhe	: azaphenylalanine
	Aze(2)	: azetidine-2-carboxylic acid
	Aze(3)	: azetidine-3-carboxylic acid
	β-Ala	: β-alanine
	Boc	: tert-butoxycarbonyl
15	Boc ₂ O	: di-tert-butyl dicarbonate
	Br-Z	: 2-bromobenzyloxycarbonyl
	Bu ^t	: tert-butyl
	Bzl	: benzyl
	CDI	: 1,1'-carbonyldiimidazole
20	Cha	: cyclohexylalanine
	CIP	: 2-chloro-1,3-dimethylimidazolium tetrafluoroborate
	Cit	: citrulline
	Clt resin	: 2-chlorotrytyl resin
	Cl-Z	: 2-chlorobenzyloxycarbonyl
25	cPr	: cyclopropyl
	Dab	: 2,4-diaminobutanoic acid
	Dap	: 2,3-diaminopropionic acid
	Dap(Ac)	: N ^β -acetyl-β-diaminopropionic acid
	Dap(For)	: N ^β -formyl-β-diaminopropionic acid
30	Dap(Gly)	: N ^β -glycyl-β-diaminopropionic acid
	Dap(GnGly)	: N ^β -(N-guanidinoglycyl)-β-diaminopropionic acid
	DCM	: dichloromethane
	DEA	: diethylamine
	DIEA	: N,N-diisopropylethylamine
35	DIPCDI	: 1,3-diisopropylcarbodiimide

	DMAP	: 4-dimethylaminopyridine
	DMF	: N,N-dimethylformamide
	EDT	: 1,2-ethanedithiol
	Fmoc	: 9-fluorenylmethoxycarbonyl
5	For	: formyl
	γ -Abu	: 4-aminobutanoic acid
	γ -MeLeu	: γ -methylleucine
	Gn	: guanidino
	GuAmb	: 4-guanidinomethylbenzoyl
10	Har	: homoarginine
	Har(Me)	: N ^ω -methylhomoarginine
	His(3Me)	: 3-methylhistidine π -methylhistidine
	HOAt	: 1-hydroxy-7-azabenzotriazole
	HOBt	: 1-hydroxybenzotriazole
15	HONB	: N-hydroxy-5-norbornene-2,3-dicarboximide
	Hph	: homophenylalanine
	cisHyp	: cis-4-hydroxyproline
	Hyp	: trans-4-hydroxyproline
	Hyp(Bzl)	: O-benzyl-trans-4-hydroxyproline
20	IndPr	: 3-(indole-3-yl)propionyl
	Izc	: imidazolidine-2-carboxylic acid
	Lys(Me ₂)	: N ^{ε,ε} -dimethyllysine
	MBHA	: p-methylbenzhydramine
	MeOH	: methanol
25	Mtt	: 4-methyltryl
	N((CH ₂) ₃ Gn)Gly	: N-(3-guanidinopropyl)glycine
	Nal(1)	: 1-naphthylalanine
	Nal(2)	: 2-naphthylalanine
	Nar	: norarginine
30	Nar(Me)	: N ^ω -methylnorarginine
	Nle	: norleucine
	NMeAla	: N ^α -methylalanine
	NMeArg	: N ^α -methylarginine
	NMeAsn	: N ^α -methylasparagine
35	NMeLeu	: N ^α -methylleucine

	NMePhe	: N ^α -methylphenylalanine
	NMeSer	: N ^α -methylserine
	NMeTrp	: N ^α -methyltryptophan
	NMeTyr	: N ^α -methyltyrosine
5	Nva	: Norvaline
	OBu ^t	: tert-butoxy
	Orn	: ornithine
	Orn(Mtt)	: N ^δ -(4-methyltrytyl)ornithine
	PAL	:
10	5-(4-(9-fluorenylmethoxycarbonyl)aminomethyl-3,5-dimethoxyphenoxy)valeric acid	
	Pbf	: 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl
	pGlu	: pyroglutamic acid
	Phe(2Cl)	: 2-chlorophenylalanine
	Phe(2F)	: 2-fluorophenylalanine
15	Phe(2Me)	: 2-methylphenylalanine
	Phe(3,4Cl ₂)	: 3,4-dichlorophenylalanine
	Phe(3,4F ₂)	: 3,4-difluorophenylalanine
	Phe(3CF ₃)	: 3-trifluoromethylphenylalanine
	Phe(3Cl)	: 3-chlorophenylalanine
20	Phe(3F)	: 3-fluorophenylalanine
	Phe(3Me)	: 3-methylphenylalanine
	Phe(4Cl)	: 4-chlorophenylalanine
	Phe(4CN)	: 4-cyanophenylalanine
	Phe(4F)	: 4-fluorophenylalanine
25	Phe(4Gn)	: 4-guanidinophenylalanine
	Phe(4NH ₂)	: 4-aminophenylalanine
	Phe(4NO ₂)	: 4-nitrophenylalanine
	Phe(4CN)	: 4-cyanophenylalanine
	Phe(4Me)	: 4-methylphenylalanine
30	Phe(F ₅)	: pentafluorophenylalanine
	αMePhe	: α-methylphenylalanine
	PheΨ(CH ₂ O)Gly	: The -CONH- bond between Phe and Gly is substituted with the -CH ₂ O- bond.
	PheΨ(CSNH)-NH ₂	: The C-terminal phenylalanylamide is substituted with the
35		phenylalanylthioamide.

	Phg	: phenylglycine
	PhOH	: phenol
	PhSMe	: thioanisole
	Pic(2)	: pipecolinic acid
5	Pic(3)	: 3-piperidinecarboxylic acid
	Pip(2)	: 2-aminopipecolinic acid
	Pro	: proline
	Pro(4F)	: trans-4-fluoroproline
	Pro(4NH ₂)	: cis-4-aminoproline
10	Pya(2)	: 2-pyridylalanine
	Pya(3)	: 3-pyridylalanine
	Pya(4)	: 4-pyridylalanine
	PyAOP	: (7-azabenzotriazole-1-yloxy)-tris(pyrrolidino)phosphonium hexafluorophosphate
15	PyBOP	: (benzotriazol-1-yloxy)-tris(pyrrolidino)phosphonium hexafluorophosphate
	PyBrop	: bromo-tris(pyrrolidino)phosphonium hexafluorophosphate
	Pzc(2)	: piperazine-2-carboxylic acid
	Sar	: N-methylglycine
20	Ser(Ac)	: O-acetylserine
	Ser(Me)	: O-methylserine
	Ser(3Phenyl)	: 3-phenylserine
	Thi	: 2-thienylalanine
	Thz	: thioproline
25	Tic	: 1,2,3,4-tetrahydroisoquinoline-2-carboxylic acid
	TIS	: triisopropylsilane
	Tle	: tert-leucine
	Tos	: tosyl
	Trp(For)	: N ⁱⁿ -formyltryptophan
30	Trt	: trytyl
	Tyr(Me)	: O-methyltyrosine
	Tyr(PO ₃ H ₂)	: O-phosphotyrosine
	TyrΨ(CH ₂ NH)Asn	: The -CONH- bond between Tyr and Asn is substituted with the -CH ₂ NH- bond.
35	TFA	: trifluoroacetic acid

TFE : trifluoroethanol
 Z : benzyloxycarbonyl

In the specification, where the codes of bases and amino acids are denoted
 5 by abbreviations, they are based on the abbreviations in accordance with the
 IUPAC-IUB Commission on Biochemical Nomenclature or the common codes in the
 art, examples of which are shown below. For amino acids that may have the optical
 isomer, L form is presented unless otherwise indicated.

10 DNA : deoxyribonucleic acid
 cDNA : complementary deoxyribonucleic acid
 A : adenine
 T : thymine
 G : guanine
 15 C : cytosine
 Y : thymine or cytosine
 N : thymine, cytosine, adenine or guanine
 R : adenine or guanine
 M : cytosine or adenine
 20 W : thymine or adenine
 S : cytosine or guanine
 RNA : ribonucleic acid
 mRNA : messenger ribonucleic acid
 dATP : deoxyadenosine triphosphate
 25 dTTP : deoxythymidine triphosphate
 dGTP : deoxyguanosine triphosphate
 dCTP : deoxycytidine triphosphate
 ATP : adenosine triphosphate
 EDTA : ethylenediaminetetraacetic acid
 30 SDS : sodium dodecyl sulfate
 TFA : trifluoroacetic acid
 EIA : enzyme immunoassay
 Gly or G : glycine
 Ala or A : alanine
 35 Val or V : valine

	Leu or L	: leucine
	Ile or I	: isoleucine
	Ser or S	: serine
	Thr or T	: threonine
5	Cys or C	: cysteine
	Met or M	: methionine
	Glu or E	: glutamic acid
	Asp or D	: aspartic acid
	Lys or K	: lysine
10	Arg or R	: arginine
	His or H	: histidine
	Phe or F	: phenylalanine
	Tyr or Y	: tyrosine
	Trp or W	: tryptophan
15	Pro or P	: proline
	Asn or N	: asparagine
	Gln or Q	: glutamine
	pGlu	: pyroglutamic acid

20 The sequence identification numbers in the sequence listing of the specification indicates the following sequence, respectively.

[SEQ ID NO: 1]

This shows the amino acid sequence of human-derived metastin (Metastin).

25 [SEQ ID NO: 2]

This shows the base sequence of DNA encoding human metastin.

[SEQ ID NO: 3]

This shows the amino acid sequence of mouse metastin precursor (A).

[SEQ ID NO: 4]

30 This shows the base sequence of DNA encoding mouse metastin precursor (A), which is the base sequence in plasmid pCMV-mKiSS-1 harbored on transformant Escherichia coli DH10B/pCMV-mKiSS-1.

[SEQ ID NO: 5]

This shows the amino acid sequence of mouse metastin precursor (B).

35 [SEQ ID NO: 6]

This shows the base sequence of DNA encoding mouse metastin precursor (B), which is the base sequence in plasmid pCR2.1-mKiSS-1.4A harbored on transformant *Escherichia coli* DH5 α /pCR2.1-mKiSS-1.4A.

[SEQ ID NO: 7]

5 This shows the amino acid sequence of rat-derived metastin precursor.

[SEQ ID NO: 8]

This shows the base sequence of DNA encoding rat metastin precursor.

[SEQ ID NO: 9]

10 This shows the amino acid sequence of human OT7T175 (metastin receptor).

[SEQ ID NO: 10]

This shows the base sequence of DNA encoding human OT7T175 (metastin receptor).

[SEQ ID NO: 11]

15 This shows the amino acid sequence of rat OT7T175 (metastin receptor).

[SEQ ID NO: 12]

This shows the base sequence of DNA encoding rat OT7T175 (metastin receptor).

[SEQ ID NO: 13]

20 This shows the amino acid sequence of mouse OT7T175 (metastin receptor).

[SEQ ID NO: 14]

This shows the base sequence of DNA encoding mouse OT7T175 (metastin receptor).

25 [SEQ ID NO: 15]

This shows the amino acid sequence of human metastin 15 (40-54).

[SEQ ID NO: 16]

This shows the amino acid sequence of human metastin 10 (45-54) (MS10).

[SEQ ID NO: 17]

30 This shows the amino acid sequence of human metastin 9 (46-54).

[SEQ ID NO: 18]

This shows the amino acid sequence of human metastin 8 (47-54).

[SEQ ID NO: 19]

This shows the base sequence of DNA encoding human metastin 15 (40-54).

35 [SEQ ID NO: 20]

This shows the base sequence of DNA encoding human metastin 10 (45-54).
[SEQ ID NO: 21]

This shows the base sequence of DNA encoding human metastin 9 (46-54).
[SEQ ID NO: 22]

5 This shows the base sequence of DNA encoding human metastin 8 (47-54).

The transformant *Escherichia coli* DH10B/pCMV-mKiSS-1 has been on deposit since January 24, 2000 with International Patent Organisms Depository, National Institute of Advanced Industrial Science and Technology (the former
10 Ministry of International Trade and Industry, Agency of Industrial Science and Technology, National Institute of Bioscience and Human Technology (NIBH)), located at Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki (postal code 305-8566), Japan, as the Accession Number FERM BP-7003 and since December 16, 1999 with Institute for Fermentation (IFO), located at 2-17-85, Juso-Honmachi, Yodogawa-ku,
15 Osaka-shi, Osaka, Japan, as the Accession Number IFO 16348.

The transformant *Escherichia coli* DH5 α /pCR2.1-mKiSS-1.4A has been on deposit since March 6, 2000 with International Patent Organisms Depository, National Institute of Advanced Industrial Science and Technology (the former
20 Ministry of International Trade and Industry, Agency of Industrial Science and Technology, National Institute of Bioscience and Human Technology (NIBH)), located at Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki (postal code 305-8566), Japan, as the Accession Number FERM BP-7073 and since February 16, 2000 with Institute for Fermentation (IFO), located at 2-17-85 Juso-Honmachi, Yodogawa-ku, Osaka-shi, Osaka, Japan, as the Accession Number IFO 16360.

25

EXAMPLES

(REFERENCE EXAMPLE 1)

Production of N-methyl-N,N'-Bis-Boc-1-guanylpurazole

Under a nitrogen flow, 720 mg of 60% NaH in oil was dissolved in 20 mL
30 of dry DMF and 20 mL of dry DMF solution of 5.59 g of N,N'-Bis-Boc-1-guanylpurazole commercially available was added to the solution at 0°C, followed by stirring for 10 minutes. After 1.68 mL of methyl iodide was added thereto, the mixture was stirred at room temperature for 24 hours. After the solvent was removed by distillation, the residue was dissolved in AcOEt and the
35 solution washed with 1N HCl aq. solution, satd. NaHCO₃ aq. solution and then satd.

NaCl aq. solution. After drying over Na_2SO_4 , the solvent was concentrated and the concentrate was purified by flush column chromatography (ethyl acetate/n-hexane = 1/4) using silica gel 60 (200 mL) to give 5.35 g (yield 91.6%) of N-methyl-N,N'-bis-Boc-1-guanylpurazole as a colorless oil.

5 ^1H NMR (300 MHz, CDCl_3): δ 8.00 (br s, 1H), 7.69 (br s, 1H), 6.42 (dd, 1H, $J=2.7$, 1.5 Hz), 3.25 (s, 3H), 1.53 (s, 9H), 1.30 (s, 9H)

Elemental analysis as $\text{C}_{15}\text{H}_{24}\text{N}_4\text{O}_4$

Calcd.: C, 55.54; H, 7.46; N, 17.27

Found: C, 55.36; H, 7.48; N, 17.06

10 Rf1: 0.64, Rf2: 0.79

Developing solvent for TLC:

Rf1 (ethyl acetate/n-hexane = 1/2), Rf2 (methanol/chloroform = 2/98)

Elution time on HPLC: 26.7 mins.

15 Elution conditions:

Column: Wakosil-II 5C18 HG (4.6 x 100 mm)

Linear density gradient elution with eluants A/B = 100/0-20/80, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (40 mins.)

Flow rate: 1.0 ml/min.

20

(REFERENCE EXAMPLE 2)

Production of N-methyl-N,N'-Bis-Z-1-guanylpurazole

In an argon atmosphere, 40 mg of 60% NaH in oil was dissolved in 5 mL of dry DMF and 5 mL of dry DMF solution of 380 mg of N,N'-Bis-Z-1-guanylpurazole commercially available was added to the solution at 0°C, followed by stirring for 10 minutes. After 125 μL of methyl iodide was added thereto, the mixture was stirred at room temperature for 15 hours. After the solvent was distilled off, the residue was dissolved in AcOEt and the solution washed with 1N HCl aq. solution, satd. NaHCO_3 aq. solution and then satd. NaCl aq. solution. After drying over Na_2SO_4 , the solvent was concentrated to give 393 mg of the crude product. From the crude product, 170 mg was purified by flush column chromatography (ethyl acetate/n-hexane = 1/4) using silica gel 60 (75 mL) to give 353 mg (yield 89.5%) of N-methyl-N,N'-bis-Z-1-guanylpurazole as a colorless oil.

35 ^1H NMR (300 MHz, CDCl_3): δ 7.97 (br s, 1H), 7.61 (d, 1H, $J=1.0\text{Hz}$), 7.37-7.32 (m, 4H), 7.29-7.26 (m, 4H), 7.16-7.13 (m, 2H), 6.36 (dd, 1H, $J=2.8$, 1.6 Hz), 5.18 (s, 2H),

5.04 (s, 2H), 3.22 (s, 3H)

Elemental analysis as $C_{21}H_{20}N_4O_4$

Calcd.: C, 64.28; H, 5.14; N, 14.28

Found: C, 64.09; H, 5.24; N, 14.43

5 Rf1: 0.50, Rf2: 0.86

Developing solvent for TLC:

Rf1 (ethyl acetate/n-hexane=1/2)

Rf2 (methanol/chloroform=2/98)

Elution time on HPLC: 28.9 mins.

10 Elution conditions:

Column: Wakosil-II 5C18 HG (4.6 x 100 mm)

Linear density gradient elution with eluants A/B = 100/0-20/80, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (40 mins.)

Flow rate: 1.0 ml/min.

15

(EXAMPLE 1)

(Synthesis A): Production of

des(1)-Ac-[D-Tyr²,D-Trp³,Thr⁵,AzaGly⁷,D-Arg⁹,Trp¹⁰]MS10 (Compound No. 708).

- 20 Trp(Boc), D-Arg(Pbf) and Leu were introduced in this order into 178 mg of Rink Amide MBHA Resin (0.56 mmol/g) on an ABI 433A peptide synthesizer to produce H-Leu-D-Arg(Pbf)-Trp(Boc)-Rink Amide MBHA resin. Separately, 116.3 mg (0.4 mmol) of Fmoc-NHNH₂·HCl was suspended in 1 mL of DMF, and under ice cooling a suspension of 61.6 mg (0.38 mmol) of CDI in 10 ml of THF was added
- 25 thereto. Subsequently, 139.4 μ l (0.8 mmol) of DIEA was added to the mixture, followed by stirring at room temperature for an hour. The resulting reaction solution was added to the H-Leu-D-Arg(Pbf)-Trp(Boc)-Rink Amide MBHA resin described above, followed by stirring at room temperature for 15 hours. After completion of the reaction, the resin was washed and, Phe, Thr(Bu^t) and Asn(Trt)
- 30 were introduced in this order into the resulting Fmoc-AzaGly-Leu-D-Arg(Pbf)-Trp(Boc)-Rink Amide MBHA resin on ABI 433A. The resin was divided into halves, one of which was taken out and the remaining half was applied again on ABI 433A, whereby D-Trp(Boc) and D-Tyr(Bu^t) were introduced in this order to give the
- 35 H-D-Tyr(Bu^t)-D-Trp(Boc)-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-D-Arg(Pbf)-Trp(Boc)

-)-Rink Amide MBHA resin. Subsequently, the product was treated for 20 minutes in 3 ml of DMF with 9.4 μ l (0.1 mmol) of Ac_2O and 17.4 μ l (0.1 mmol) of DIEA for N-terminal acetylation. The resin was washed and dried to give 202.2 mg of the Ac-D-Tyr(Bu^t)-D-Trp(Boc)-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-D-Arg(Pbf)-Trp(Bo
- 5 c)-Rink Amide MBHA resin. To the resin obtained, 1.5 mL of TFA/PhSMe/m-cresol/ H_2O /TIS/EDT (80/5/5/5/2.5/2.5) was added and the mixture was stirred for 90 minutes. Diethyl ether was added to the reaction solution, the resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an
- 10 aqueous acetic acid solution and the extract was filtered to remove the resin. Then, linear density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with eluants A/B: 70/30-60/40 using: eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the
- 15 product were collected and lyophilized to give 16.2 mg of white powders. Mass spectrum ($\text{M}+\text{H}^+$) 1284.9 (calcd. 1284.6) Elution time on HPLC: 13.3 mins. Elution conditions:
- Column: YMC-AM301 (4.6 x 100 mm)
- 20 Linear density gradient elution with eluants A/B = 80/20 - 30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.) Flow rate: 1.0 ml/min.

(EXAMPLE 2)

- 25 Production of Fmoc-AzaGly-Leu-Arg(Boc_2 , Me)-Trp(Boc)-Rink Amide MBHA resin
- After 5 g (0.4 mmol/g) of commercially available Rink Amide MBHA resin was swollen in DMF, the resin was treated with 50 ml of 20% piperidine/DMF solution for 20 minutes to remove the Fmoc group. The resin obtained was washed with DMF and treated with 4.213 g (8 mmol) of Fmoc-Trp(Boc)-OH, 1.272 mL (8
- 30 mmol) of DIPCDI and 16 mL (8 mmol) of 0.5M HOAt/DMF solution at room temperature for 90 minutes, whereby Trp(Boc) was introduced to give the Fmoc-Trp(Boc)-Rink Amide MBHA resin. In a similar manner, Orn(Mtt) was introduced to give 2 mmol of the Fmoc-Orn(Mtt)-Trp(Boc)-Rink Amide MBHA resin. After the resin obtained was washed and swollen with DCM, 50 mL of
- 35 TFA/TIS/DCM (1/5/94) was added thereto, followed by shaking for 10 minutes and

removing the solution through filtration. This procedure was repeated until yellow color caused by free Mtt group in a TFA/TIS/DCM (1/5/94) solution disappeared when the solution was added; thus the Mtt group was removed.

The resulting Fmoc-Orn-Trp(Boc)-Rink Amide MBHA resin was
5 neutralized with 5%-DIEA/DCM solution. After washing with DCM, 25 mL of DCM-TFE (4:1) and 1.946 g (6 mmol) of N-methyl-N',N'-bis-Boc-1-guanylpurazole obtained in REFERENCE EXAMPLE 1 were added to the resin. DIEA was added to the mixture to adjust pH of the resulting solution to 10. The solution was shaken for 15 hours to give 6.195 g of Fmoc-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA
10 resin. Fmoc-Leu was introduced into the obtained resin as in the same manner described above. The resin was divided in halves and the Fmoc group was removed from the thus obtained Fmoc-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin (1 mmol) to give H-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin (1 mmol).

15 Separately, 1.745 g (6 mmol) of Fmoc-NHNH₂ HCl was suspended in 20 mL of DMF-THF (4:1). Under ice cooling, 973 mg (6 mmol) of CDI and 2.09 mL (12 mmol) of DIEA were added to the suspension, followed by stirring at room temperature for an hour. The resulting reaction solution was added to the H-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin described above, and the
20 mixture was stirred at room temperature for 15 hours. After completion of the reaction, the resin was washed and dried to give 3.314 g of the Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin.

(EXAMPLE 3)

25 (Synthesis B): Production of des(1-3)-Ac-[Thr5,AzaGly7,Arg(Me)9,Trp10]MS10 (Compound No. 709)

After 5.455 g (0.55 mmol/g) of commercially available Rink Amide MBHA resin was swollen in DMF, the resin was treated with 50 ml of 20% piperidine/DMF solution for 20 minutes to remove the Fmoc group. The resin obtained was washed
30 with DMF and then treated with 6.319 g (12 mmol) of Fmoc-Trp(Boc)-OH, 1.908 mL (12 mmol) of DIPCDI and 24 mL (12 mmol) of 0.5M HOAt/DMF solution at room temperature for 90 minutes, whereby Trp(Boc) was introduced to give the Fmoc-Trp(Boc)-Rink Amide MBHA resin. In a similar manner, Orn(Mtt) was introduced to give 3 mmol of the Fmoc-Orn(Mtt)-Trp(Boc)-Rink Amide MBHA
35 resin. The resin obtained was washed with DCM and swollen, and then 50 mL of

TFA/TIS/TFE/DCM (1/5/19/75) was added to the resin, followed by shaking for 10 minutes and removing the solution through filtration. This procedure was repeated until yellow color caused by free Mtt group in a TFA/TIS/TFE/DCM (1/5/19/75) solution disappeared when the solution was added thereto; thus, the Mtt group was removed.

The resulting Fmoc-Orn-Trp(Boc)-Rink Amide MBHA resin was neutralized with 5%-DIEA/DCM solution. After washing with DCM, 20 mL of DCM-TFE (4:1) and 2.919 g (9 mmol) of N-methyl-N',N'-bis-Boc-1-guanylpurazole obtained in REFERENCE EXAMPLE 1 were added to the resin. DIEA was added to the mixture to adjust pH of the solution to 10. The mixture was shaken for 15 hours to give the Fmoc-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin. Fmoc-Leu was introduced into the obtained resin as in the same manner described above. The Fmoc group was then removed from the thus obtained Fmoc-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin (3 mmol) to give the H-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin (3 mmol).

Separately, 3.489 g (12 mmol) of Fmoc-NHNH₂.HCl was suspended in 20 mL of DMF. Under ice cooling, a suspension of 1.849 g (11.4 mmol) of CDI in 20 mL of THF was added and then 4.181 mL (24 mmol) of DIEA was added to the mixture, followed by stirring at room temperature for an hour. The resulting reaction solution was added to the H-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin described above, followed by stirring at room temperature for 15 hours. After completion of the reaction, the resin was washed and dried to give the Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin.

The resin obtained was swollen in DMF and then treated with 30 mL of 20% piperidine/DMF solution for 20 minutes to remove the Fmoc group. After the resin obtained was washed with DMF, the resin was treated with 5.419 g (12 mmol) of Trt-Phe-OH.0.5AcOEt, 6.257 g (12 mmol) of PyAOP, 24 mL (12 mmol) of 0.5M HOAt/DMF and 7.316 mL (42 mmol) of DIEA at room temperature for 90 minutes to give the Trt-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin. After the resin obtained was washed with DCM and swollen, 50 mL of TFA/TIS/TFE/DCM (1/5/19/75) was added to the resin, followed by shaking for 10 minutes and removing the solution through filtration. This procedure was repeated until yellow brown color caused by the free Trt group in a TFA/TIS/TFE/DCM (1/5/19/75) solution disappeared when the solution was added; thus the Trt group was removed. The H-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide

MBHA resin obtained was neutralized with 5%-DIEA/DMF solution and washed in DMF. Thereafter, the resin was treated with 4.780 g (12 mmol) of Fmoc-Thr(Bu^t)-OH, 1.908 mL (12 mmol) of DIPCDI and 24 mL (12 mmol) of 0.5M HOAt/DMF at room temperature for 90 minutes to introduce Thr(Bu^t).

- 5 Subsequently, the Fmoc deprotection through treatment with 50 ml of 20% piperidine/DMF solution for 20 minutes and condensation by the DIPCDI/HOAt method as in the introduction of Thr(Bu^t) were performed to introduce Asn(Trt). The resin was then washed and dried to give 10.624 g of the Fmoc-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin. After 100 mg (0.03 mmol) of the resulting resin was swollen in 10 DMF, the resin was treated with 3 ml of 20% piperidine/DMF solution for 20 minutes to remove the Fmoc group. The resin obtained was washed with DMF and treated in 1 ml of DMF with 9.4 μ L (0.1 mmol) of Ac₂O and 17.4 μ L (0.1 mmol) of DIEA at room temperature for 30 minutes for N-terminal acetylation. The resin 15 was washed and dried to give 94.2 mg of the Ac-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin.

- To the resin obtained, 0.75 mL of TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added and the mixture was stirred for 90 minutes. Diethyl- 20 ether was added to the reaction solution, the resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the extract was filtered to remove the resin. Thereafter, linear density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with eluants A/B: 76/24-66/34 using: 25 0.1% TFA in water and eluant B: eluant A: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the product were collected and lyophilized to give 2.8 mg of white powders.

Mass spectrum (M+H)⁺ 949.8 (Calcd. 949.5)

- 30 Elution time on HPLC: 10.2 mins.

Elution conditions:

Column YMC-AM301 (4.6 x 100 mm)

Linear density gradient elution with eluants A/B = 80/20-30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

- 35 Flow rate: 1.0 ml/min.

(EXAMPLE 4)

(Synthesis C): Production of des(1-2)-Ac-[Acp3,Thr5,AzaGly7,Arg(Me)9,Trp10] MS10 (Compound No. 713)

- 5 After 100 mg (0.03 mmol) of the
Fmoc-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu- Arg(Boc₂,Me)-Trp(Boc)-Rink Amide
MBHA resin was swollen in DMF, the resin was treated with 3 ml of 20%
piperidine/DMF solution for 20 minutes to remove the Fmoc group. The resin was
washed with DMF and then treated in 1 mL of DMF with 70.6 mg (0.2 mmol) of
10 Fmoc-Acp-OH, 104.2 mg (0.2 mmol) of PyAOP and 52.4 μ L (0.2 mmol) of DIEA at
room temperature for 90 minutes to introduce Acp; thus, the
Fmoc-Acp-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink
Amide MBHA resin was obtained. The obtained resin was treated with 3 ml of
20% piperidine/DMF solution and then washed with DMF to remove the Fmoc
15 group. Subsequently, the resin was treated in 1 mL of DMF with 9.4 μ L (0.1 mmol)
of Ac₂O and 17.4 μ L (0.1 mmol) of DIEA at room temperature for 30 minutes to
acetylate the N terminus. The treated resin was then washed and dried to give 101.2
mg of the
Ac-Acp-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide
20 MBHA resin. To the resin obtained, 0.75 mL of
TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by
stirring for 90 minutes. Diethyl ether was added to the reaction solution, the
resulting precipitate was centrifuged, and the supernatant was removed. This
procedure was repeated twice for washing. The residue was extracted with an
25 aqueous acetic acid solution and the extract was filtered to remove the resin. Then,
linear density gradient elution (60 minutes) was performed at a flow rate of 15
ml/min with eluants A/B: 77/23-67/33 using: eluant A: 0.1% TFA in water and eluant
B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack
R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the
30 product were collected and lyophilized to give 7.3 mg of white powders.
Mass spectrum (M+H)⁺ 1062.7 (Calcd. 1062.6)
Elution time on HPLC: 10.7 mins.
Elution conditions:

Column YMC-AM301 (4.6 x 100 mm)

- 35 Linear density gradient elution with eluants A/B = 80/20-30/70, using 0.1%

TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

Flow rate: 1.0 ml/min.

(EXAMPLE 5)

5 (Synthesis D): Production of
Ac-D-Tyr-D-Trp-Asp(NHPen)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound
No. 714)

Using 661 mg (0.25 mmol) of the
Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin, the peptide
10 chain was extended on a peptide synthesizer ABI 433A (Fmoc/DCC/HOBt) to give the
the
H-D-Tyr(Bu^t)-D-Trp(Boc)-Asp(OBu^t)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp
(Boc)-Rink amide MBHA resin. To the resin, 5 mL of DMF, 111 mg of AcONB
and 87 µL of DIEA were added, followed by shaking for 3 hours. The resin was
15 washed and then dried to give the Ac-D-Tyr(Bu^t)-D-Trp(Boc)-
Asp(OBu^t)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA
resin. To the 9/10 amount of the resin, 5 mL of
TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added and the mixture
was shaken for 2 hours and ether was added for precipitation. The washing
20 procedure with ether was repeated and then dried to give 218.4 mg of
Ac-D-Tyr-D-Trp-Asp-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂. To 32.5 mg of the
product, 1 mL of DMF, 5.7 µL of aminopentane, 13.5 mg of HOBt, 26.0 mg of
PyBOP and 26.1 µL of DIEA, and the mixture was stirred for 24 hours. After the
solvent was removed by distillation, ether was added for precipitation. The
25 washing procedure with ether was repeated and the dried. The residue was
dissolved in an aqueous acetic acid solution. After insoluble matters were removed
by filtration, linear density gradient elution (60 minutes) was performed with eluants
A/B: 65/35-55/45 using: eluant A: 0.1% TFA in water and eluant B: 0.1%
TFA-containing acetonitrile on preparative HPLC using YMC D-ODS-5-ST S-5
30 120A column (20 x 150 mm). The fractions containing the product were collected
and lyophilized to give 11.0 mg of white powders.

Mass spectrum (M+H)⁺ 1368.9 (Calcd. 1368.7)

Elution time on HPLC: 20.9 mins.

Elution conditions:

35 Column: Wakosil-II 5C18 (4.6 x 100 mm)

Linear density gradient elution with eluants A/B = 100/0-50/50, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)
Flow rate: 1.0 ml/min.

5 (EXAMPLE 6)

(Synthesis E): Production of

des(1)-Ac-[D-Tyr²,D-Trp³,Alb⁴,Thr⁵,AzaGly⁷,Arg(Me)⁹,Trp¹⁰]MS¹⁰ (Compound No. 717)

Using 132 mg (0.05 mmol) of the

- 10 Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin, the peptide chain was extended on a peptide synthesizer ABI 433A (Fmoc/DCC/HOBt) to give the
H-D-Tyr(Bu¹)-D-Trp(Boc)-Alb-Thr(Bu¹)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-
Rink amide MBHA resin. To the resin, 2 mL of DMF, 23 mg of AcONB and 17 μ L
15 of DIEA were added, followed by shaking for 3 hours. The resin was washed and then dried to give the Ac- D-Tyr(Bu¹)-D-Trp(Boc)- Alb-Thr(Bu¹)-Phe-AzaGly-Leu-
Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin. To the resin, 1 mL of
TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by
stirring for 2 hours. Ether was added to the reaction solution, the resulting
20 precipitate was centrifuged and the supernatant was removed. This procedure was repeated for washing. The residue was extracted with an aqueous acetic acid solution and insoluble matters were removed by filtration. Then, linear density
gradient elution (60 minutes) was performed with eluants A/B: 69/31-59/41 using:
eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on
25 preparative HPLC using YMC D-ODS-5-ST S-5 120A column (20 x 150 mm). The fractions containing the product were collected and lyophilized to give 4.8 mg of white powders.

Mass spectrum (M+H)⁺ 1313.9 (Calcd. 1313.7)

Elution time on HPLC: 18.3 mins.

- 30 Elution conditions:

Column: Wakosil-II 5C18 (4.6 x 100 mm)

Linear density gradient elution with eluants A/B = 100/0-50/50, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

Flow rate: 1.0 ml/min.

35

(EXAMPLE 7)

Production of des(1)-Ac-[D-Tyr²,Hyp³,Thr⁵,AzaGly⁷,Arg(Me)⁹,Trp¹⁰]MS¹⁰
(Compound No. 723)

The Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin,
5 800 mg (0.3 mmol), was used as a starting material, and the starting material was
reacted on peptide synthesizer ABI 433A (according to the Fmoc/DCC/HOBt 0.25
mmol protocol) to introduce Phe, Thr(Bu^t), Asn(Trt), Hyp(Bu^t) and D-Tyr(Bu^t) in
this order thereby to give the H-D-Tyr(Bu^t)-
Hyp(Bu^t)-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide
10 MBHA resin. Subsequently, the resin was treated for 20 minutes in DMF with 94.4
μl (1 mmol) of Ac₂O and 174.2 μl (1 mmol) of DIEA for N-terminal acetylation to
give 1.049 g of the
Ac-D-Tyr(Bu^t)-Hyp(Bu^t)-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Bo
c)-Rink amide MBHA resin. The same procedure was carried out again to give
15 1.035 g of the resin. To each resin, 8 mL of TFA/PhSMe/m-cresol/H₂O/TIS/EDT
(80/5/5/5/2.5/2.5) was added, followed by stirring for 90 minutes. Diethyl ether
was added to the reaction solution, the resulting precipitate was centrifuged, and the
supernatant was removed. This procedure was repeated twice for washing. The
residue was extracted with an aqueous acetic acid solution and the resin was removed
20 by filtration. Thereafter, linear density gradient elution (60 minutes) was performed
at a flow rate of 15 ml/min with eluants A/B: 76/24-66/34 using: eluant A: 0.1% TFA
in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using
YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions
containing the product were collected and lyophilized. The obtained white
25 powders, 159.5 mg, were dissolved in 200 mL of water and 550 μL of ion exchange
resin BioRAD AG1 x 8 AcO⁻ form was added to the solution. While manually
stirring the reaction solution sometimes, the solution was allowed to stand for an
hour. The solution was filtered through a membrane filter to remove the resin and
give 134.5 mg of white powders as the acetate. To the purified sample (the acetate)
30 obtained, 6.725 ml of glacial acetic acid was added and the mixture was sonicated for
5 minutes. Subsequently, 20.175 ml of pure water was added to prepare 5
mg/ml/25% aqueous acetic acid solution. The resulting 5 mg/ml solution was
dispensed by 4 ml each in six vials, and the rest of the solution was transferred to a
vial. These vials were frozen at -80°C for 2 hours and then lyophilized, in which
35 the lyophilization was performed while cooling at -40°C for an hour, -20°C for 2

hours, 0°C for 12 hours, 5°C for 8 hours and 20°C for 5 hours. As a result, six vials of 20 mg each and one vial of 12.34 mg were obtained.

Mass spectrum (M+H)⁺ 1225.9 (Calcd. 1225.6)

Anal. for amino acids (20% HCl containing 4% thioglycolic acid, 110°C, hydrolysis for 24 hours; figures in parentheses show theoretical values.): Asp 0.99 (1); Thr 0.96 (1); Leu 0.98 (1); Tyr 0.98 (1); Phe 1.00 (1)

Elution time on HPLC: 11.4 mins.

Elution conditions:

Column: YMC ODS AM-301 (4.6 x 100 mm) (4.6 x 100 mm)

10 Linear-density gradient elution with eluants A/B = 80/20-30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

Flow rate: 1.0 ml/min.

(EXAMPLE 8)

15 Production of des(1)-Ac-[D-Tyr², Gly³, Thr⁵, AzaGly⁷, Arg(Me)⁹, Trp¹⁰]MS10 (Compound No. 726)

The Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin, 661 mg (0.25 mmol), was used as a starting material, and the starting material was reacted on peptide synthesizer ABI 433A (according to the Fmoc/DCC/HOBt 0.25 mmol protocol) to introduce Phe, Thr(Bu^t), Asn(Trt), Gly and D-Tyr(Bu^t) in this order thereby to give the H-D-Tyr(Bu^t)-Gly-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin. Subsequently, the resin was treated for 20 minutes in DMF with 94.4 μl (1 mmol) of Ac₂O and 174.2 μl (1 mmol) of DIEA for N-terminal acetylation to give 866.6 mg of the Ac-D-Tyr(Bu^t)-Gly-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin. To the resin obtained, 8 mL of TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by stirring for 90 minutes. Diethyl ether was added to the reaction solution, the resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the resin was removed by filtration. Then, linear density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with eluants A/B: 76/24-66/34 using: eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B

20
25
30
35

- S-5, 120A column (30 x 250 mm). The fractions containing the product were collected and lyophilized. The obtained white powders, 42.9 mg, were dissolved in 50 mL of water and 153 μ L of ion exchange resin BioRAD AG1 x 8 AcO⁻ form was added to the solution. While manually stirring the reaction solution sometimes, the solution was allowed to stand for an hour. The solution was then filtered through a membrane filter to remove the resin and give 39.9 mg of white powders as the acetate. To the purified sample (the acetate) obtained, 1.995 ml of glacial acetic acid was added and the mixture was sonicated for 5 minutes. Subsequently, 5.985 ml of pure water was added to prepare 5 mg/ml/25% aqueous acetic acid solution.
- The resulting 5 mg/ml solution was dispensed in one vial of 4 ml, and the rest of the solution was transferred to a vial. These vials were frozen at -80°C for 2 hours and then lyophilized, in which the lyophilization was performed while cooling at -40°C for an hour, -20°C for 2 hours, 0°C for 12 hours, 5°C for 8 hours and 20°C for 5 hours. As a result, one vial of 20 mg and one vial of 19.08 mg prepared from the remaining solution were obtained.
- Mass spectrum (M+H)⁺ 1169.7 (Calcd. 1169.6)
- Anal. for amino acids (20% HCl containing 4% thioglycolic acid, 110°C, hydrolysis for 24 hours; figures in parentheses show theoretical values.): Asp 0.98 (1); Thr 0.93 (1); Gly 0.97 (1); Leu 0.94 (1); Tyr 0.97 (1); Phe 1.00 (1)
- Elution time on HPLC: 11.3 mins.
- Elution conditions:
- Column: YMC ODS AM-301 (4.6 x 100 mm)
- Linear density gradient elution with eluants A/B = 80/20-30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)
- Flow rate: 1.0 ml/min.

(EXAMPLE 9)

Production of des(1)-Ac-[D-Tyr², Aib³, Thr⁵, AzaGly⁷, Arg(Me)⁹, Trp¹⁰]MS10 (Compound No. 727)

- The Fmoc-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂, Me)-Trp(Boc)-Rink amide MBHA resin, 1325 mg (0.4 mmol), was used as a starting material, and the starting material was reacted by the manual solid phase synthesis (Fmoc/DIPCDI/HOAt) to introduce Aib and D-Tyr(Bu^t) in this order. Thus, the H-D-Tyr(Bu^t)-Aib-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂, Me)-Trp(Boc)-Rink amide MBHA resin was obtained.

Subsequently, 5 mL of DMF, 265 mg of AcONB and 209 μ L of DIEA were added and the mixture was shaken for 3 hours for N-terminal acetylation to give the Ac-D-Tyr(Bu^t)-Aib-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin. To the resin
5 obtained, 8 mL of TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by stirring for 90 minutes. Diethyl ether was added to the reaction solution, the resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the resin was removed by filtration. Then, linear
10 density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with eluants A/B: 72/28-62/38 using: eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the product were collected and lyophilized. The obtained white powders were dissolved in 50 mL of
15 water and 150 μ L of ion exchange resin BioRAD AG1 x 8 AcO⁻ form was added to the solution. While manually stirring the reaction solution sometimes, the solution was allowed to stand for an hour. The solution was then filtered through a membrane filter to remove the resin and give 41.9 mg of white powders as the acetate. To the purified sample (the acetate) obtained, 2.095 ml of glacial acetic
20 acid was added and the mixture was sonicated for 5 minutes. Subsequently, 6.285 ml of pure water was added to prepare 5 mg/ml/25% aqueous acetic acid solution. The resulting 5 mg/ml solution was dispensed in one vial of 4 ml and one vial of 3 ml, and the rest of the solution was transferred to a vial. These vials were frozen at -80°C for 2 hours and then lyophilized, in which the lyophilization was performed
25 while cooling at -40°C for an hour, -20°C for 2 hours, 0°C for 12 hours, 5°C for 8 hours and 20°C for 5 hours. As a result, one vial of 20 mg, one vial of 15 mg and one vial of 4.8 mg prepared from the remaining solution were obtained.

Mass spectrum (M+H)⁺ 1197.7 (Calcd. 1197.6)

Anal. for amino acids (20% HCl containing 4% thioglycolic acid, 110°C, hydrolysis
30 for 24 hours; figures in parentheses show theoretical values.): Asp 0.98 (1); Thr 0.94 (1); Leu 0.95 (1); Tyr 0.97 (1); Phe 1.00 (1)

Elution time on HPLC: 13.0 mins.

Elution conditions:

Column YMC-AM301 (4.6 x 100 mm)

35 Linear density gradient elution with eluants A/B = 80/20-30/70, using 0.1%

TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

Flow rate: 1.0 ml/min.

(EXAMPLE 10)

- 5 Production of des(1)-Ac-[D-Tyr²,Glu³,Thr⁵,AzaGly⁷,Arg(Me)⁹,Trp¹⁰]MS10
(Compound No. 746)

The Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin, 661 mg (0.25 mmol), was used as a starting material, and the starting material was reacted on a peptide synthesizer ABI 433A (according to the Fmoc/DCC/HOBt 0.25 mmol protocol), whereby Phe, Thr(Bu^t), Asn(Trt), Glu(OBu^t) and D-Tyr(Bu^t) were introduced in this order to give the H-D-Tyr(Bu^t)-Glu(OBu^t)-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin. Subsequently, the resin was treated for 20 minutes in DMF with 94.4 μ l (1 mmol) of Ac₂O and 174.2 μ l (1 mmol) of DIEA for N-terminal acetylation to give 940.0 mg of the Ac-D-Tyr(Bu^t)-Glu(OBu^t)-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin. To the resin obtained, 6 mL of TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by stirring for 90 minutes. Diethyl ether was added to the reaction solution, the resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the resin was removed by filtration. Then, linear density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with eluants A/B: 76/24-66/34 using: eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the product were collected and lyophilized. The obtained white powders, 57.3 mg, were dissolved in 100 mL of water and 192 μ L of ion exchange resin AG1 x 8 AcO⁻ form, which was obtained by converting commercially available BioRAD AG1 x 8 Cl⁻ form into the acetate type in a conventional manner, was added to the solution. While manually stirring the reaction solution sometimes, the solution was allowed to stand for an hour. The solution was then filtered through a membrane filter to remove the resin and the resin was lyophilized to give 42.7 mg of white powders. To the purified sample (the acetate) obtained, 2.135 ml of glacial acetic acid was added and the mixture was sonicated for 5 minutes. Subsequently, 6.405 ml of pure water was

added to prepare 5 mg/ml/25% aqueous acetic acid solution. The resulting solution was dispensed by 4 ml each in two vials, and the rest of the solution was transferred to a vial. These vials were frozen at -80°C for 2 hours and then lyophilized, in which the lyophilization was performed while cooling at -40°C for 4 hours, -20°C for 5 2 hours, 0°C for 12 hours, 5°C for 8 hours and 20°C for 5 hours. As a result, two vials of 20 mg each and one vial of 0.52 mg prepared from the remaining solution were obtained.

Mass spectrum (M+H)⁺ 1241.8 (Calcd. 1241.4)

Anal. for amino acids (20% HCl containing 4% thioglycolic acid, 110°C, hydrolysis 10 for 24 hours; figures in parentheses show theoretical values.): Asp 0.99 (1); Thr 0.95 (1); Glu 0.96 (1); Leu 0.98 (1); Tyr 0.98 (1); Phe 1.00 (1)

Elution time on HPLC: 11.4 mins.

Elution conditions:

Column: YMC ODS AM-301 (4.6 x 100 mm)

15 Linear density gradient elution with eluants A/B = 80/20-30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

Flow rate: 1.0 ml/min.

(EXAMPLE 11)

20 Production of
des(1)-Ac-[D-Tyr², Lys³, Thr⁵, Phe(3F)⁶, AzaGly⁷, Arg(Me)⁹, Trp¹⁰]MS10
(Compound No. 756)

The Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin, 661 mg (0.25 mmol), was used as a starting material, and the starting material was 25 reacted on ABI 433A peptide synthesizer (according to the Fmoc/DCC/HOBt 0.25 mmol protocol), whereby Phe(3F), Thr(Bu^t), Asn(Trt), Lys(Boc) and D-Tyr(Bu^t) were introduced in this order to give the
H-D-Tyr(Bu^t)-Lys(Boc)-Asn(Trt)-Thr(Bu^t)-Phe(3F)-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin. Subsequently, the resin was treated for 20 minutes 30 in DMF with 94.4 μl (1 mmol) of Ac₂O and 174.2 μl (1 mmol) of DIEA for N-terminal acetylation to give 881.7 mg of the
Ac-D-Tyr(Bu^t)-Lys(Boc)-Asn(Trt)-Thr(Bu^t)-Phe(3F)-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin. To the resin obtained, 6 mL of
TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by 35 stirring for 90 minutes. Diethyl ether was added to the reaction solution, the

resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the resin was removed by filtration. Then, linear density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with eluants A/B: 77/23-67/33 using: eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the product were collected and lyophilized. The obtained white powders, 47.7 mg, were dissolved in 50 mL of water and 316 μ L of ion exchange resin BioRAD AG1 x 8 AcO⁻ form was added to the solution. While manually stirring the reaction solution sometimes, the solution was allowed to stand for an hour. The solution was then filtered through a membrane filter to remove the resin and give 42.2 mg of white powders as the acetate. To the purified sample (the acetate) obtained, 2.11 ml of glacial acetic acid was added and the mixture was sonicated for 5 minutes. Subsequently, 6.33 ml of pure water was added to prepare 5 mg/ml/25% aqueous acetic acid solution. The resulting 5 mg/ml solution was dispensed in two vials of 4 ml each, and the rest of the solution was transferred to a vial. These vials were frozen at -80°C for 2 hours and then lyophilized, in which the lyophilization was performed while cooling at -40°C for an hour, -20°C for 2 hours, 0°C for 12 hours, 5°C for 8 hours and 20°C for 5 hours. As a result, two vials of 20 mg and one vial of 0.20 mg prepared from the remaining solution were obtained.

Mass spectrum (M+H)⁺ 1258.8 (Calcd. 1258.6)

Anal. for amino acids (20% HCl containing 4% thioglycolic acid, 110°C, hydrolysis for 24 hours; figures in parentheses show theoretical values.): Asp 0.99 (1); Thr 0.95 (1); Leu 0.95 (1); Tyr 0.99 (1); Phe(3F) 1.00 (1); Lys 0.97 (1)

Elution time on HPLC: 10.8 mins.

Elution conditions:

Column: YMC ODS AM-301 (4.6 x 100 mm)

Linear density gradient elution with eluants A/B = 80/20-30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

Flow rate: 1.0 ml/min.

(EXAMPLE 12)

Production of

des(1)-Ac-[D-Tyr², Glu³, Thr⁵, Phe(3F)⁶, AzaGly⁷, Arg(Me)⁹, Trp¹⁰]MS10

(Compound No. 757)

The Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin, 661 mg (0.25 mmol), was used as a starting material, and the starting material was reacted on ABI 433A peptide synthesizer (according to the Fmoc/DCC/HOBt 0.25 mmol protocol), whereby Phe(3F), Thr(Bu^t), Asn(Trt), Glu(OBu^t) and D-Tyr(Bu^t) were introduced in this order to give the H-D-Tyr(Bu^t)-Glu(OBu^t)-Asn(Trt)-Thr(Bu^t)-Phe(3F)-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin. Subsequently, the resin was treated for 20 minutes in DMF with 94.4 μ l (1 mmol) of Ac₂O and 174.2 μ l (1 mmol) of DIEA for N-terminal acetylation to give 872.5 mg of the Ac-D-Tyr(Bu^t)-Glu(OBu^t)-Asn(Trt)-Thr(Bu^t)-Phe(3F)-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin. To the resin obtained, 8 mL of TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by stirring for 90 minutes. Diethyl ether was added to the reaction solution, the resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the resin was removed by filtration. Then, linear density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with eluants A/B: 74.5/25.5-64.5/35.5 using eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the product were collected and lyophilized. The obtained white powders, 60.0 mg, were dissolved in 100 mL of water and 198 μ l of ion exchange resin AG1 x 8 AcO⁻ form, which was obtained by converting commercially available BioRAD AG1 x 8 Cl⁻ form into the acetate type in a conventional manner, was added to the solution. While manually stirring the reaction solution sometimes, the solution was allowed to stand for an hour. The solution was then filtered through a membrane filter to remove the resin and give 45.2 mg of white powders as the acetate. To the purified sample (the acetate) obtained, 2.26 ml of glacial acetic acid was added and the mixture was sonicated for 5 minutes. Subsequently, 6.78 ml of pure water was added to prepare 5 mg/ml/25% aqueous acetic acid solution. The resulting 5 mg/ml solution was dispensed by 4 ml each in two vials, and the rest of the solution was transferred to a vial. These vials were frozen at -80°C for 2 hours and then lyophilized, in which the lyophilization was performed while cooling at -40°C for an hour, -20°C for 2 hours, 0°C for 12 hours, 5°C for 8 hours and 20°C for 5 hours.

As a result, two vials of 20 mg and one vial of 3.81 mg prepared from the remaining solution were obtained.

Mass spectrum (M+H)⁺ 1259.9 (Calcd. 1259.6)

Anal. for amino acids (20% HCl containing 4% thioglycolic acid, 110°C, hydrolysis for 24 hours; figures in parentheses show theoretical values.): Asp 0.99 (1); Thr 0.94 (1); Glu 0.99 (1); Leu 0.94 (1); Tyr 0.97 (1); Phe(3F) 1.00 (1)

Elution time on HPLC: 11.9 mins.

Elution conditions:

Column: YMC ODS AM-301 (4.6 x 100 mm)

10 Linear density gradient elution with eluants A/B = 80/20-30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)
Flow rate: 1.0 ml/min.

(EXAMPLE 13)

15 Production of
des(1)-Ac-[D-Tyr²,Hyp³,Thr⁵,Phe(4F)⁶,AzaGly⁷,Arg(Me)⁹,Trp¹⁰]MS10
(Compound No. 787)

The Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin, 661 mg (0.25 mmol), was used as a starting material, and the starting material was
20 reacted on peptide synthesizer ABI 433A (according to the Fmoc/DCC/HOBt 0.25 mmol protocol), whereby Phe(4F), Thr(Bu^t), Asn(Trt), Hyp(Bu^t) and D-Tyr(Bu^t) were introduced in this order to give the
H-D-Tyr(Bu^t)-Hyp(Bu^t)-Asn(Trt)-Thr(Bu^t)-Phe(4F)-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin. Subsequently, the resin was treated for 20
25 minutes in DMF with 94.4 μl (1 mmol) of Ac₂O and 174.2 μl (1 mmol) of DIEA for N-terminal acetylation to give 832.8 mg of the
Ac-D-Tyr(Bu^t)-Hyp(Bu^t)-Asn(Trt)-Thr(Bu^t)-Phe(4F)-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA resin. The same procedure was carried out again to
give 823.9 mg of the resin. To each resin, 6 mL of
30 TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by stirring for 90 minutes. Diethyl ether was added to the reaction solution, the resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the resin was removed by filtration. Then, linear
35 density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with

eluant A/B: 74/26-64/36 using: eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the product were collected and lyophilized. The obtained white powders, 117.7 mg, were dissolved in 200 mL of water and 394 μ L of ion exchange resin AG1 x 8 AcO⁻ form, which was obtained by converting commercially available BioRAD AG1 x 8 Cl⁻ form into the acetate type in a conventional manner, was added to the solution. While manually stirring the reaction solution sometimes, the solution was allowed to stand for an hour. The solution was then filtered through a membrane filter to remove the resin and give 115.5 mg of white powders as the acetate. To the purified sample (the acetate) obtained, 5.775 ml of glacial acetic acid was added and the mixture was sonicated for 5 minutes. Subsequently, 17.325 ml of pure water was added to prepare 5 mg/ml/25% aqueous acetic acid solution. The resulting 5 mg/ml solution was dispensed by 4 ml each in five vials, and the rest of the solution was transferred to a vial. These vials were frozen at -80°C for 2 hours and then lyophilized, in which the lyophilization was performed while cooling at -40°C for 4 hours, -20°C for 2 hours, 0°C for 12 hours, 5°C for 8 hours and 20°C for 5 hours. As a result, five vials of 20 mg and one vial of 11.95 mg prepared from the remaining solution were obtained.

Mass spectrum (M+H)⁺ 1243.6 (Calcd. 1243.6)

Anal. for amino acids (20% HCl containing 4% thioglycolic acid, 110°C, hydrolysis for 24 hours; figures in parentheses show theoretical values.): Asp 1.01 (1); Thr 0.96 (1); Leu 0.97 (1); Tyr 1.02 (1); Phe(4F) 1.00 (1)

Elution time on HPLC: 12.0 mins.

Elution conditions:

Column: YMC ODS AM-301 (4.6 x 100 mm)

Linear density gradient elution with eluants A/B =80/20-30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

Flow rate: 1.0 ml/min.

30

(EXAMPLE 14)

Production of des(1)-Ac-[D-Tyr2,Hyp3,Thr5,AzaGly7,Trp10]MS10 (Compound No. 797)

The Rink amide MBHA resin, 357 mg (0.25 mmol), was used as a starting material, and the starting material were reacted on peptide synthesizer ABI 433A

(according to the Fmoc/DCC/HOBt 0.25 mmol protocol), whereby Trp(Boc), Arg(Pbf) and Leu were introduced to give the H-Leu-Arg(Pbf)-Trp(Boc)-Rink amide MBHA resin. In a separate reactor, 290.75 mg (1 mmol) of the Fmoc-NHNH₂.HCl was weighed and dissolved in DMF. Under ice cooling, a suspension of 156.9 mg (0.95 mmol) of CDI in THF and 339.7 µl of DIEA were added and the mixture was stirred at room temperature for an hour. The mixture was added to the H-Leu-Arg(Pbf)-Trp(Boc)-Rink amide MBHA resin, followed by stirring at room temperature overnight. After the resin was washed, Phe, Thr(Bu^t), Asn(Trt), Hyp(Bu^t) and D-Tyr(Bu^t) were introduced in this order again on peptide synthesizer ABI 433A (according to the Fmoc/DCC/HOBt 0.25 mmol protocol) to give the H-D-Tyr(Bu^t)-Hyp(Bu^t)-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Pbf)-Trp(Boc)-Rink amide MBHA resin. Subsequently, the resin was treated for 20 minutes in DMF with 94.4 µl (1 mmol) of Ac₂O and 174.2 µl (1 mmol) of DIEA for N-terminal acetylation to give 596.6 mg of the Ac-D-Tyr(Bu^t)-Hyp(Bu^t)-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Pbf)-Trp(Boc)-Rink amide MBHA resin. To the resin obtained, 4 mL of TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by stirring for 90 minutes. Diethyl ether was added to the reaction solution, the resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the resin was removed by filtration. Then, linear density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with eluants A/B: 74/26-64/36 using: eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the product were collected and lyophilized. The obtained white powders, 146.3 mg, were dissolved in 100 mL of water and 530 µL of ion exchange resin BioRAD AG1 x 8 AcO⁻ was added to the solution, followed by stirring for an hour. After the resin was removed by cotton plug filtration through silica wool, the same amount of the resin was again added and the mixture was stirred for an hour. The solution was filtered through a membrane filter to remove the resin and lyophilized to give 127.3 mg of white powders as the acetate.

Mass spectrum (M+H)⁺ 1211.1 (Calcd. 1211.6)

Anal. for amino acids (20% HCl containing 4% thioglycolic acid, 110°C, hydrolysis for 24 hours; figures in parentheses show theoretical values.): Asp 0.99 (1); Thr 0.96

(1); Leu 0.93 (1); Tyr 0.98 (1); Phe 1.00 (1); Arg 0.99 (1)

Elution time on HPLC: 11.4 mins.

Elution conditions:

Column YMC ODS AM-301 (4.6 x 100 mm)

5 Linear density gradient elution with eluants A/B = 80/20-30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)
Flow rate: 1.0 ml/min.

(EXAMPLE 15)

10 Production of des(1)-Ac-[D-Tyr²,Hyp³,Alb⁴,Thr⁵,AzaGly⁷,Arg(Me)⁹,Trp¹⁰]MS10
(Compound No. 800)

The Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin (0.378 mmol/g), 661 mg (0.25 mmol), was used as a starting material, and the starting material were reacted on peptide synthesizer ABI 433A (according to the
15 Fmoc/DCC/HOBt 0.25 mmol protocol), whereby Phe, Thr(Bu^t), Alb, Hyp(Bu^t) and D-Tyr(Bu^t) were introduced in this order thereby to give the H-D-Tyr(Bu^t)-Hyp(Bu^t)-Alb-Thr(Bu^t)-Phe-Gly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin. Subsequently, the resin was treated for 20 minutes in DMF with 94.4 µl (1 mmol) of Ac₂O and 174.2 µl (1 mmol) of DIEA for N-terminal
20 acetylation to give 1.173 g of the Ac-D-Tyr(Bu^t)-Hyp(Bu^t)-Asn(Trt)-Thr(Bu^t)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin. To the resin obtained, 7 mL of TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by stirring for 90 minutes. Diethyl ether was added to the reaction solution, the
25 resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the resin was removed by filtration. Thereafter, linear density gradient elution (60 minutes) was performed at a flow rate of 8 ml/min with eluants A/B: 74/26-64/36 using: eluant A: 0.1% TFA in water and eluant B:
30 0.1% TFA-containing acetonitrile on preparative HPLC using Daisopak-SP100-5-ODS-P (20 x 250 mm). The fractions containing the product were collected and lyophilized. The obtained white powders, 94.8 mg, were dissolved in 20 mL of water and 320 µL of ion exchange resin BioRAD AG1 x 8 AcO⁻ form was added to the solution. The solution was filtered through a
35 membrane filter to remove the resin and lyophilized to give 79.4 mg of white

powders as the acetate.

Mass spectrum (M+H)⁺ 1240.4 (Calcd. 1240.6)

Anal. for amino acids (20% HCl containing 4% thioglycolic acid, 110°C, hydrolysis for 24 hours; figures in parentheses show theoretical values.): Thr 1.02 (1); Leu 0.99

5 (1); Tyr 1.00 (1); Phe 1.00 (1)

Elution time on HPLC: 12.3 mins.

Elution conditions:

Column: YMC ODS AM-301 (4.6 x 100 mm)

Linear density gradient elution with eluants A/B =80/20-30/70, using 0.1%

10 TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

Flow rate: 1.0 ml/min.

(EXAMPLE 16)

(Synthesis F): Production of

15 des(1-5)-4-[bis-(2-pyridylmethyl)aminomethyl]benzoyl-[AzaGly7,Arg(Me)9,Trp10]
MS10 (Compound No. 801)

In a reactor, 265 mg (0.1 mmol) of the

Fmoc-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink amide MBHA resin (0.378 mmol/g) was weighed, and washed and swollen in DMF. The resin was then

20 treated in 5 ml of 20% piperidine/DMF solution for 20 minutes for Fmoc

deprotection. The resin was treated with 155.0 mg (0.4 mmol) of Fmoc-Phe-OH, 63.6 µL (0.4 mmol) of DIPCDI and 0.8 mL (0.4 mmol) of 0.5M HOAt/DMF at room temperature for 90 minutes to introduce Phe. Subsequently, Fmoc deprotection

through treatment in 5 ml of 20% piperidine/DMF solution for 20 minutes and

25 condensation by the DIPCDI/HOAt method as in the Phe introduction, whereby

Ambz(4) was introduced to give the

Fmoc-Ambz(4)-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(Boc)-Rink Amide MBHA

resin. After Fmoc deprotection, the resulting resin was treated for 10 minutes in

DMF with 11.4 µl (0.12 mmol) of 2-pyridinecarboxyaldehyde in the presence of 50

30 µl of AcOH, 41.1 mg (0.4 mmol) of NaBH₃CN was added to thereto and the mixture was stirred for an hour. After washing in DMF, the same procedure was repeated again. The resin obtained was washed in DMF and MeOH and dried to give 281.4 mg of the

4-[Bis-(2-Pyridylmethyl)aminomethyl]benzoyl-Phe-AzaGly-Leu-Arg(Boc₂,Me)-Trp(

35 Boc)-Rink Amide MBHA resin.

To the resin obtained, 2 mL of TFA/PhSMe/m-cresol/H₂O/TIS/EDT (80/5/5/5/2.5/2.5) was added, followed by stirring for 90 minutes. Diethyl ether was added to the reaction solution, the resulting precipitate was centrifuged, and the supernatant was removed. This procedure was repeated twice for washing. The residue was extracted with an aqueous acetic acid solution and the resin was removed by filtration. Then, linear density gradient elution (60 minutes) was performed at a flow rate of 15 ml/min with eluants A/B: 76/24-66/34 using: eluant A: 0.1% TFA in water and eluant B: 0.1% TFA-containing acetonitrile on preparative HPLC using YMC Pack R&D-ODS-5-B S-5, 120A column (30 x 250 mm). The fractions containing the product were collected and lyophilized. The obtained white powders, 17.3 mg, were dissolved in 20 mL of AcCN-water (9:1) and 210 µL of ion exchange resin BioRAD AG1 x 8 AcO⁻ was added to the solution. While manually stirring the reaction solution sometimes, the solution was filtered through a membrane filter to remove the resin and lyophilized to give 10.8 mg of white powders as the acetate.

Mass spectrum (M+H)⁺ 1007.3 (Calcd. 1007.5)

Elution time on HPLC: 10.7 mins.

Elution conditions:

Column: YMC ODS AM-301 (4.6 x 100 mm)

Linear density gradient elution with eluants A/B = 80/20-30/70, using 0.1% TFA in water as eluant A and acetonitrile containing 0.1% TFA as eluant B (25 mins.)

Flow rate: 1.0 ml/min.

The structures of compounds synthesized as in EXAMPLES 1 to 16 or compounds synthesized by synthesis methods similar to EXAMPLES 1 to 6 and physicochemical properties of these compounds are shown in TABLE 1A below.

The description of the column "Synthesis method" in the Table represents that the compound may be synthesized or was synthesized by or according to the synthesis method described therein.

More specifically, "A" in the column "Synthesis method" of Compound number 708 represents that the compound was synthesized by the synthesis method A as described in EXAMPLE 1.

"B" in the column "Synthesis method" of Compound number 709 represents that the compound was synthesized by the synthesis method B as described in

EXAMPLE 3.

“C” in the column “Synthesis method” of Compound number 713 represents that the compound was synthesized by the synthesis method C as described in EXAMPLE 4.

5 “D” in the column “Synthesis method” of Compound number 714 represents that the compound was synthesized by the synthesis method D as described in EXAMPLE 5.

“E” in the column “Synthesis method” of Compound number 717 represents that the compound was synthesized by the synthesis method E as described in EXAMPLE 6.

10 Each description in the column “Synthesis method” of each compound described in EXAMPLES 7 to 16 represents that the compound may be synthesized according to each synthesis method described therein.

Each description in the column “Synthesis method” of each compound which are not described in EXAMPLES 1 to 16 represents that the compound was
15 synthesized according to each synthesis method described therein.

The description of the column “HPLC condition” in the Table represents that each compound was eluted by each HPLC condition of “a”, “b” or “c” described therein.

[TABLE 1A]

Compound number	Structure	M+H ⁺ (obs.)	M+H ⁺ (cal.)	HPLC (min.)	HPLC condition	Synthesis method
708	des(1)-Ac-[D-Tyr2,D-Trp3,Thr5,AzaGly7,D-Arg9,Trp10]MS10	1284.9	1284.6	13.3	a	A
709	des(1)-3-Ac-[Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	949.8	949.5	10.2	a	B
710	des(1)-3-Decanoyl-[Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1061.9	1061.6	19.4	a	B
712	des(1)-2-Ac-[AcP3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1020.7	1020.6	9.2	a	C
713	des(1)-2-Ac-[AcP3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1062.7	1062.6	10.7	a	C
714	des(1)-Ac-[D-Tyr2,D-Trp3,Asp(NHCH2)4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1368.9	1368.7	20.9	b	D
715	des(1)-Ac-[D-Tyr2,D-Trp3,Asp(NHCH2)4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1339.1	1338.7	18.4	b	D
716	des(1)-Ac-[D-Tyr2,D-Trp3,Asp(NHCH2)4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1389.0	1388.7	20.2	b	D
717	des(1)-Ac-[D-Tyr2,D-Trp3,Alb4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1313.9	1313.7	18.3	b	E
718	des(1)-Ac-[D-Tyr2,D-Pya(4)3,Alb4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1275.9	1275.6	14.5	b	E
719	des(1)-Ac-[D-Tyr2,D-Trp3,D-Pro5,AzaGly7,Arg(Me)9,Trp10]MS10	1294.8	1294.6	18.4	b	E
720	des(1)-Ac-[D-Tyr2,Aze(2)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1195.8	1195.6	11.7	a	E
721	des(1)-Ac-[D-Tyr2,Pic(2)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1223.8	1223.6	12.5	a	E
722	des(1)-Ac-[D-Tyr2,Pic(3)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1225.9	1225.6	11.4	a	E
723	des(1)-Ac-[D-Tyr2,Hyp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1227.8	1227.6	12.6	a	E
724	des(1)-Ac-[D-Tyr2,Thz3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1197.8	1197.6	12.0	a	E
725	des(1)-Ac-[D-Tyr2,NMeAla3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1169.7	1169.6	11.3	a	E
726	des(1)-Ac-[D-Tyr2,Gly3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1197.7	1197.6	13.0	a	E
727	des(1)-Ac-[D-Tyr2,Alb3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1231.6	1231.7	13.1	a	E
728	des(1)-Ac-[D-Tyr2,Abz(2)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1195.8	1195.6	11.0	a	E
730	des(1)-Ac-[D-Tyr2,Aze(3)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1183.8	1183.6	11.5	a	E
731	des(1)-Ac-[D-Tyr2,Sar3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1197.8	1197.6	12.9	a	E
732	des(1)-Ac-[D-Tyr2,D-NMeAla3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1210.7	1210.6	10.5	a	E
734	des(1)-Ac-[D-Tyr2,Izc3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1227.9	1227.6	11.4	a	E
735	des(1)-Ac-[D-Tyr2,D-Asp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1198.9	1198.6	10.1	a	E
736	des(1)-Ac-[D-Tyr2,D-Dap3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1199.8	1199.6	11.0	a	E
737	des(1)-Ac-[D-Tyr2,D-Ser3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1249.4	1249.4	10.9	a	E
738	des(1)-Ac-[D-Tyr2,D-Gln3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1284.5	1284.7	16.2	b	E
739	des(1)-Ac-[D-Tyr2,D-His3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1183.8	1183.6	11.8	a	E
740	des(1)-Ac-[D-Tyr2,D-Trp3,Dab4,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1265.0	1265.6	13.6	a	E
742	des(1)-Ac-[D-Tyr2,Ala3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1199.8	1199.3	11.3	a	E
743	des(1)-Ac-[D-Tyr2,Leu3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1241.0	1240.7	10.3	a	E
744	des(1)-Ac-[D-Tyr2,Ser3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10					
745	des(1)-Ac-[D-Tyr2,Lys3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10					





746	des(1)-Ac-[D-Tyr2,Glu3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1241.8	1241.4	11.4	a	E
747	des(1)-Ac-[D-Tyr2,β-Ala3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1183.8	1183.3	11.0	a	E
748	des(1)-Ac-[D-Tyr2,D-Trp3,Thr5,Phe(4CI)6,AzaGly7,Arg(Me)9,Trp10]MS10	1333.0	1332.6	15.2	a	E
749	des(1)-Ac-[D-Tyr2,D-Trp3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1316.9	1316.6	14.2	a	E
750	des(1)-Ac-[D-Tyr2,D-Trp3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1316.9	1316.6	14.4	a	E
754	des(1)-Ac-[D-Tyr2,Lys3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1258.8	1258.6	10.6	a	E
755	des(1)-Ac-[D-Tyr2,Glu3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1259.8	1259.6	11.7	a	E
756	des(1)-Ac-[D-Tyr2,Lys3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1258.8	1258.6	10.8	a	E
757	des(1)-Ac-[D-Tyr2,Glu3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1259.9	1259.6	11.9	a	E
758	des(1)-Ac-[D-Tyr2,Lys3,Thr5,Phe(4CI)6,AzaGly7,Arg(Me)9,Trp10]MS10	1274.8	1274.6	11.6	a	E
759	des(1)-Ac-[D-Tyr2,Glu3,Thr5,Phe(4CI)6,AzaGly7,Arg(Me)9,Trp10]MS10	1275.8	1275.6	12.8	a	E
760	des(1)-Ac-[D-Tyr2,Pzc(2)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1224.8	1224.6	10.2	a	E
763	des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1243.8	1243.6	11.5	a	E
764	des(1)-Ac-[D-Tyr2,Trp3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1316.9	1316.6	14.6	a	E
765	des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1243.6	1243.6	12.0	a	E
766	des(1)-Ac-[D-Tyr2,Trp3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1316.4	1316.6	14.8	a	E
767	des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(4CI)6,AzaGly7,Arg(Me)9,Trp10]MS10	1259.7	1259.6	12.9	a	E
768	des(1)-Ac-[D-Tyr2,Trp3,Thr5,Phe(4CI)6,AzaGly7,Arg(Me)9,Trp10]MS10	1332.4	1332.6	15.6	a	E
769	des(1)-Ac-[D-Tyr2,Gly3,Thr5,Phe(4CI)6,AzaGly7,Arg(Me)9,Trp10]MS10	1203.5	1203.5	13.0	a	E
770	des(1)-Ac-[D-Tyr2,Alb3,Thr5,Phe(4CI)6,AzaGly7,Arg(Me)9,Trp10]MS10	1261.4	1231.6	14.7	a	E
771	des(1)-Ac-[D-Tyr2,Orn3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1226.3	1226.4	14.2	b	E
772	des(1)-Ac-[D-Tyr2,Trp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1213.4	1213.6	15.6	b	E
773	des(1)-Ac-[D-Tyr2,His(3Me)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1263.3	1263.6	14.4	b	E
774	des(1)-Ac-[D-Tyr2,DL-Ala(Pip)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1266.3	1266.7	14.5	b	E
775	des(1)-Ac-[D-Tyr2,Tyr(PO ₃ H ₂)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1254.8	1254.6	15.7	b	E
776	des(1)-Glycolyl-[D-Tyr2,Hyp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1241.3	1241.6	15.3	b	E
777	des(1-2)-Ac-[D-Tyr3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1112.6	1112.3	15.4	b	E
780	des(1)-Ac-[D-Tyr2,Pro(4NH ₂)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1224.7	1224.6	10.3	a	E
781	des(1)-Ac-[D-Tyr2,Hyp(Bzl)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1315.8	1315.7	15.7	a	E
782	des(1)-Ac-[D-Tyr2,D-NMePhe3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1273.8	1273.6	15.4	a	E
783	des(1)-Ac-[D-Tyr2,Gly3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1187.8	1187.6	11.7	a	E
784	des(1)-Ac-[D-Tyr2,Alb3,Thr5,Phe(2F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1215.8	1215.6	13.4	a	E
785	des(1)-Ac-[D-Tyr2,Gly3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1187.7	1187.6	11.9	a	E
786	des(1)-Ac-[D-Tyr2,Alb3,Thr5,Phe(3F)6,AzaGly7,Arg(Me)9,Trp10]MS10	1215.8	1215.6	13.5	a	E
787	des(1)-Ac-[D-Tyr2,Hyp3,Thr5,Phe(4F)8,AzaGly7,Arg(Me)9,Trp10]MS10	1243.6	1243.6	12.0	a	E






788	des(1)-Ac-[D-Tyr2, Glu3, Thr5, Phe(4F)6, AzaGly7, Arg(Me)9, Trp10]MS10	1259.5	1259.6	11.9	a	E
789	des(1)-Ac-[D-Tyr2, Lys3, Thr5, Phe(4F)6, AzaGly7, Arg(Me)9, Trp10]MS10	1258.6	1258.6	10.8	a	E
790	des(1)-Ac-[D-Tyr2, Gly3, Thr5, Phe(4F)6, AzaGly7, Arg(Me)9, Trp10]MS10	1187.4	1187.6	11.9	a	E
791	des(1)-Ac-[D-Tyr2, Ala3, Thr5, Phe(4F)6, AzaGly7, Arg(Me)9, Trp10]MS10	1215.2	1215.6	13.6	a	E
794	des(1)-Ac-[D-Tyr2, Hyp3, Thr5, D-Phe6, AzaGly7, Arg(Me)9, Trp10]MS10	1225.5	1225.6	11.8	a	E
797	des(1)-Ac-[D-Tyr2, Hyp3, Thr5, AzaGly7, Trp10]MS10	1211.1	1211.6	11.4	a	A
800	des(1)-Ac-[D-Tyr2, Hyp3, Ala4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1240.4	1240.6	12.3	a	E
801	des(1)-5-4-[Bis-(2-Pyridylmethyl)aminomethyl]benzoyl-[AzaGly7, Arg(Me)9, Trp10]MS10	1007.3	1007.5	10.7	a	F
809	des(1)-Ac-[D-Tyr2, Hyp3, NMeSer5, AzaGly7, Arg(Me)9, Trp10]MS10	1225.6	1225.6	10.3	c	E
810	des(1)-Ac-[D-Tyr2, Hyp3, Hyp5, AzaGly7, Arg(Me)9, Trp10]MS10	1237.6	1237.6	10.5	c	E
813	des(1)-Ac-[D-Tyr2, Hyp3, Gly5, AzaGly7, Arg(Me)9, Trp10]MS10	1181.5	1181.6	10.1	c	E
814	des(1)-Ac-[D-Tyr2, Hyp3, Ala5, AzaGly7, Arg(Me)9, Trp10]MS10	1195.4	1195.6	11.0	c	E
815	des(1)-Ac-[D-Tyr2, Hyp3, D-Ala5, AzaGly7, Arg(Me)9, Trp10]MS10	1195.6	1195.6	10.8	c	E
816	des(1)-Ac-[D-Tyr2, Hyp3, His4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1248.6	1248.6	9.7	c	E
843	des(1)-Ac-[D-Tyr2, Hyp3, Gln4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1239.8	1239.6	10.4	c	E
844	des(1)-Ac-[D-Tyr2, Hyp3, D-Asn4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1235.6	1225.6	10.2	c	E
845	des(1)-Ac-[D-Tyr2, Hyp3, Cit4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1269.0	1268.7	10.3	c	E
846	des(1)-Ac-[D-Tyr2, Hyp3, D-Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1226.1	1225.6	10.3	c	E
856	des(1)-Ac-[D-Tyr2, Hyp3, Thr5, AzaGly7, Ala(cP)8, Arg(Me)9, Trp10]MS10	1223.9	1223.6	10.9	a	E
860	des(1)-5-4-Ureidobenzoyl-[AzaGly7, Arg(Me)9, Trp10]MS10	854.7	854.4	17.7	b	F
861	des(1)-Ac-[D-Tyr2, Hyp3, Arg4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1268.1	1267.7	10.5	c	E
862	des(1)-Ac-[D-Tyr2, Hyp3, Gly4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1168.9	1168.6	11.2	c	E
863	des(1)-Ac-[D-Tyr2, Hyp3, Dap4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1198.2	1197.6	9.9	c	E
864	des(1)-Ac-[D-Tyr2, Hyp3, Dab4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1212.3	1211.6	9.9	c	E
868	des(1)-Ac-[D-Tyr2, Hyp3, Thr5, α MePhe6, AzaGly7, Arg(Me)9, Trp10]MS10	1240.0	1239.6	13.4	a	E
870	des(1)-Ac-[D-Tyr2, Hyp3, Thr5, Phe(2Me)6, AzaGly7, Arg(Me)9, Trp10]MS10	1239.7	1239.6	12.5	a	E
872	des(1)-Ac-[D-Tyr2, Hyp3, Thr5, Phe(3Me)6, AzaGly7, Arg(Me)9, Trp10]MS10	1239.6	1239.6	12.7	a	E
874	des(1)-Ac-[D-Tyr2, Hyp3, Thr5, Phe(4Me)6, AzaGly7, Arg(Me)9, Trp10]MS10	1239.6	1239.6	12.5	a	E
877	des(1)-Ac-[D-Tyr2, Hyp3, Thr5, threo-Ser(3Phenyl)6, AzaGly7, Arg(Me)9, Trp10]MS10	1241.1	1241.6	10.5	a	E
882	des(1)-Ac-[D-Tyr2, Hyp3, Thr5, erythro-Ser(3Phenyl)6, AzaGly7, Arg(Me)9, Trp10]MS10	1241.6	1241.6	10.9	a	E
886	des(1)-Ac-[D-Tyr2, Hyp3, Nva4, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1210.8	1210.6	14.0	a	E
887	des(1)-2-Ac-[Hyp3, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1062.7	1062.5	10.2	a	B
888	des(1)-2-3-(p-Hydroxyphenyl)propionyl-[Hyp3, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1168.7	1168.6	12.2	a	B
889	des(1)-2-[p-Glu3, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1018.5	1018.5	10.3	a	B
896	des(1)-Ac-[D-Tyr2, cisHyp3, Thr5, AzaGly7, Arg(Me)9, Trp10]MS10	1225.6	1225.4	12.4	a	E






897	des(1)-Ac-[D-Tyr2,Pro(4F)3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1227.4	1227.6	12.9	a	E
899	des(1)-Ac-[Tyr2,Hyp3,Thr5,AzaGly7,Arg(Me)9,Trp10]MS10	1225.7	1225.4	17.0	b	E
	a:20-70% AcCN/25min, flow 1ml/min.					
	YMC ODS AM-301 (4.6 x 100mm)					
	b:0-50% AcCN/25min, flow 1ml/min.					
	Wakosil-II 5C18 HG (4.6 x 100mm)					
	c:20-70% AcCN/25min flow 1ml/min					
	SHISEIDO CAPCELL PAK C18 MGII (4.6 x 100mm)					






The structures of compounds in TABLE 1A above are shown in TABLE 1B.






[TABLE 1B]






Compound No.	Structure
708	
709	
710	
712	






713	
714	
715	
716	
717	






718	
719	
720	
721	
722	






723	
724	
725	
726	
727	






728	
730	
731	
732	
734	






735	
736	
737	
738	
739	






740	
742	
743	
744	
745	

746	
747	
748	
749	
750	


754	
755	
756	
757	
758	

759	
760	
763	
764	
765	


766	
767	
768	
769	
770	

771	
772	
773	
774	
775	


776




777




780













781













782













783	
784	
785	
786	
787	






788	
789	
790	
791	
794	






797	
800	
801	
809	
810	



813	
814	
815	
816	
843	

844	
845	
846	
856	
860	

861	
862	
863	
864	
868	

870	
872	
874	
877	
882	

886	
887	
888	
889	
896	

897	
899	

TEST EXAMPLE 1

Assay for Intracellular Ca Ion Level-Increasing Activity Using FLIPR

In accordance with the method described in JPA 2000-312590, the intracellular Ca ion level-increasing activity was measured using FLIPR.

5 The stable expression cell line hOT7T175 was obtained by transfection of the expression plasmid pAK-rOT175 for animal cells into CHO/dhfr⁻ cells, using CellPfect Transfection Kit (Amersham Pharmacia Biotech, Inc.). First, 240 μ L of Buffer A (attached to CellPfect Transfection Kit) was added to 9.6 μ g of plasmid DNA dissolved in 240 μ L of distilled water followed by stirring. After the mixture
10 was settled for 10 minutes, 480 μ L of Buffer B (attached to CellPfect Transfection Kit) was added to the mixture, which was vigorously stirred to form liposomes containing the DNA. Then, 4×10^5 CHO/dhfr⁻ cells (obtained from ATCC) were inoculated on a 60 mm Petri dish. After culturing the cells in Ham's F-12 medium (Nissui Seiyaku Co., Ltd.) supplemented with 10% fetal bovine serum (BIO
15 WHITTAKER, Inc.) at 37°C for 2 days in 5% carbon dioxide gas, 480 mL of the liposomes were dropwise added to the cells on the Petri dish. After culturing the cells at 37°C for 6 hours in 5% carbon dioxide gas, the cells were washed twice with serum-free Ham's F-12 medium and 3 mL of 15% glycerol was added to the cells on the Petri dish followed by treatment for 2 minutes. The cells were again washed
20 twice with serum-free Ham's F-12 medium followed by incubation in Ham's F-12 medium supplemented with 10% fetal bovine serum at 37°C for 15 hours in 5% carbon dioxide gas. The cells were dispersed by trypsin treatment to recover from the Petri dish. The recovered cells were inoculated on a 6-well plate in 1.25×10^4 cells/well and incubation was initiated at 37°C in Dulbecco's modified Eagle medium
25 (DMEM) medium (Nissui Seiyaku Co., Ltd.) containing 10% dialyzed fetal bovine serum (JRH BIOSCIENCES, Inc.) in 5% carbon dioxide gas. The plasmid-transfected CHO transformants grew in the medium but the non-transfected cells gradually died, accordingly the medium was exchanged on Days 1 and 2 after the initiation of incubation to remove the cells died. Approximately 20 colonies of
30 the CHO transformants that kept growing on Days 8 to 10 after the incubation were isolated. From the cells in these colonies, cells showing high reactivity with the ligand peptide metastin (hereinafter merely referred to as hOT7T175/CHO) were selected to provide for the following experiment.

35 The intracellular Ca ion level-increasing activity of the synthetic peptide in hOT7T175/CHO was determined using FLIPR (Molecular Devices, Inc.).

hOT7T175/CHO was subcultured in DMEM supplemented with 10% dialyzed fetal bovine serum (hereinafter abbreviated as dFBS) (hereinafter referred to as 10% dFBS/DMEM) and provided for the experiment. The hOT7T175/CHO was suspended in 10% dFBS-DMEM in 15×10^4 cells/ml. The suspension was
5 inoculated on a 96-well plate for FLIPR (Black Plate Clear Bottom, Coster, Inc.) at 200 μ l/well (3.0×10^4 cells/200 μ L/well), followed by incubation at 37°C overnight in a 5% CO₂ incubator (hereinafter referred to as the cell plate). Then, 21 ml of HANKS/HBSS (9.8 g of HANKS, 0.35 g of sodium hydrogencarbonate, 20 ml of 1M HEPES; after adjusting the pH to 7.4 with 1N sodium hydroxide, the mixture
10 was sterilized by filtration), 210 μ l of 250 mM Probenecid and 210 μ l of fetal bovine serum (FBS) (HANKS/HBSS-Probenecid-FBS) were mixed.

Furthermore, 2 vials of Fluo3-AM (50 μ g/vial) were dissolved in 21 μ L of dimethylsulfoxide and 21 μ L of 20% Pluronic acid. The resulting solution was added to and mixed with 10 ml of the HANKS/HBSS-Probenecid-FBS described
15 above. After the culture medium was removed, the mixture was dispensed onto the cell plate in 100 μ l each/well, followed by incubation at 37°C for an hour in a 5% CO₂ incubator (pigment loading). The peptide was dissolved in dimethylsulfoxide in 1×10^{-3} M. The peptide solution was diluted with HANKS/HBSS containing 2.5 mM Probenecid, 0.2% BSA and 0.1% CHAPS. The dilution was then transferred to
20 a 96-well plate for FLIPR (V-Bottom plate, Coster, Inc.) (hereinafter referred to as the sample plate). After completion of the pigment loading onto the cell plate, the cell plate was washed 4 times with wash buffer, which was prepared by adding 2.5 mM Probenecid to HANKS/HBSS, using a plate washer to leave 100 μ L of the wash buffer after the washing. The cell plate and the sample plate were set in FLIPR and
25 0.05 ml of a sample from the sample plate was automatically transferred to the cell plate with the FLIPR device to promote the cell response. A change in the intracellular calcium ion level for 180 seconds was measured with passage of time. The intracellular Ca ion level-increasing activity [specific activity to Metastin (1-54)] is shown in TABLE 2.

[TABLE 2]

Compound No.	Specific Activity	Compound No.	Specific Activity	Compound No.	Specific Activity
708	12.8	747	2.6	789	3.4
709	9.8	748	0.4	790	2.3
710	7.2	749	0.5	791	2.9
712	8.9	750	1.0	794	2.3
713	3.7	754	3.1	797	1.4
714	0.7	755	3.5	800	0.8
715	1.6	756	2.4	801	11.1
716	3.6	757	2.7	809	1.0
717	1.0	758	1.8	810	1.5
718	2.6	759	1.9	813	0.5
719	3.8	760	1.7	814	0.5
720	0.8	763	6.4	815	0.5
721	1.1	764	1.5	816	1.1
722	2.2	765	2.6	843	0.8
723	2.0	766	0.2	844	0.9
724	1.4	767	1.5	845	1.0
725	2.7	768	0.6	846	0.9
726	3.1	769	1.5	856	0.6
727	1.5	770	2.6	860	1.3
728	3.2	771	1.9	861	1.4
730	4.1	772	1.3	862	0.7
731	8.8	773	1.1	863	0.9
732	3.2	774	2.0	864	1.0
734	2.3	775	1.3	868	1.4
735	3.4	776	1.5	870	0.6
736	2.4	777	1.8	872	0.7
737	3.0	780	2.1	874	0.8
738	1.4	781	1.0	877	1.0
739	1.6	782	1.8	882	0.8
740	1.1	783	8.8	886	1.2

742	2.3	784	3.4	888	1.1
743	3.4	785	3.2	896	0.6
744	2.4	786	3.3	897	0.7
745	3.0	787	2.3	899	1.1
746	2.2	788	2.7		

TEST EXAMPLE 2

Evaluation of Blood Testosterone Level Reducing Effect of Metastin Peptide Derivatives Using Mature Male Rats

- 5 A metastin peptide derivative (hereinafter referred to as peptide) was dissolved in 50% DMSO aqueous solution (Otsuka Pharmaceutical Co., Ltd.) to prepare a peptide solution with a concentration of 0.03, 0.01 or 0.003 mM. This peptide solution was filled in five ALZET osmotic pumps (Model 2001, 0.2 ml in volume, release rate: 0.001 ml/hr, DURECT Corporation). The ALZET pumps
- 10 filled with the peptide solution were implanted subcutaneously in 5 CD(SD)IGS male rats of 9 weeks old after birth (Charles River Japan, Inc.) on the back under ether anesthesia by one pump/animal. For negative control, 50% DMSO aqueous solution was filled in 5 ALZET osmotic pumps, which were similarly implanted in 5 male CD(SD)IGS rats (Charles River Japan, Inc.), respectively. These
- 15 pump-implanted rats were fed for 6 days under normal feeding conditions. After weighing, the animal was decapitated to collect blood. After 0.03 ml/ml blood of aprotinin solution (Trasylol, Bayer) containing 0.1 g/ml EDTA.2Na was added to blood, the plasma was separated and recovered by centrifugation at 1,800 x g for 25 min. From the plasma obtained, 0.05 ml was applied to radioimmunoassay
- 20 (DPC Total Testosterone Kit, Diagnostic Products Corporation) to measure the plasma testosterone level in each rat. The peptides are listed in TABLE 3, when the number of rats showing the testosterone level below the measurement limit (0.04 ng/ml of plasma level) in radioimmunoassay was 3 or more in the 5 rats receiving the peptides.

[TABLE 3]


Compound No.	Compound No.
720	763
721	765
723	767
726	773
727	775
732	776
734	787
738	788
739	797
742	814
744	856
745	870
746	872
750	874
755	877
756	882
757	888
758	896
759	897

INDUSTRIAL APPLICABILITY

According to the present invention, there are provided stable metastin derivatives having excellent biological activities (a cancer metastasis suppressing activity, a cancer growth suppressing activity, a gonadotrophic hormone secretion stimulating activity, sex hormone secretion stimulating activity, etc.).

CLAIMS

I. A metastatin derivative represented by formula:



(wherein:

XX0 represents formyl, a C₁₋₂₀ alkanoyl, cyclopropanecarbonyl, 6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl, 6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl, 3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl, acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl, 6-aminocaproyl, 6-acetylaminocaproyl, 4-[bis-(2-pyridylmethyl)aminomethyl]benzoyl or 4-ureidobenzoyl;

XX2 represents Tyr, D-Tyr, D-Ala, D-Leu, D-Phe, D-Lys, D-Trp or a chemical bond;

XX3 represents:

i) an amino acid selected from Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Ser, Thr, Trp, Tyr and Val wherein the α -amino group may optionally be methylated;

ii) a cyclic amino acid selected from Pro, Aze(2), Aze(3), Pic(2), Pic(3), Hyp, Thz, Abz(2), Abz(3), Pzc(2), Pro(4NH₂), Hyp(Bzl), cisHyp, Pro(4F) and lzc;

iii) an amino acid selected from D-Dap, D-Pya(4), DL-Ala(Pip), Orn, Aib and Tyr(PO₃H₂); or,

iv) a chemical bond;

XX4 represents Asn, 2-amino-3-ureidopropionic acid, N ^{β} -formyl- β -diaminopropionic acid, N ^{β} -acetyl- β -diaminopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine, 2,4-diaminobutanoic acid, 2,3-diaminopropionic acid, His, Gln, Gly, Arg, Cit, Nva, D-Asn or a chemical bond;

XX5 represents Ser, Thr, Val, NMeSer, Gly, Ala, Hyp, D-Ala, D-Thr, D-Pro or a chemical bond;

XX6 represents Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), threo-Ser(3Phenyl), erythro-Ser(3Phenyl), or optionally substituted phenylalanine;


AzaGly represents azaglycine;

XX8 represents Leu, Nva, Val or Ala(cPr);

XX9 represents optionally substituted arginine, optionally substituted lysine or optionally substituted ornithine; and,

XX10 represents 2-naphthylalanine, 2-thienylalanine, tyrosine, optionally substituted phenylalanine or optionally substituted tryptophan); or a salt thereof.

2. A metastatin derivative represented by formula:



(wherein:

XX0 represents formyl, a C₁₋₂₀ alkanoyl, cyclopropanecarbonyl, 6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl, 6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl, 3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl, acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl, 6-aminocaproyl, 6-acetylaminocaproyl, 4-[bis-(2-pyridylmethyl)aminomethyl]benzoyl or 4-ureidobenzoyl;

XX2 represents Tyr, D-Tyr, D-Ala, D-Leu, D-Phe, D-Lys, D-Trp or a chemical bond;

XX3 represents D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-Trp, D-Tyr, D-Pya(4), D-NMeAla, D-NMePhe, Aze(2), Aze(3), Pic(2), Pic(3), Hyp, Thz, Gly, Aib, Abz(2), Abz(3), Sar, Izc, Leu, Lys, Glu, Thr, Trp, Ser, Ala, NMeAla, β -alanine, DL-Ala(Pip), Pzc(2), Orn, His(3Me), Tyr(PO₃H₂), Pro(4NH₂), Hyp(Bzl), cisHyp, Pro(4F) or a chemical bond;

XX4 represents Asn, 2-amino-3-ureidopropionic acid, N ^{β} -formyl- β -diaminopropionic acid, N ^{β} -acetyl- β -diaminopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine, 2,4-diaminobutanoic acid, 2,3-diaminopropionic acid, His, Gln, Gly, Arg, Cit, Nva, D-Asn or a chemical bond;

XX5 represents Ser, Thr, Val, NMeSer, Gly, Ala, Hyp, D-Ala, D-Thr, D-Pro or a chemical bond;

XX6 represents Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), Phe(2F), Phe(3F), Phe(4F), Phe(4Cl), α MePhe, Phe(2Me), Phe(3Me), Phe(4Me), threo-Ser(3Phenyl), erythro-Ser(3Phenyl) or D-Phe;


AzaGly represents azaglycine;

XX8 represents Leu, Nva, Val or Ala(cPr);

XX9 represents Arg, Orn, Arg(Me), D-Arg or Arg(asymMe₂); and,

XX10 represents Phe, Trp, 2-naphthylalanine, 2-thienylalanine, tyrosine or 4-fluorophenylalanine); or a salt thereof.

3. A metastin derivative represented by formula:



(wherein:

XX0 represents formyl, a C₁₋₂₀ alkanoyl, cyclopropanecarbonyl, 6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl, 6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl, 3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl, acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl or 6-aminocaproyl;

XX2 represents Tyr, D-Tyr, D-Ala, D-Leu, D-Phe, D-Lys, D-Trp or a chemical bond;

XX3 represents D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-NMeAla, D-NMePhe, Aze(2), Pic(2), Pic(3), Hyp, Thz, NMeAla, Gly, Aib, Abz(2), Abz(3), Sar, Leu, Lys, Glu, β -alanine, Pzc(2), Orn, His(3Me), Tyr(PO₃H₂), Pro(4NH₂) or Hyp(Bzl);

XX4 represents Asn, 2-amino-3-ureidopropionic acid, N ^{β} -formyl- β -diaminopropionic acid, N ^{β} -acetyl- β -diaminopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine, 2,4-diaminobutanoic acid, His, Gln, Cit or D-Asn;

XX5 represents Ser, Thr, Val, NMeSer, Gly, Ala, Hyp, D-Ala or D-Thr;

XX6 represents Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), Phe(2F), Phe(3F), Phe(4F), Phe(4Cl) or D-Phe;

AzaGly represents azaglycine;

XX8 represents Leu, Nva or Val;

XX9 represents Arg, Orn, Arg(Me) or Arg(asymMe₂); and,

XX10 represents Phe, Trp, 2-naphthylalanine, 2-thienylalanine, tyrosine or 4-fluorophenylalanine); or a salt thereof.

4. A metastin derivative represented by formula:

XX0-XX2-XX3-XX4-XX5-XX6-AzaGly-XX8-XX9-XX10-NH₂

(wherein:

XX0 represents formyl, a C₁₋₁₂ alkanoyl, cyclopropanecarbonyl, 6-(acetyl-D-arginylamino)caproyl, 6-((R)-2,3-diaminopropionylamino)caproyl, 6-(D-norleucylamino)caproyl, 4-(D-arginylamino)butyryl, 3-(4-hydroxyphenyl)propionyl, glycyl, tyrosyl, acetylglycyl, acetyltyrosyl, D-tyrosyl, acetyl-D-tyrosyl, pyroglutamyl, 3-(pyridin-3-yl)propionyl, adipoyl, glycoloyl or 6-aminocaproyl;

XX2 represents Tyr, D-Tyr, D-Ala, D-Leu, D-Phe, D-Lys, D-Trp or a chemical bond;

XX3 represents D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-NMeAla, D-NMePhe, Aze(2), Pic(2), Pic(3), Hyp, Thz, NMeAla, Gly, Aib, Abz(2), Abz(3), Sar, Leu, Lys, Glu, β -alanine, Pzc(2), Orn, His(3Me), Tyr(PO₃H₂), Pro(4NH₂) or Hyp(Bzl);

XX4 represents Asn, 2-amino-3-ureidopropionic acid, N ^{β} -formyldiaminopropionic acid, N ^{β} -acetyldiaminopropionic acid, N ^{ω} -pentylasparagine, N ^{ω} -cyclopropylasparagine, N ^{ω} -benzylasparagine or 2,4-diaminobutanoic acid;

XX5 represents Ser, Thr or Val;

XX6 represents Phe, Tyr, Trp, Tyr(Me), Thi, Nal(2), Cha, Pya(4), Phe(2F), Phe(3F), Phe(4F) or Phe(4Cl);

AzaGly represents azaglycine;

XX8 represents Leu, Nva or Val;

XX9 represents Arg, Orn, Arg(Me) or Arg(asymMe₂); and,

XX10 represents Phe, Trp, 2-naphthylalanine, 2-thienylalanine, tyrosine or 4-fluorophenylalanine); or a salt thereof.

5. The metastin derivative or a salt thereof according to claim 1, wherein:

XX0 represents a C₁₋₁₂ alkanoyl, 6-aminocaproyl, 6-acetylaminocaproyl, glycoloyl, 4-[bis-(2-pyridylmethyl)aminomethyl]benzoyl, 4-ureidobenzoyl, 3-(4-hydroxyphenyl)propionyl or pyroglutamyl;

XX2 represents D-Tyr, Tyr or a chemical bond;

XX3 represents D-Asp, D-Dap, D-Ser, D-Gln, D-His, D-Trp, D-Tyr, D-Pya(4) D-NMeAla, D-NMePhe, Aze(2), Aze(3), Pic(2), Pic(3), Hyp, Thz, Gly, Aib, Abz(2), Sar, Izc, Leu, Lys, Glu, Thr, Trp, Ser, Ala, NMeAla, β -alanine,

DL-Ala(Pip), Pzc(2), Orn, His(3Me), Tyr(PO_3H_2), Pro(4NH_2), Hyp(Bzl), cisHyp, Pro(4F) or a chemical bond;

XX4 represents Asn, 2-amino-3-ureidopropionic acid, N^ω -pentylasparagine, N^ω -cyclopropylasparagine, N^ω -benzylasparagine, 2,4-diaminobutanoic acid, 2,3-diaminopropionic acid, His, Gln, Gly, Arg, Cit, Nva, D-Asn or a chemical bond;

XX5 represents Thr, NMeSer, Gly, Ala, Hyp, D-Ala, D-Thr, D-Pro or a chemical bond;


XX6 represents Phe, Phe(2F), Phe(3F), Phe(4F), Phe(4Cl), αMePhe , Phe(2Me), Phe(3Me), Phe(4Me), threo-Ser(3Phenyl), erythro-Ser(3Phenyl) or D-Phe;

XX8 represents Leu or Ala(cPr);

XX9 represents Arg, Arg(Me) or D-Arg; and,

XX10 represents Trp.

6. The metastatin derivative according to claim 1, which is represented by formula:



(wherein:

XX0 represents formyl, a C_{1-6} alkanoyl or glycoloyl;

XX2 represents D-Tyr or a chemical bond;

XX3 represents Aze(2), Hyp, Gly, Aib, Leu, Lys, Glu, His(3Me), Tyr(PO_3H_2), Hyp(Bzl), cisHyp or Pro(4F);

XX4 represents Asn or 2-amino-3-ureidopropionic acid;

XX5 represents Ser, Thr or Ala;

XX6 represents Phe, Cha, Phe(2F), Phe(3F), Phe(4F), Phe(4Cl), Phe(2Me), Phe(3Me), Phe(4Me), threo-Ser(3Phenyl) or erythro-Ser(3Phenyl);

AzaGly represents azaglycine;

XX8 represents Leu or Ala(cPr);

XX9 represents Arg or Arg(Me); and,

XX10 represents Phe or Trp); or a salt thereof.

7. Ac-D-Tyr-D-Trp-Asn-Thr-Phe-AzaGly-Leu-D-Arg-Trp- NH_2 (Compound No. 708),

Ac-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp- NH_2 (Compound No. 709),

Decanoyl-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp- NH_2 (Compound No. 710),

Ac-p-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 712),
Ac-Acp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 713),
Ac-D-Tyr-D-Trp-Asp(NHPen)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 714),
Ac-D-Tyr-D-Trp-Asp(NHcPr)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 715),
Ac-D-Tyr-D-Trp-Asp(NHBzl)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 716),
Ac-D-Tyr-D-Trp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 717),
Ac-D-Tyr-D-Pya(4)-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 718),
Ac-D-Tyr-D-Trp-Asn-D-Pro-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 719),
Ac-D-Tyr-Aze(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 720),
Ac-D-Tyr-Pic(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 721),
Ac-D-Tyr-Pic(3)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 722),
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 723),
Ac-D-Tyr-Thz-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 724),
Ac-D-Tyr-NMeAla-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 725),
Ac-D-Tyr-Gly-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 726),
Ac-D-Tyr-Aib-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 727),
Ac-D-Tyr-Abz(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 728),
Ac-D-Tyr-Aze(3)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 730),
Ac-D-Tyr-Sar-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 731),
Ac-D-Tyr-D-NMeAla-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 732),
Ac-D-Tyr-Izc-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 734),
Ac-D-Tyr-D-Asp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 735),

Ac-D-Tyr-D-Dap-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 736),
Ac-D-Tyr-D-Ser-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 737),
Ac-D-Tyr-D-Gln-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 738),
Ac-D-Tyr-D-His-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 739),
Ac-D-Tyr-D-Trp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 740),
Ac-D-Tyr-Ala-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 742),
Ac-D-Tyr-Leu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 743),
Ac-D-Tyr-Ser-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 744),
Ac-D-Tyr-Lys-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 745),
Ac-D-Tyr-Glu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 746),
Ac-D-Tyr-β-Ala-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 747),
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 748),
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 749),
Ac-D-Tyr-D-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 750),
Ac-D-Tyr-Lys-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 754),
Ac-D-Tyr-Glu-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 755),
Ac-D-Tyr-Lys-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 756),
Ac-D-Tyr-Glu-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 757),
Ac-D-Tyr-Lys-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 758),
Ac-D-Tyr-Glu-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 759),
Ac-D-Tyr-Pzc(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.

- 760),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 763),
Ac-D-Tyr-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 764),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 765),
Ac-D-Tyr-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 766),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 767),
Ac-D-Tyr-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 768),
Ac-D-Tyr-Gly-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 769),
Ac-D-Tyr-Aib-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 770),
Ac-D-Tyr-Orn-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 771),
Ac-D-Tyr-Thr-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 772),
Ac-D-Tyr-His(3Me)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 773),
Ac-D-Tyr-DL-Ala(Pip)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 774),
Ac-D-Tyr-Tyr(PO₃H₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 775),
Glycoloyl-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 776),
Ac-D-Tyr-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 777),
Ac-D-Tyr-Pro(4NH₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 780),
Ac-D-Tyr-Hyp(Bzl)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 781),
Ac-D-Tyr-D-NMePhe-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 782),
Ac-D-Tyr-Gly-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.

783),
Ac-D-Tyr-Aib-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 784),
Ac-D-Tyr-Gly-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 785),
Ac-D-Tyr-Aib-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 786),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 787),
Ac-D-Tyr-Glu-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 788),
Ac-D-Tyr-Lys-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 789),
Ac-D-Tyr-Gly-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 790),
Ac-D-Tyr-Aib-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 791),
Ac-D-Tyr-Hyp-Asn-Thr-D-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 794),
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg-Trp-NH₂ (Compound No. 797),
Ac-D-Tyr-Hyp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 800),
4-[Bis-(2-Pyridylmethyl)aminomethyl]benzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 801),
Ac-D-Tyr-Hyp-Asn-NMeSer-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 809),
Ac-D-Tyr-Hyp-Asn-Hyp-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 810),
Ac-D-Tyr-Hyp-Asn-Gly-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 813),
Ac-D-Tyr-Hyp-Asn-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 814),
Ac-D-Tyr-Hyp-Asn-D-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 815),
Ac-D-Tyr-Hyp-His-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 816),
Ac-D-Tyr-Hyp-Gln-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 843),
Ac-D-Tyr-Hyp-D-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 844),
Ac-D-Tyr-Hyp-Cit-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 845),

Ac-D-Tyr-Hyp-Asn-D-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 846),
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Ala(cPr)-Arg(Me)-Trp-NH₂ (Compound No. 856),
4-Ureidobenzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 860),
Ac-D-Tyr-Hyp-Arg-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 861),
Ac-D-Tyr-Hyp-Gly-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 862),
Ac-D-Tyr-Hyp-Dap-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 863),
Ac-D-Tyr-Hyp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 864),
Ac-D-Tyr-Hyp-Asn-Thr- α MePhe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 868),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(2Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 870),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(3Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 872),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 874),
Ac-D-Tyr-Hyp-Asn-Thr-threo-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 877),
Ac-D-Tyr-Hyp-Asn-Thr-erythro-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 882),
Ac-D-Tyr-Hyp-Nva-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 886),
Ac-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 887),
3-(p-Hydroxyphenyl)propionyl-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 888),
pGlu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 889),
Ac-D-Tyr-cisHyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 896),
Ac-D-Tyr-Pro(4F)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 897),
Ac-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 899), or
a salt thereof.

8. Ac-D-Tyr-D-Trp-Asp(NHPen)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 714),
Ac-D-Tyr-D-Trp-Asp(NHcPr)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound

No. 715),

Ac-D-Tyr-D-Trp-Asp(NHBzl)-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 716),

Ac-D-Tyr-D-Trp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 717),

Ac-D-Tyr-D-Pya(4)-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 718),

Ac-D-Tyr-Aze(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 720),

Ac-D-Tyr-Pic(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 721),

Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 723),

Ac-D-Tyr-Thz-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 724),

Ac-D-Tyr-Gly-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 726)

Ac-D-Tyr-Aib-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 727),

Ac-D-Tyr-D-NMeAla-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 732),

Ac-D-Tyr-D-Gln-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 738),

Ac-D-Tyr-D-His-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 739),

Ac-D-Tyr-D-Trp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 740),

Ac-D-Tyr-Ala-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 742),

Ac-D-Tyr-Leu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 743),

Ac-D-Tyr-Ser-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 744),

Ac-D-Tyr-Lys-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 745),

Ac-D-Tyr-Glu-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 746),

Ac-D-Tyr-D-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 748),

Ac-D-Tyr-D-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 749),

Ac-D-Tyr-D-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 750),

Ac-D-Tyr-Lys-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No.

754),
Ac-D-Tyr-Glu-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 755),
Ac-D-Tyr-Lys-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 756),
Ac-D-Tyr-Glu-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 757),
Ac-D-Tyr-Lys-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 758),
Ac-D-Tyr-Glu-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 759),
Ac-D-Tyr-Pzc(2)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 760),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 763),
Ac-D-Tyr-Trp-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 764),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 765),
Ac-D-Tyr-Trp-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 766),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 767),
Ac-D-Tyr-Trp-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 768),
Ac-D-Tyr-Gly-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 769),
Ac-D-Tyr-Aib-Asn-Thr-Phe(4Cl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 770),
Ac-D-Tyr-Orn-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 771),
Ac-D-Tyr-Thr-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 772),
Ac-D-Tyr-His(3Me)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 773),
Ac-D-Tyr-Tyr(PO₃H₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 775),

- Glycoloyl-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 776),
- Ac-D-Tyr-Pro(4NH₂)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 780),
- Ac-D-Tyr-Hyp(Bzl)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 781),
- Ac-D-Tyr-D-NMePhe-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 782),
- Ac-D-Tyr-Gly-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 783),
- Ac-D-Tyr-Aib-Asn-Thr-Phe(2F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 784),
- Ac-D-Tyr-Gly-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 785),
- Ac-D-Tyr-Aib-Asn-Thr-Phe(3F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 786),
- Ac-D-Tyr-Hyp-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 787),
- Ac-D-Tyr-Glu-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 788),
- Ac-D-Tyr-Lys-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 789),
- Ac-D-Tyr-Gly-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 790),
- Ac-D-Tyr-Aib-Asn-Thr-Phe(4F)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 791),
- Ac-D-Tyr-Hyp-Asn-Thr-D-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 794),
- Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg-Trp-NH₂ (Compound No. 797),
- Ac-D-Tyr-Hyp-Alb-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 800),
- 4-[Bis-(2-Pyridylmethyl)aminomethyl]benzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 801),
- Ac-D-Tyr-Hyp-Asn-NMeSer-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 809),
- Ac-D-Tyr-Hyp-Asn-Hyp-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 810),

Ac-D-Tyr-Hyp-Asn-Gly-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 813),
Ac-D-Tyr-Hyp-Asn-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 814),
Ac-D-Tyr-Hyp-Asn-D-Ala-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 815),
Ac-D-Tyr-Hyp-His-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 816),
Ac-D-Tyr-Hyp-Gln-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 843),
Ac-D-Tyr-Hyp-D-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 844),
Ac-D-Tyr-Hyp-Cit-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 845),
Ac-D-Tyr-Hyp-Asn-D-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 846),
Ac-D-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Ala(cPr)-Arg(Me)-Trp-NH₂ (Compound No. 856),
4-Ureidobenzoyl-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 860),
Ac-D-Tyr-Hyp-Arg-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 861),
Ac-D-Tyr-Hyp-Gly-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 862),
Ac-D-Tyr-Hyp-Dap-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 863),
Ac-D-Tyr-Hyp-Dab-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 864),
Ac-D-Tyr-Hyp-Asn-Thr- α MePhe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 868),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(2Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 870),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(3Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 872),
Ac-D-Tyr-Hyp-Asn-Thr-Phe(4Me)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 874),
Ac-D-Tyr-Hyp-Asn-Thr-threo-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 877),
Ac-D-Tyr-Hyp-Asn-Thr-erythro-Ser(3Phenyl)-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 882),
Ac-D-Tyr-Hyp-Nva-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 886),
3-(p-Hydroxyphenyl)propionyl-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 888),
Ac-D-Tyr-cisHyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 896),

Ac-D-Tyr-Pro(4F)-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 897),

Ac-Tyr-Hyp-Asn-Thr-Phe-AzaGly-Leu-Arg(Me)-Trp-NH₂ (Compound No. 899), or a salt thereof.

9. A prodrug of the metastin derivative according to claim 1 or a salt thereof.

10. A prodrug of the metastin derivative according to claim 7 or a salt thereof.

11. A medicament comprising the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

12. The medicament according to claim 11, which is an agent for suppressing cancer metastasis or an agent for suppressing cancer growth.

13. The medicament according to claim 11, which is an agent for preventing or treating cancer.

14. The medicament according to claim 11, which is an agent for controlling placental function.

15. The medicament according to claim 11, which is an agent for preventing or treating choriocarcinoma, hydatid mole, invasive mole, miscarriage, fetal hypoplasia, abnormal glucose metabolism, abnormal lipid metabolism or induction of delivery.

16. The medicament according to claim 11, which is an agent for improving gonadal function.

17. The medicament according to claim 11, which is an agent for preventing or treating hormone-dependent cancer, infertility, endometriosis, early puberty or myoma of the uterus.

18. The medicament according to claim 11, which is an agent for inducing or stimulating ovulation.

19. The medicament according to claim 11, which is a gonadotropic hormone secretagogue agent or a sex hormone secretagogue agent.

20. The medicament according to claim 11, which is an agent for preventing or treating Alzheimer's disease, Autism or moderate cognitive impairment.

21. A method for suppressing cancer metastasis or cancer growth, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

22. A method for preventing or treating cancer, which comprises administering to a mammal an effective dose of the metastin derivative according to

any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

23. A method for controlling placental function, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

24. A method for preventing or treating choriocarcinoma, hydatid mole, invasive mole, miscarriage, fetal hypoplasia, abnormal glucose metabolism, abnormal lipid metabolism or induction of delivery, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

25. A method for improving gonadal function, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

26. A method for preventing or treating hormone-dependent cancer, infertility, endometriosis, early puberty or myoma of the uterus, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

27. A method for inducing or stimulating ovulation, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

28. A method for promoting gonadotropic hormone secretion or promoting sex hormone secretion, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

29. A method for preventing or treating Alzheimer's disease, Autism or moderate cognitive impairment, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

30. The medicament according to claim 11, which is a down-regulating agent of gonadotropic hormone or sex hormone.

31. The agent according to claim 11, which is a down-regulating agent for human OT7T175 (metastin receptor) protein consisting of the amino acid sequence represented by SEQ ID NO: 9.

32. The medicament according to claim 30 or 31, which is an agent for preventing or treating hormone-dependent cancer.

33. A method for down-regulating gonadotropic hormone or sex hormone,

which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

34. A method for down-regulating human OT7T175 (metastin receptor) protein consisting of the amino acid sequence represented by SEQ ID NO: 9, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

35. A method for preventing or treating hormone-dependent cancer, which comprises administering to a mammal an effective dose of the metastin derivative according to any one of claims 1 through 8, or a salt thereof, or a prodrug thereof.

[SEQUENCE LISTING]

<110> Takeda Pharmaceutical Company Limited

<120> Metastin Derivatives And Use Thereof

<130> PCT06-0113

<150> JP2005-370388

<151> 2005-12-22

<150> JP2006-275843

<151> 2006-10-06

<160> 22

<210> 1

<211> 54

<212> PRT

<213> Homo sapiens

<400> 1

Gly Thr Ser Leu Ser Pro Pro Pro Glu Ser Ser Gly Ser Arg Gln Gln

1 5 10 15

Pro Gly Leu Ser Ala Pro His Ser Arg Gln Ile Pro Ala Pro Gln Gly

20 25 30

Ala Val Leu Val Gln Arg Glu Lys Asp Leu Pro Asn Tyr Asn Trp Asn

35 40 45

Ser Phe Gly Leu Arg Phe

50

<210> 2

<211> 162

<212> DNA

<213> Homo sapiens

<400> 2

```

ggtacttctc tgtctccgcc gccggaatct tctggttctc gtcagcagcc gggctctgtct    60
gctccgcact ctcgtcagat cccggctccg caggggtgctg ttctggttca gcgtgaaaaa   120
gacctgccga actacaactg gaactctttc ggtctgcgtt tc                        162

```

<210> 3

<211> 152

<212> PRT

<213> Mus musculus

<400> 3

```

Met Tyr Leu Arg Phe Gly Val Asp Val Cys Ser Leu Ser Pro Trp Lys
      5              10              15
Glu Thr Val Asp Leu Pro Leu Pro Pro Arg Met Ile Ser Met Ala Ser
      20              25              30
Trp Gln Leu Leu Leu Leu Leu Cys Val Ala Thr Tyr Gly Glu Pro Leu
      35              40              45
Ala Lys Val Ala Pro Gly Ser Thr Gly Gln Gln Ser Gly Pro Gln Glu
      50              55              60
Leu Val Asn Ala Trp Glu Lys Glu Ser Arg Tyr Ala Glu Ser Lys Pro
      65              70              75              80
Gly Ser Ala Gly Leu Arg Ala Arg Arg Ser Ser Pro Cys Pro Pro Val
      85              90              95
Glu Gly Pro Ala Gly Arg Gln Arg Pro Leu Cys Ala Ser Arg Ser Arg
      100             105             110
Leu Ile Pro Ala Pro Arg Gly Ala Val Leu Val Gln Arg Glu Lys Asp

```

115 120 125
 Leu Ser Thr Tyr Asn Trp Asn Ser Phe Gly Leu Arg Tyr Gly Arg Arg
 130 135 140
 Gln Ala Ala Arg Ala Ala Arg Gly
 145 150

<210> 4

<211> 456

<212> DNA

<213> Mus musculus

<400> 4

atgtatctga gatttggcgt tgatgtctgc agcctgagtc cctggaagga gactgtagac 60
 ctgcccttc ctcccagaat gatctcaatg gcttcttggc agctgctgct tctcctctgt 120
 gtcgccacct atggggagcc gctggcaaaa gtgaagcctg gatccacagg ccagcagtc 180
 ggaccccagg aactcgtaa tgcctgggaa aaggaatcgc ggtatgcaga gagcaagcct 240
 gggctctgcag ggctgcgcgc tcgtaggtcg tcgccatgcc cgccggttga gggccccgcg 300
 gggcgccagc ggccctgtg tgcctcccgc agtcgcctga tccctgcgcc ccgcggagcg 360
 gtgctggtgc agcgggagaa ggacctgtcc acctacaact ggaactcctt cggcctgcgc 420
 tacggcagga ggcaggcggc gcgggcagca cggggc 456

<210> 5

<211> 156

<212> PRT

<213> Mus musculus

<400> 5

Met Tyr Leu Arg Phe Gly Val Asp Val Cys Ser Leu Ser Pro Trp Lys
 5 10 15
 Glu Thr Val Asp Leu Pro Leu Pro Pro Arg Met Ile Ser Met Ala Ser
 20 25 30

Trp Gln Leu Leu Leu Leu Cys Val Ala Thr Tyr Gly Glu Pro Leu
 35 40 45

Ala Lys Val Ala Pro Leu Val Lys Pro Gly Ser Thr Gly Gln Gln Ser
 50 55 60

Gly Pro Gln Glu Leu Val Asn Ala Trp Glu Lys Glu Ser Arg Tyr Ala
 65 70 75 80

Glu Ser Lys Pro Gly Ser Ala Gly Leu Arg Ala Arg Arg Ser Ser Pro
 85 90 95

Cys Pro Pro Val Glu Gly Pro Ala Gly Arg Gln Arg Pro Leu Cys Ala
 100 105 110

Ser Arg Ser Arg Leu Ile Pro Ala Pro Arg Gly Ala Val Leu Val Gln
 115 120 125

Arg Glu Lys Asp Leu Ser Thr Tyr Asn Trp Asn Ser Phe Gly Leu Arg
 130 135 140

Tyr Gly Arg Arg Gln Ala Ala Arg Ala Ala Arg Gly
 145 150 155

<210> 6

<211> 468

<212> DNA

<213> Mus musculus

<400> 6

atgtatctga gatttggcgt tgatgtctgc agcctgagtc cctggaagga gactgtagac 60
 ctgcccccttc ctcccagaat gatctcaatg gcttcttgge agctgctgct tctcctctgt 120
 gtcgccacct atggggagcc gctggcaaaa gtggcacctt tggtaagcc tggatccaca 180
 ggccagcagt ccggacccca ggaactcggt aatgcctggg aaaaggaatc gcggtatgca 240
 gagagcaagc ctgggtctgc agggctgcgc gctcgtaggt cgtcgccatg cccgccggtt 300
 gagggccccc cggggcgcca gcggcccctg tgtgcctccc gcagtcgctt gatecctgcg 360
 ccccgcgag cggtgctggt gcagcgggag aaggacctgt ccacctacaa ctggaactcc 420
 ttcggcctgc gctacggcag gaggcaggcg gcgcgggcag cacggggc 468

<210> 7

<211> 130

<212> PRT

<213> Rattus sp.

<400> 7

Met Thr Ser Leu Ala Ser Trp Gln Leu Leu Leu Leu Cys Val Ala

5

10

15

Ser Phe Gly Glu Pro Leu Ala Lys Met Ala Pro Val Val Asn Pro Glu

20

25

30

Pro Thr Gly Gln Gln Ser Gly Pro Gln Glu Leu Val Asn Ala Trp Gln

35

40

45

Lys Gly Pro Arg Tyr Ala Glu Ser Lys Pro Gly Ala Ala Gly Leu Arg

50

55

60

Ala Arg Arg Thr Ser Pro Cys Pro Pro Val Glu Asn Pro Thr Gly His

65

70

75

80

Gln Arg Pro Pro Cys Ala Thr Arg Ser Arg Leu Ile Pro Ala Pro Arg

85

90

95

Gly Ser Val Leu Val Gln Arg Glu Lys Asp Met Ser Ala Tyr Asn Trp

100

105

110

Asn Ser Phe Gly Leu Arg Tyr Gly Arg Arg Gln Val Ala Arg Ala Ala

115

120

125

Arg Gly

130

<210> 8

<211> 390

<212> DNA

<213> Rattus sp.

atgacctcgc	tggcttcttg	gcagctgctg	cttctcctcl	gtgtggcctc	ttttggggag	60
ccactggcaa	aaatggcacc	tgtggtgaac	cctgaacca	caggccaaca	gtccggaccc	120
caggaactcg	ttaatgcctg	gcaaaagggc	ccgcggtatg	cagagagcaa	gcctggggct	180
gcaggactgc	gcgctgcg	aacatcgcca	tgcccgccg	tggagaacc	cacggggcac	240
cagcgcccc	cgtgtgccac	ccgcagtcgc	ctgatccctg	cgccccgcg	atcgggtctg	300
gtgcagcgcg	agaaggacat	gtcagcctac	aactggaact	cctttggcct	gcgctacggc	360
aggaggcagg	tggcgcgggc	ggcacggggc				390

<213> Homo sapiens

Met	His	Thr	Val	Ala	Thr	Ser	Gly	Pro	Asn	Ala	Ser	Trp	Gly	Ala	Pro
				5					10					15	
Ala	Asn	Ala	Ser	Gly	Cys	Pro	Gly	Cys	Gly	Ala	Asn	Ala	Ser	Asp	Gly
				20				25						30	
Pro	Val	Pro	Ser	Pro	Arg	Ala	Val	Asp	Ala	Trp	Leu	Val	Pro	Leu	Phe
				35				40					45		
Phe	Ala	Ala	Leu	Met	Leu	Leu	Gly	Leu	Val	Gly	Asn	Ser	Leu	Val	Ile
				50				55				60			
Tyr	Val	Ile	Cys	Arg	His	Lys	Pro	Met	Arg	Thr	Val	Thr	Asn	Phe	Tyr
				65				70				75			80
Ile	Ala	Asn	Leu	Ala	Ala	Thr	Asp	Val	Thr	Phe	Leu	Leu	Cys	Cys	Val
				85						90				95	
Pro	Phe	Thr	Ala	Leu	Leu	Tyr	Pro	Leu	Pro	Gly	Trp	Val	Leu	Gly	Asp
				100				105					110		
Phe	Met	Cys	Lys	Phe	Val	Asn	Tyr	Ile	Gln	Gln	Val	Ser	Val	Gln	Ala
				115				120				125			

Thr Cys Ala Thr Leu Thr Ala Met Ser Val Asp Arg Trp Tyr Val Thr
 130 135 140
 Val Phe Pro Leu Arg Ala Leu His Arg Arg Thr Pro Arg Leu Ala Leu
 145 150 155 160
 Ala Val Ser Leu Ser Ile Trp Val Gly Ser Ala Ala Val Ser Ala Pro
 165 170 175
 Val Leu Ala Leu His Arg Leu Ser Pro Gly Pro Arg Ala Tyr Cys Ser
 180 185 190
 Glu Ala Phe Pro Ser Arg Ala Leu Glu Arg Ala Phe Ala Leu Tyr Asn
 195 200 205
 Leu Leu Ala Leu Tyr Leu Leu Pro Leu Leu Ala Thr Cys Ala Cys Tyr
 210 215 220
 Ala Ala Met Leu Arg His Leu Gly Arg Val Ala Val Arg Pro Ala Pro
 225 230 235 240
 Ala Asp Ser Ala Leu Gln Gly Gln Val Leu Ala Glu Arg Ala Gly Ala
 245 250 255
 Val Arg Ala Lys Val Ser Arg Leu Val Ala Ala Val Val Leu Leu Phe
 260 265 270
 Ala Ala Cys Trp Gly Pro Ile Gln Leu Phe Leu Val Leu Gln Ala Leu
 275 280 285
 Gly Pro Ala Gly Ser Trp His Pro Arg Ser Tyr Ala Ala Tyr Ala Leu
 290 295 300
 Lys Thr Trp Ala His Cys Met Ser Tyr Ser Asn Ser Ala Leu Asn Pro
 305 310 315 320
 Leu Leu Tyr Ala Phe Leu Gly Ser His Phe Arg Gln Ala Phe Arg Arg
 325 330 335
 Val Cys Pro Cys Ala Pro Arg Arg Pro Arg Arg Pro Arg Arg Pro Gly
 340 345 350
 Pro Ser Asp Pro Ala Ala Pro His Ala Glu Leu His Arg Leu Gly Ser
 355 360 365
 His Pro Ala Pro Ala Arg Ala Gln Lys Pro Gly Ser Ser Gly Leu Ala

370

375

380

Ala Arg Gly Leu Cys Val Leu Gly Glu Asp Asn Ala Pro Leu

385

390

395

<210> 10

<211> 1194

<212> DNA

<213> Homo sapiens

<400> 10

```

atgcacaccg tggctacgtc cggacccaac gcgtcctggg gggcaccggc caacgcctcc 60
ggctgcccgg gctgtggcgc caacgcctcg gacggcccag tcccttcgcc gcgggcccgtg 120
gacgcctggc tcgtgccgct cttcttcgcg gcgctgatgc tgctgggcct ggtggggaac 180
tcgtggta tctacgtcat ctgccccac aagccgatgc ggaccgtgac caacttctac 240
atgcccaacc tggcggccac ggacgtgacc ttctcctgt gctgcgtccc cttcacggcc 300
ctgctgtacc cgctgcccgg ctgggtgctg ggcgaactca tgtgcaagtt cgtcaactac 360
atccagcagg tctcgggtgca ggccacgtgt gccactctga ccgccatgag tgtggaccgc 420
tggtacgtga cgggtgtccc gttgcgcgcc ctgcaccgcc gcacgccccg cctggcgctg 480
gctgtcagcc tcagcatctg ggtaggctct gcggcggtgt ctgcgccggt gctcgccctg 540
caccgcctgt caccggggcc gcgcgcctac tgcagtgagg ccttccccag ccgcgcctg 600
gagcgcgcct tcgcaactgta caacctgctg gcgctgtacc tgctgccgct gctcgccacc 660
tgcgccctgt atgcggccat gctgcgccac ctgggcccgg tcgccgtgcg ccccgcgccc 720
gccgatagcg ccttcgaggg gcaggtgctg gcagagcgcg caggcgccgt gcgggccaag 780
gtctcgcgcc tgggtggcgc cgtggtcctg ctcllcgccg cctgctgggg ccccatccag 840
ctgttcctgg tgctgcaggc gctgggcccc gcgggctcct ggcacccacg cagctacgcc 900
gcctacgcgc ttaagacctg ggctcactgc atgtcctaca gcaactccgc gctgaacccg 960
ctgctctacg ccttcctggg ctgcacttc cgacaggcct tccgccggt ctgcccctgc 1020
gcgccgcgcc gccccgcgc cccccgcgg cccggaccct cggacccgc agccccacac 1080
gcggagctgc accgcctggg gtcccacccg gccccgccca gggcgagaa gccagggagc 1140
agtgggctgg ccgcgcgcgg gctgtgcgtc ctgggggagg acaacgcccc tctc 1194

```

<210> 11

<211> 396

<212> PRT

<213> Rattus sp.

<400> 11

Met Ala Ala Glu Ala Thr Leu Gly Pro Asn Val Ser Trp Trp Ala Pro

5

10

15

Ser Asn Ala Ser Gly Cys Pro Gly Cys Gly Val Asn Ala Ser Asp Gly

20

25

30

Pro Gly Ser Ala Pro Arg Pro Leu Asp Ala Trp Leu Val Pro Leu Phe

35

40

45

Phe Ala Ala Leu Met Leu Leu Gly Leu Val Gly Asn Ser Leu Val Ile

50

55

60

Phe Val Ile Cys Arg His Lys His Met Gln Thr Val Thr Asn Phe Tyr

65

70

75

80

Ile Ala Asn Leu Ala Ala Thr Asp Val Thr Phe Leu Leu Cys Cys Val

85

90

95

Pro Phe Thr Ala Leu Leu Tyr Pro Leu Pro Thr Trp Val Leu Gly Asp

100

105

110

Phe Met Cys Lys Phe Val Asn Tyr Ile Gln Gln Val Ser Val Gln Ala

115

120

125

Thr Cys Ala Thr Leu Thr Ala Met Ser Val Asp Arg Trp Tyr Val Thr

130

135

140

Val Phe Pro Leu Arg Ala Leu His Arg Arg Thr Pro Arg Leu Ala Leu

145

150

155

160

Thr Val Ser Leu Ser Ile Trp Val Gly Ser Ala Ala Val Ser Ala Pro

165

170

175

Val Leu Ala Leu His Arg Leu Ser Pro Gly Pro His Thr Tyr Cys Ser

180

185

190

Glu Ala Phe Pro Ser Arg Ala Leu Glu Arg Ala Phe Ala Leu Tyr Asn

195 200 205
 Leu Leu Ala Leu Tyr Leu Leu Pro Leu Leu Ala Thr Cys Ala Cys Tyr
 210 215 220
 Gly Ala Met Leu Arg His Leu Gly Arg Ala Ala Val Arg Pro Ala Pro
 225 230 235 240
 Thr Asp Gly Ala Leu Gln Gly Gln Leu Leu Ala Gln Arg Ala Gly Ala
 245 250 255
 Val Arg Thr Lys Val Ser Arg Leu Val Ala Ala Val Val Leu Leu Phe
 260 265 270
 Ala Ala Cys Trp Gly Pro Ile Gln Leu Phe Leu Val Leu Gln Ala Leu
 275 280 285
 Gly Pro Ser Gly Ala Trp His Pro Arg Ser Tyr Ala Ala Tyr Ala Leu
 290 295 300
 Lys Ile Trp Ala His Cys Met Ser Tyr Ser Asn Ser Ala Leu Asn Pro
 305 310 315 320
 Leu Leu Tyr Ala Phe Leu Gly Ser His Phe Arg Gln Ala Phe Cys Arg
 325 330 335
 Val Cys Pro Cys Gly Pro Gln Arg Gln Arg Arg Pro His Ala Ser Ala
 340 345 350
 His Ser Asp Arg Ala Ala Pro His Ser Val Pro His Ser Arg Ala Ala
 355 360 365
 His Pro Val Arg Val Arg Thr Pro Glu Pro Gly Asn Pro Val Val Arg
 370 375 380
 Ser Pro Ser Val Gln Asp Glu His Thr Ala Pro Leu
 385 390 395

<210> 12

<211> 1188

<212> DNA

<213> Rattus sp.

<400> 12

```

atggccgcag aggcgacgtt gggtcggaac gtgagctggt gggctccgtc caacgcttcg    60
ggatgcccg gctgcggtgt caatgcctcg gatggcccag gctccgcgcc aaggcccctg    120
gatgcctggc tgggtcccct gtttttcgct gccctaattg tgctggggct agtcgggaac    180
tcactggtea tcttcgttat ctgccgccac aagcacatgc agaccgtcac caatttctac    240
atcgctaacc tggcggccac agatgicact ttccttctgt gctgcgtacc cttcaccgcg    300
ctcctctatc cgctgcccac ctgggtgctg ggagacttca tgtgcaaatt cgtcaactac    360
atccagcagg tctcggtgca agccacatgt gccactttga cagccatgag tgtggaccgc    420
tggtacgtga ctgtgttccc gctgcgtgca cttcaccgcc gcaactccgc cctggccclg    480
actgtcagcc ttagcatctg ggtgggttcc gcagctgttt ccgccccggt gctggctctg    540
caccgcctgt cggccgggcc tcacacctac tgcagtgagg cgtttcccag ccgtgccttg    600
gagcgcgctt tcgcgtcta caacctgctg gccctatacc tgctgccgt gctcgccacc    660
tgcgctgct acggtgccat gctgcgccac ctgggccgcg ccgctgtacg cccgcacccc    720
actgatggcg ccctgcaggg gcagctgcta gcacagcgcg ctggagcagt gcgcaccaag    780
gtctcccggc tgggtggccg tgctgtcctg ctcttcgccg cctgctgggg ccgatccag    840
ctgttcttgg tgcttcaagc cctggggccc tggggggcct ggcaccctcg aagctatgcc    900
gcctacgcgc tcaagatctg ggtcactgc atgtcctaca gcaattctgc gctcaaccgc    960
ctgctctatg ccttcttggg ttcccacttc agacaggcct tctgccggt gtgccctgc   1020
ggcccgaac gccagcgtcg gcccacgcg tcagcgcact cggaccgagc cgcaccccat   1080
agtgtgccgc acagccgggc tgcgcaccct gtccgggtca ggacccccga gcctgggaac   1140
cctgtggtgc gctgcacctc tgttcaggat gaacacactg cccactc                   1188

```

<210> 13

<211> 396

<212> PRT

<213> Mus musculus

<400> 13

Met Ala Thr Glu Ala Thr Leu Ala Pro Asn Val Thr Trp Trp Ala Pro

1

5

10

15

Ser Asn Ala Ser Gly Cys Pro Gly Cys Gly Val Asn Ala Ser Asp Asp

20 25 30
 Pro Gly Ser Ala Pro Arg Pro Leu Asp Ala Trp Leu Val Pro Leu Phe
 35 40 45
 Phe Ala Thr Leu Met Leu Leu Gly Leu Val Gly Asn Ser Leu Val Ile
 50 55 60
 Tyr Val Ile Cys Arg His Lys His Met Gln Thr Val Thr Asn Phe Tyr
 65 70 75 80
 Ile Ala Asn Leu Ala Ala Thr Asp Val Thr Phe Leu Leu Cys Cys Val
 85 90 95
 Pro Phe Thr Ala Leu Leu Tyr Pro Leu Pro Ala Trp Val Leu Gly Asp
 100 105 110
 Phe Met Cys Lys Phe Val Asn Tyr Ile Gln Gln Val Ser Val Gln Ala
 115 120 125
 Thr Cys Ala Thr Leu Thr Ala Met Ser Val Asp Arg Trp Tyr Val Thr
 130 135 140
 Val Phe Pro Leu Arg Ala Leu His Arg Arg Thr Pro Arg Leu Ala Leu
 145 150 155 160
 Ala Val Ser Leu Ser Ile Trp Val Gly Ser Ala Ala Val Ser Ala Pro
 165 170 175
 Val Leu Ala Leu His Arg Leu Ser Pro Gly Pro Arg Thr Tyr Cys Ser
 180 185 190
 Glu Ala Phe Pro Ser Arg Ala Leu Glu Arg Ala Phe Ala Leu Tyr Asn
 195 200 205
 Leu Leu Ala Leu Tyr Leu Leu Pro Leu Leu Ala Thr Cys Ala Cys Tyr
 210 215 220
 Gly Ala Met Leu Arg His Leu Gly Arg Ala Ala Val Arg Pro Ala Pro
 225 230 235 240
 Thr Asp Gly Ala Leu Gln Gly Gln Leu Leu Ala Gln Arg Ala Gly Ala
 245 250 255
 Val Arg Thr Lys Val Ser Arg Leu Val Ala Ala Val Val Leu Leu Phe
 260 265 270

Ala Ala Cys Trp Gly Pro Ile Gln Leu Phe Leu Val Leu Gln Ala Leu
 275 280 285

Gly Pro Ser Gly Ala Trp His Pro Arg Ser Tyr Ala Ala Tyr Ala Val
 290 295 300

Lys Ile Trp Ala His Cys Met Ser Tyr Ser Asn Ser Ala Leu Asn Pro
 305 310 315 320

Leu Leu Tyr Ala Phe Leu Gly Ser His Phe Arg Gln Ala Phe Cys Arg
 325 330 335

Val Cys Pro Cys Cys Arg Gln Arg Gln Arg Arg Pro His Thr Ser Ala
 340 345 350

His Ser Asp Arg Ala Ala Thr His Thr Val Pro His Ser Arg Ala Ala
 355 360 365

His Pro Val Arg Ile Arg Ser Pro Glu Pro Gly Asn Pro Val Val Arg
 370 375 380

Ser Pro Cys Ala Gln Ser Glu Arg Thr Ala Ser Leu
 385 390 395

<210> 14

<211> 1188

<212> DNA

<213> Mus musculus

<400> 14

atggccaccg aggcgacatt ggctcccaat gtgacctggt gggtccgctc caacgcttca 60
 ggatgcccag gctgcggtgt caacgcctcg gatgaccag gctctgcgcc aaggcccctg 120
 gatgcctggc tggttcccct gtttttcgct aactcatgt tgcttgggct ggtcggaac 180
 tcattggtca tctacgttat ctgccgccac aagcacatgc agacagttac caacttctac 240
 atcgctaacc tggctgccac agacgtcact ttctactgt gctgcgtgcc cttcaccgca 300
 ctctctacc cgctgcccgc ctgggtgctg ggagacttca tgtgcaaatt cgtaactac 360
 atccagcagg tctcggtgca agccacatgt gccactctga cggccatgag tgtggaccgc 420
 tggatatgtga ctgtgtccc gctgcgtgca cttcaccgcc gactccgcg cctggccctg 480

gctgtcagcc tcagcatctg ggtgggggtca gcagctgtgt ccgccccggt gctggccctg 540
 caccgcctgt cgccagggcc tcgcacctac tgcagcgagg cgtttcccag ccgcgcctg 600
 gagcgcgctt tcgcgtctta caacctgctg gctctatatc tgctgcccgt gctcgccacc 660
 tgcgcctgct acggcgccat gctgcgccac ctgggcccgt cggtgttacg ccccgacccc 720
 actgacggcg ccctgcaggg acagctgcta gcacagcgcg ccggagcagt gcgcaccaag 780
 gtctcccggc tgggtggccgc tgcgtcctg ctcttcgccc cctgctgggg ccgatccag 840
 ctgttcctgg tgcttcaagc cctgggcccc tcgggggcct ggcaccctcg aagctatgcc 900
 gcctacgcgg tcaagatctg ggctcactgc atgtcctaca gcaactcggc gctcaatccg 960
 ctgctctatg ccttcctggg ttcacacttc agacaggcct tctgcccggt gtgcccctgc 1020
 tgccggcaac gccagcgccg gcccacacg tcagcgcaact cggaccgagc tgcaactcac 1080
 actgtgccgc acagccgtgc tgcgcacct gtgcggatca ggagcccgga gcctgggaac 1140
 cctgtggtgc gctcgccctg cgctcagagt gaacgcactg cctcactc 1188

<210> 15

<211> 15

<212> PRT

<213> Artificial

<220>

<223> the C-terminus of the polypeptide is amide (-CONH₂) form

<400> 15

Lys Asp Leu Pro Asn Tyr Asn Trp Asn Ser Phe Gly Leu Arg Phe

1

5

10

15

<210> 16

<211> 10

<212> PRT

<213> Artificial

<220>

<223> the C-terminus of the polypeptide is amide (-CONH₂) form

<400> 16

Tyr Asn Trp Asn Ser Phe Gly Leu Arg Phe

1 5 10

<210> 17

<211> 9

<212> PRT

<213> Artificial

<220>

<223> the C-terminus of the polypeptide is amide (-CONH₂) form

<400> 17

Asn Trp Asn Ser Phe Gly Leu Arg Phe

1 5 9

<210> 18

<211> 8

<212> PRT

<213> Artificial

<220>

<223> the C-terminus of the polypeptide is amide (-CONH₂) form

<400> 18

Trp Asn Ser Phe Gly Leu Arg Phe

1 5 8

<210> 19

<211> 45

<212> DNA

<213> Homo sapiens

<400> 19

aaggacctgc cgaactacaa ctggaactcc ttcggcctgc gcttc 45

<210> 20

<211> 30

<212> DNA

<213> Homo sapiens

<400> 20

tacaactgga actccttcgg cctgcgcttc 30

<210> 21

<211> 27

<212> DNA

<213> Homo sapiens

<400> 21

aactggaact ccttcggcct gcgcttc 27

<210> 22

<211> 24

<212> DNA

<213> Homo sapiens

<400> 22

tggaactcct tcggcctgcg cttc 24

INTERNATIONAL SEARCH REPORT

International application No

PCT/JP2006/326176

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07K14/47 A61K38/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1 577 323 A (TAKEDA PHARMACEUTICAL COMPANY [JP]) 21 September 2005 (2005-09-21) See for instance: compounds 242, 277, 322, 385 and 386, and the claims.	1, 2, 11-35
A	WO 2004/060264 A (TAKEDA CHEMICAL INDUSTRIES LTD [JP] TAKEDA PHARMACEUTICAL COMPANY [JP]) 22 July 2004 (2004-07-22) sequences 3, 4	1-35
X, P	WO 2006/001499 A (TAKEDA PHARMACEUTICAL COMPANY [JP]; KITADA CHIEKO [JP]; ASAMI TAIJI [J]) 5 January 2006 (2006-01-05) claims	1-22, 30-32
	----- -/--	

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

11 April 2007

Date of mailing of the international search report

27/04/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Vogt, Titus

INTERNATIONAL SEARCH REPORT

International application No
PCT/JP2006/326176

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2004/106289 A (COMBIO AS [DK]; BONDEBJERG JON [DK]; FUGLSANG HENRIK [DK]; NAERUM LARS) 9 December 2004 (2004-12-09) -----	
A	EP 1 604 682 A (TAKEDA CHEMICAL INDUSTRIES LTD [JP]) 14 December 2005 (2005-12-14) -----	

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box II.1

Although claims 21-29, 33-35 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box II.2

Claims Nos.: 9,10

Present claims 9 and 10 relate to a prodrug of the metastatin derivatives of claims 1 and 7, respectively. The expression "prodrug" indicates a given desired property or effect, namely a compound that can be converted into a drug.

However, the description does not provide support and disclosure in the sense of Article 6 and 5 PCT for any such prodrugs and there is no common general knowledge of this kind available to the person skilled in the art. The lack of support and disclosure is to such an extent that a search for the subject matter of claims 9 and 10 is not possible.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.5), should the problems which led to the Article 17(2) declaration be overcome.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/JP2006/326176

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 9, 10, 21-29, 33-35
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 21-29, 33-35 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 9, 10
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/JP2006/326176

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1577323	A	21-09-2005	AU 2003292696 A1	10-08-2004
			CA 2511858 A1	29-07-2004
			CN 1761680 A	19-04-2006
			WO 2004063221 A1	29-07-2004
			KR 20050090420 A	13-09-2005
			US 2006241051 A1	26-10-2006
WO 2004060264	A	22-07-2004	CA 2512553 A1	22-07-2004
			CN 1756765 A	05-04-2006
			EP 1581557 A2	05-10-2005
			JP 2006517922 T	03-08-2006
			US 2004142875 A1	22-07-2004
WO 2006001499	A	05-01-2006	AR 049938 A1	13-09-2006
			AU 2005257484 A1	05-01-2006
			CA 2571420 A1	05-01-2006
			EP 1758932 A2	07-03-2007
WO 2004106289	A	09-12-2004	EP 1633702 A1	15-03-2006
EP 1604682	A	14-12-2005	CA 2518541 A1	23-09-2004
			WO 2004080479 A1	23-09-2004
			US 2006287227 A1	21-12-2006